Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Guide to the Experimental Plots - 1913



Full Table of Content

Hoos Field - Leguminous Plants

Rothamsted Research

Rothamsted Research (1914) *Hoos Field - Leguminous Plants*; Guide To The Experimental Plots - 1913, pp 41 - 41 - **DOI:** https://doi.org/10.23637/ERADOC-1-121

HOOS FIELD

41

HOOS FIELD—LEGUMINOUS PLOTS

1848-9 ONWARDS

The small plots (see Plan on page 40) represent portions of the original plots on which attempts have been made to grow leguminous plants continuously since 1848. Various combinations of mineral manures have been used up till 1898, but after the first few years very small crops have been grown, and the clovers in particular generally fail. After fallowing in 1903 to clean the plots, they were resown as before in 1904.

The remainder of the area was formerly occupied by similar small plots of the same leguminous plants. These were ploughed up in 1898, and five crops of wheat were taken without manure in order to test the amount of nitrogen accumulated by the leguminous crop and left in the soil.

In 1904 black tartarian oats were sown, and in the oats, lucerne, red clover, and alsike clover were sown on three strips; a fourth strip, fallowed in 1904, was sown with vetches in October of that year, as shown in the Plan on page 40. The new plots run across the old ones at right angles. The following table shows the crop obtained in 1905 and each year since to 1912 inclusive.

Table XIX.—Produce, Hoos Field Leguminous Land.

	Season 1905.	Season 1906.	Season 1907.	Season 1908.	Season 1909.	Season 1910.	Season 1911.	Season 1912.	
Lucerne .	Cwt. 38·1	Cwt. 55.2	Cwt. 90.6	Cwt. 83.9	Cwt. 15.3	Cwt. 53.3	Cwt. 56.9	Oats Bush.	Straw Cwt. 29.5
Red Clover Alsike Clover	. 47.2	Barley Straw bush. cwt. 36.2 25.6	∫ 67·5 27·8	***	2.4	60.4	23·0 35·7	37·2 29·1	23.5
Vetches .	. 45.8	22.3	24.2	12.2	19.6		8.9	cwt. 5°6	

Dates of sowing leguminous seeds :-

HOOS FIELD—POTATO PLOTS

RESIDUE OF MANURES

On ten plots potatoes were grown with various manures for 26 years (1876-1901). In 1902 the manuring was discontinued and barley sown. Table XX. shows the yields obtained.

Digitized by Microsult &