

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Guide to the Experimental Plots - 1913

[Full Table of Content](#)



Agdell Field - Crops Grown in Rotation

Rothamsted Research

Rothamsted Research (1914) *Agdell Field - Crops Grown in Rotation* ; Guide To The Experimental Plots - 1913, pp 7 - 11 - DOI: <https://doi.org/10.23637/ERADOC-1-121>

THE ROTHAMSTED SOIL

The Rothamsted soil was described by Lawes in 1847, as follows:—
“The soil upon which my experiments were tried consists of rather a heavy loam resting upon chalk, capable of producing good wheat when well manured; not sufficiently heavy for beans, but too heavy for good turnips or barley. The average produce of wheat in the neighbourhood is said to be less than 22 bushels per acre, wheat being grown once in five years. The rent varies from 20s. to 26s. per acre, tithe free.”

It is fairly uniform in the different fields, and consists essentially of a heavy loam containing little coarse sand or grit, but a considerable amount of fine sand and silt, and a large body of clay. In consequence, the soil has to be worked with care, becoming very sticky and drying to impracticable clods if moved when wet. It “runs together” if heavy rain falls after a tilth has been established, and then dries with a hard, unkindly surface, these difficulties being much exaggerated on the plots which have been farmed for a long time without any supply of organic matter in the manures.

The most notable feature in the Rothamsted soil is the amount of calcium carbonate in the surface layer; analyses of the earliest samples available (1856) show more than 5 per cent. in the surface soil of Broadbalk field. The subsoil below the depth of 9 inches contains little or no calcium carbonate, and this fact, together with the varying proportion in the surface soil, indicate that the original soil was almost devoid of calcium carbonate, and that the quantity now found in the surface soil has all been applied artificially.


AGDELL FIELD

Crops grown in Rotation.

The Agdell field, which was put under experiment in the year 1848, differs from the other Rothamsted fields in that it is farmed on a four-course rotation of Swedes, barley, clover (or beans) or fallow, and wheat, instead of growing one crop continuously. It is divided into three main plots, one of which (O) has received no manure, the second (M) mineral manures only, and the third (C) a complete manure, containing the same minerals, but also nitrogen in the form of rape cake and ammonium-salts. The manures are applied to the Swedes only, the other three crops of each course being grown without manure. Each of the three plots is further subdivided:—Half the plots carry clover or beans as the third crop of the course, and half the plots are bare fallow. This shows the effect of introducing the leguminous crop into the rotation, as compared with the bare fallow.

On this field the long-continued cropping without manure on O affects the successive crops in the rotation very differently. The Swede crop is least capable of growing in the impoverished soil, and is reduced to less than a ton per acre; the clover and barley crops are also small, but the deep-rooted wheat crop is comparatively little affected, and yielded as much as 24.2 bushels per acre in 1911, the sixty-fourth year without any

A.—Plan of the Plots in Agdell Field on which Experiments have been made on Four-Course Rotation, 1848 and onwards.



Total area of ploughed land, about 3 acres. Area of each of the 6 divisions, $\frac{1}{2}$ acre.
 The 2 lower divisions, Unmanured continuously (Plots 5 and 6).
 The 2 middle divisions, Mineral Manure, for the Roots, each Course (Plots 3 and 4).
 The 2 upper divisions, Mineral and Nitrogenous Manure, for the Roots, each Course (Plots 1 and 2).
 The 3 left-hand divisions, Clover (or Beans), 3rd year each Course.
 The 3 right-hand divisions, Fallow, 3rd year each Course.
 The double lines indicate division paths between plot and plot.

Digitized by Microsoft

CROPS IN ROTATION

manure. With minerals, but without nitrogen, the Swedes continue to give a fair crop; the barley and wheat are better than on the unmanured plot, while the clover grows almost as freely as on the completely manured plot.

TABLE I.—*Effect of Manure on Crops grown in rotation, Agdell Field. Average produce per acre over the seven last Courses, 1884-1911.*

	O.	M.	C.
	Unmanured. 5 and 6.	Mineral Manure. 3 and 4.	Complete Manure. 1 and 2.
Roots (Swedes) Cwt.	15·0	201·4	373·6
Barley Grain Bush.	14·5	19·4	28·0
Barley Straw Cwt.	10·9	12·8	18·6
Clover Hay * Cwt.	9·9	42·3	39·9
Bean Corn † Bush.	15·8	23·2	19·5
Bean Straw † Cwt.	8·7	16·9	11·5
Wheat Grain Bush.	24·8	34·7	35·5
Wheat Straw Cwt.	21·3	32·2	33·0

* Average of 5 courses.

† Average of 2 courses.

TABLE II.—*Crops grown in rotation, Agdell Field. Produce per acre over the last complete Course (16th), 1908-1911.*

Year.	Crop.	O.		M.		C.	
		Unmanured.		Mineral Manure.		Complete Mineral and Nitrogenous Manures.	
		5.	6.	3.	4.	1.	2.
		Fallow.	Beans or Clover.	Fallow.	Beans or Clover.	Fallow.	Beans or Clover.
1908	Roots (Swedes) . Cwt.	21·6	6·4	179·0	235·8	395·4	314·0
1909	Barley Grain . Bush.	11·4	10·0	17·4	22·1	26·8	33·4
	Barley Straw . Cwt.	10·1	11·3	12·7	16·9	18·7	23·8
1910	Clover Hay . . Cwt.	...	17·4	...	64·1	...	76·7
1911	Wheat Grain . Bush.	23·9	24·5	31·9	37·8	33·3	38·0
	Wheat Straw . Cwt.	20·4	21·4	28·6	33·5	29·3	32·5

When the Plots 2 and 4 grow a good crop of clover, the residues of the crop have a very beneficial effect upon the succeeding crops of the rotation, as compared with the crops of Plots 1 and 3, which are bare fallowed; the wheat is increased by something like 15 per cent., the roots (although manured) are slightly better, and the barley, following the roots, still shows the value of the preceding clover crop. No such residue seems to be left behind by the bean crop, whenever that is taken in the rotation instead of clover. On the unmanured Plot 6, only, the clover shows little or no effect on succeeding crops, because there its growth is too small to leave behind any residue of nitrogen.

AGDELL FIELD




FIG. 1.—Effect of Manure upon Crops grown in Rotation. Total Produce. Average of Five Courses (1884-1903). Swedes in 100 cwt.; Barley and Wheat in 1000 lb.; and Clover in 10 cwt.

TABLE III.—Crops grown in rotation, Agdell Field. Effect of the largest Clover or Bean Crop on the following Wheat Crop. Total produce per acre.

	Clover, 1910.	Wheat, 1911.			Beans, 1862.	Wheat, 1863.		
		After Fallow.	After Clover.	Difference due to Clover.		After Fallow.	After Beans.	Difference due to Beans.
O. Unmanured	Cwt. 17.4	Lb. 3876	Lb. 4052	Per cent. +4.5	Lb. 3603	Lb. 7222	Lb. 5281	Per cent. -26.9
M. Mineral Manure	64.1	5338	6292	+17.9	4033	7910	6090	-23.0
C. Complete Manure	76.6	5454	6163	+13.0	5755	8792	7674	-12.7

TABLE IV.—Crops grown in rotation, Agdell Field. Effect of Clover or Beans on the following Wheat Crops. Total produce per acre.

	Clover Crops.*	Wheat.†			Bean Crops.‡	Wheat.§		
		After Fallow.	After Clover.	Difference due to Clover.		After Fallow.	After Beans.	Difference due to Beans.
O. Unmanured	Cwt. 14.0	Lb. 4028	Lb. 3696	Per cent. -8.2	Lb. 1888	Lb. 4907	Lb. 4373	Per cent. -10.9
M. Mineral Manure	47.0	5147	6052	+17.6	2615	5528	5447	-1.5
C. Complete Manure	50.1	5493	6093	+0.9	3177	6092	5929	-2.7

* 7 years (1874, 1882, 1886, 1894, 1902, 1906, and 1910).

† 7 years (1875, 1883, 1887, 1895, 1903, 1907, and 1911).

‡ 8 years (1854, 1858, 1862, 1866, 1870, 1878, 1890, and 1898).

§ 8 years (1855, 1859, 1863, 1867, 1871, 1879, 1891, and 1899).

The diagram, Fig. 2, shows in a graphic form the benefit the whole rotation receives from the growth of clover, even when the root crop receives nitrogenous manures.




FIG. 2.—Comparative Effect of Clover or Bare Fallow on the succeeding Crops in the Rotation. Total Produce—In 1000 lb. for Clover, Wheat, and Barley, and in 100 cwt. for Roots.

BARN FIELD

Mangolds.

The experiments upon mangolds began in 1876, but the land had been receiving similar manures for other root crops since 1856.

The field is divided longitudinally into eight strips running the whole length of the field; each of these strips receives one manure throughout its length; farmyard manure alone on Strip 1, and in combination with superphosphate and sulphate of potash on Strip 2, nothing on Strip 8, superphosphate alone on Strip 5, superphosphate and sulphate of potash