


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Details of the Classical and Long-term Experiments 1968-73

[Full Table of Content](#)



### R/PG/5 Park Grass - Grass

#### Rothamsted Research

Rothamsted Research (1977) *R/PG/5 Park Grass - Grass* ; Details Of The Classical And Long-Term Experiments 1968-73, pp 17 - 18 - DOI: <https://doi.org/10.23637/ERADOC-1-193>

## PARK GRASS

(R/PG/5)

The manuring, liming and general management of this experiment has continued as set out in *Details 1967*, pp 35-39, but the rates of application are given for convenience in metric terms.

### Symbols, materials and rates of application

Manures applied annually except where indicated.

N1, N2, N3:	Sulphate of ammonia to supply 48, 96, 144 kg N
N1 <sup>x</sup> , N2 <sup>x</sup>	Nitrate of soda to supply 48, 96 kg N
P	Powdered superphosphate (approx. 20% P <sub>2</sub> O <sub>5</sub> ) to supply 34 kg P (except plot 20)
K	Sulphate of potash (approx. 50% K <sub>2</sub> O) to supply 224 kg K (except plot 20)
Na	Sulphate of soda (approx. 14% Na) to supply 16 kg Na
Mg	Sulphate of magnesia (approx. 10% Mg) to supply 11 kg Mg
Si	Silicate of soda at 448 kg water soluble powder
FYM	35 t farmyard manure every fourth year (applied autumn 1968 and 1972)
F	Fish meal (about 6.5% N) to supply 63 kg N every fourth year (applied autumn 1970)
P, K, Na, Mg	applied in winter
N1, N2, N1 <sup>x</sup>	applied in one dressing about March
N2 <sup>x</sup>	Half in March and half in April/May
N3	Two-thirds applied in March and one-third in April/May

**NOTES:** Plot 20: In the three years between applications of FYM mineral fertilisers are applied: 30 kg N (as nitrate of soda), 17 kg P (as superphosphate) and 45 kg K (as muriate of potash)

Plots 5-1, 5-2, 6-N were used for microplot experiments during the period (see 'NPK to Old Grass' and 'N levels to Old Grass')

Plot 6-S: Excluded during the period 1965-71 and used for microplot experiments 1967-70 (simulated grazing R/CS/23; received PKNaMg 1965-71 as previous to 1965). Received N1 in 1972 and N1PKNaMg in 1973.

### Liming.

Ground chalk (t CaCO<sub>3</sub>)

Sub-plots	Dec. 1967			Nov. 1971
	a	b	c	a
1	2.00	—	3.14	2.00
2, 3, 4/1	2.00	—	—	2.00
4/2	2.00	1.26	5.65	—
7/8	2.00	—	—	2.00
9	2.00	2.51	4.39	2.00
10	2.00	1.26	5.02	—
11/1	4.00	6.28	5.02	—
11/2	4.00	3.77	5.02	—
13	2.00	—	1.26	2.00
14, 16, 17	2.00	—	—	2.00

	Dec. 1967			Nov. 1971	
	a	b	c		a
18		1.14	—	2.51	1.14
Whole plots					
5/1		6.15			—
5/2		5.53			—
6		7.53			—
1		—			—
15		—			—
18/2		1.14			1.14
19, 20		1.14			—

Soil series. Batcombe series

### References

1. Edwards, C.A., Butler, C.G. & Lofty, J.R. (1976)  
The Invertebrate Fauna of the Park Grass Plots II. Surface fauna.  
*Rothamsted Experimental Station, Report for 1975, Part 2, 63-89.*
2. Edwards, C.A. & Lofty, J.R. (1976)  
The Invertebrate Fauna of the Park Grass plots I. Soil fauna.  
*Rothamsted Experimental Station. Report for 1975, Part 2, 133-54.*
3. Johnston, A.E. (1972)  
Changes in the soil properties caused by the new liming scheme on Park Grass.  
*Rothamsted Experimental Station, Report for 1971, Part 2, 177-80.*
4. Nutman, P.S. & Ross, G.J.S. (1970)  
Rhizobium in the soils of the Rothamsted and Woburn Farms.  
*Rothamsted Experimental Station. Report for 1969, Part 2, 148-67.*
5. Williams, E.D. (1974)  
Changes in yield and botanical composition caused by the new liming scheme on Park Grass.  
*Rothamsted Experimental Station. Report for 1973, Part 2, 67-73.*