


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Details of the Classical and Long-term Experiments 1968-73

[Full Table of Content](#)



R/RN/7 Residual Phosphate - Barley, Potatoes, Swedes

Rothamsted Research

Rothamsted Research (1977) *R/RN/7 Residual Phosphate - Barley, Potatoes, Swedes* ; Details Of The Classical And Long-Term Experiments 1968-73, pp 53 - 54 - DOI:

<https://doi.org/10.23637/ERADOC-1-193>

RESIDUAL PHOSPHATE

ROTHAMSTED, GREAT FIELD IV AND SAWYERS I

(R/RN/7)

The crop rotation and phosphate manuring programmes for the second period 1967-72 (*Details 1967*, p.89) have been followed. In 1973 the rotation was continued for one more year before changes in 1974, and the treatments were slightly altered on metrication in 1973.

NOTE: The potash applied as a basal manure is the sulphate, not the muriate as stated on p. 90 (*Details 1967*).

Treatments (all as granular superphosphate kg P₂O₅/ha)

Symbol	Frequency	Total 1967-72	1973
0	None	—	0
A1	Annual	188*	29*
A2	Annual	377*	57*
A3	Annual	753*	115*
A4	Annual	1130*	172*
T1	1969, 1972	188*	—
T2	1969, 1972	377*	—
R2	1967	377 ^o	344+
R3	1967	753 ^o	688+
R4	1967	1130 ^o	1032+
G1	None since		
S1	1960		

For details of manuring during the first period 1960-65 and the fallow of 1966 see *Details 1967*, pp 88-89.

* applied in seedbed

+ half applied in autumn before ploughing and half in spring before cultivation

o half applied in June and half in September 1966 and worked into fallow

Basal dressings (kg/ha)

Broadcast in spring before sowing or planting as 'Nitro-Chalk' and sulphate of potash; except for barley since 1970 as (25-0-16) combine drilled.

		N	K ₂ O
Barley:	1968-69	50 Great Field	126
		100 Sawyers	126
	1970-	100	63
Potatoes:	1968-69	150	188
	1970-	251	251
Swedes:	1968-73	63	126

Liming

Ground chalk applied before ploughing in autumn to land to be cropped with barley t/ha

1968	None
1969	2.9

	1970-71	3.1
	1972	2.9
	1973	None
Varieties		
Barley:	1968-69 1970-73	Maris Badger Julia (dressed with ethirimol 1972 and 1973.)
Potatoes:	Majestic, chitted seed: 1968 1969-71 1972 1973	Irish A Once grown Scotch Foundation Stock
Swedes:	1968-73	Wilhelms dressed with Gamma-BHC with captan (1970 Green Globe turnips were sown in place of swedes which failed)
Weedkillers		
Barley:	1969-70 1971-72 1973	2,4-D with dichloroprop Ioxynil, bromoxynil with dichloroprop Dicamba, mecoprop with MCPA
Potatoes:	1968	Paraquat with linuron
Swedes:	1972	Paraquat to previous barley stubble
Other chemicals applied		
Potatoes:	1968 1968-	Mancozeb and demeton-S-methyl Sulphuric acid (BOV) to burn off tops
Areas harvested		
	Great Field IV	Sawyers I
Barley and potatoes:	0.00520	0.00572
Swedes:	0.00390	0.00429
Soil series		
Great Field	Batcombe series	
Sawyers I	Batcombe series with sandier variants.	