

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Details of the Classical and Long-term Experiments 1968-73

[Full Table of Content](#)



### R/CS/14 N P K to Old Grass - Grass

#### Rothamsted Research

Rothamsted Research (1977) *R/CS/14 N P K to Old Grass - Grass* ; Details Of The Classical And Long-Term Experiments 1968-73, pp 74 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-193>

## NPK TO OLD GRASS ROTHAMSTED PARK GRASS

(R/CS/14)

This experiment, started in 1965, studies the effects of a range of P and K levels on yields of permanent pasture and is located on Park Grass plots 5/1 and 5/2 which provide sites with little and much P and K respectively.

### Design

On each site a single replicate of 2 x 4 x 4 in 2 blocks of 16 plots each, with 2 x 2 additional plots in each block.

### Treatments

- (1) The sites differ in previous history:
- |            |           |                                                                                |
|------------|-----------|--------------------------------------------------------------------------------|
| Plot 5/1:  | 1856-97   | 96 kg N annually                                                               |
|            | 1898-1964 | Unmanured                                                                      |
| Plots 5/2: | 1856-97   | 96 kg N annually                                                               |
|            | 1898-1964 | Superphosphate and sulphate of potash to supply 34 kg P and 224 kg K annually. |
- (2) On each site all combinations of:
- (a) Nitrogen fertiliser (kg N for each cut)
- |         |      |      |
|---------|------|------|
|         | N1   | N2   |
| 1965    | 37.5 | 75   |
| 1966    | 56.0 | 112  |
| 1967-73 | 33.6 | 67.2 |
- (b) Phosphate (Kg P) annually as superphosphate
- |    |      |
|----|------|
| P0 | None |
| P1 | 16.8 |
| P2 | 33.6 |
| P4 | 67.2 |
- (c) Potassium (kg K) annually as potassium chloride
- |    |      |
|----|------|
| K0 | None |
| K2 | 112  |
| K4 | 224  |
| K8 | 448  |
- (3) Together with extra treatments on each site:  
all combinations of:
- (a) nitrogen fertiliser as 2(a)
- (b) residues of PK fertiliser applied 1965 only
- 33.6 kg P + 56.1 kg K  
33.6 kg P + 336 kg K
- Number of nitrogen applications have been:
- |             |                  |
|-------------|------------------|
| 1965        | 4                |
| 1966        | 3                |
| 1967        | 5/1 - 3; 5/2 - 4 |
| 1968 & 1969 | 3                |

1970 & 1971	2
1972	3
1973	3

**Liming**

Ground chalk applied (t/ha)

1965:	Plot 5/1 – 12.8	plot 5/2 – 11.5
1968:	Plot 5/1 – 6.2	plot 5/2 – 5.5
1970:	Both plots – 8.8	

**Area harvested**      0.00085 – 0.00090

**Soil series**              Batcombe series