

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

The Long Term Experiments

[Full Table of Content](#)



Agdell

Rothamsted Research

Rothamsted Research (2006) *Agdell* ; The Long Term Experiments, pp 39 - 39 - DOI:
<https://doi.org/10.23637/ROTHAMSTED-LONG-TERM-EXPERIMENTS-GUIDE-2006>

AGDELL

This was the only Classical in which crops were originally grown in rotation. From 1848 to 1951, three different manurial combinations (none, PKNaMg and NPKNaMg plus rape cake, castor meal) were applied to the root crops of two four-course rotations. The rotations differed only in their third course - roots, barley, fallow or legume, wheat. There were only six plots and only one course of the rotation was present each year. The root crop was turnips or swedes, the legume clover or beans. From 1920, club-root (*Plasmodiophora brassicae*) became progressively more damaging to the root crop, especially on the NPKNaMg plots as a result of increasing soil acidity. By 1948 the produce was too small to weigh, and the four-course rotation ceased in 1951. Soil acidity was corrected and the plots were then used to evaluate the P and K reserves accumulated up to 1951. During this period the original six plots were halved and two levels of soil organic matter were established by growing leys on one half. Subsequently, the plots were further sub-divided to build up different amounts of P and K in the soil. Crop yields were then related to the reserves of P and K in the soil and the effect of adding fresh P and K. The experiment ended in 1990.