


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Exhaustion Land Experiment

[Full Table of Content](#)




Default Title

Rothamsted Research

Rothamsted Research (1973-1986) *Default Title* ; Collection Of Plans For The Exhaustion Land Experiment, pp 1 - 17

Rothamsted EXHAUSTION LAND Barley. HOOSFIELD 1968 ↑



TREATMENTS per acre
Residuals of manures applied to Potatoes annually 1876-1901

- No manure since 1852
- N 400 lb sulphate of ammonia
- N* 550 lb nitrate of soda
- P 400 lb superphosphate
- K 300 lb sulphate of potash
- 100 lb sulphate of soda
- 100 lb sulphate of magnesia
- D 14 ton Farmyard manure.

} ^K with ^{N, P, Mg}

Variety: Maris Badger
 Sown at 140 lb per acre: 28 Feb 191

Basal manuring per acre
 0.7 cwt N as Nitro-Chalk 21
 combine drilled.


Plot area: 17'6" x 346' = 0.139 acre
Harvested: 9'4" x 346' = 0.074 acre

White concrete post

Note: Plot dimensions revised spring 1968

4
681

Rothamsted EXHAUSTION LAND Barley HOOSFIELD 114th year 1969



TREATMENTS per acre
Residuals of Manures applied to POTATOES
annually 1876 - 1901

- No manure since 1852
- N 400lb sulphate of ammonia
- N^x 550lb nitrate of soda
- P 400lb superphosphate
- K 300lb sulphate of potash } ^Kwith Na, Mg
- 100lb sulphate of soda
- 100lb sulphate of magnesia
- D 14 tons Farmyard manure.

Variety: Maris Badger
sown at 140lb per acre on 10th March.

Basal manuring, per acre
0.7cwt N as 'Nitro-Chalk' 21 combine drilled


Plot area: 17'6" x 346' = 0.139 acre
Harvested: 9'4" x 346' = 0.074 acre

HOOSFIELD 116th year 1971 4

Barley

Rothamsted EXHAUSTION LAND

71/R/EX/4



TREATMENTS per acre (hectare)
Residuals of Manures applied to POTATOES
 annually 1876 - 1901

- No manure since 1852
 - N 400 lb (448 Kg) sulphate of ammonia
 - N^x 550 lb (617 Kg) nitrate of soda
 - P 400 lb (448 Kg) superphosphate
 - K 300 lb (336 Kg) sulphate of potash
 - 100 lb (112 Kg) sulphate of soda
 - 100 lb (112 Kg) sulphate of magnesias
 - D 14 tons (35 tonnes) farmyard manure
- } 'K' with Na, Mg

Basal manuring, per acre (hectare)

0.7 cwt (87.5 Kg) N as 'Nitro-chalk' 2:1 combine drilled

Variety: Julia

Sown at 140 lb per acre (157 Kg/ha) on 25 Feb 1971

Plot area: 17'6" x 346' = 0.139 acre (0.056 ha)

Harvested: 9'4" x 346' = 0.074 acre (0.03 ha)

Note:


Julia barley drilled on North sideland

HOOSEFIELD 117th year 1972 4

Rothamsted EXHAUSTION LAND

Barley

72/R/EX/4



TREATMENTS per acre (hectare)
Residuals of Manures applied to POTATOES
 annually 1876 - 1901

- No manure since 1852
- N^x 400 lb (448 kg) sulphate of ammonia
- N^x 550 lb (617 kg) nitrate of soda
- P 400 lb (448 kg) superphosphate
- K 300 lb (336 kg) sulphate of potash
- 100 lb (112 kg) sulphate of soda } 'K' with Na, Mg
- 100 lb (112 kg) sulphate of magnesias }
- D 14 tons (35 tonnes) farmyard manure

Basal manuring, per acre (hectare)

0.7 cwt (87.5 kg) N as 'Nitro-chalk' 21 combine drille.

Variety: Julia (seed dressed with ethirimol)

Sown at 140 lb per acre (157 kg/ha) on 16th March

Plot area: 17'6" x 346' = 0.139 acre (0.056 ha)


Harvested: 9'4" x 346' = 0.074 acre (0.03 ha)

Note: 10 Sep. 1971: Sprayed with Weedazol (aminotriazole/amm. thiocyanate) Julia barley drilled on North sideland

Rothamsted EXHAUSTION LAND HOOSFIELD 118th year 1973 4

Barley

73/R/EX/4



TREATMENTS per acre (hectare)
Residuals of Manures applied to POTATOES
 annually 1876 - 1901

- N^x No manure since 1852
 - N^x 400 lb (448 kg) sulphate of ammonia
 - P 550 lb (617 kg) nitrate of soda
 - K 400 lb (448 kg) superphosphate
 - 300 lb (336 kg) sulphate of potash
 - 100 lb (112 kg) sulphate of soda
 - 100 lb (112 kg) sulphate of magnesia
 - D 14 tons (35 tonnes) farmyard manure
- } 'K' with Na, Mg

Basal manuring, per acre (hectare)

0.7 cwt (87.5 kg) N as 'Nitro-chalk' 2.5 combine drilled


Variety: Julia (seed dressed with ethirimol)

Sown at 140 lb per acre (157 kg/ha) on 9 March

Plot area: 17'6" x 346" = 0.139 acre (0.056 ha)
 Harvested: 9'4" x 346" = 0.074 acre (0.03 ha)

4

Rothamsted EXHAUSTION LAND 119th year BARLEY HOOSFIELD 1974



TREATMENTS per acre (hectare)
Residuals of Manures applied to POTATOES
 annually 1876-1901

- No manure since 1852
 - N^x 400 lb (448 Kg) sulphate of ammonia
 - N 550 lb (617 Kg) nitrate of soda
 - P 400 lb (448 Kg) superphosphate
 - K 300 lb (336 Kg) sulphate of potash
 - 100 lb (112 Kg) sulphate of soda
 - 100 lb (112 Kg) sulphate of magnesium
 - D 14 tons (35 tonnes) farmyard manure
- } "K" with Na, Mg

Basal manuring per acre (hectare)
 0.7 cwt (87.5 Kg) N as 'Nitro-Chalk' 25 combine drilled

Variety: Julia (seed dressed with ethirimol)
 Sown at 14.0 lb per acre (157 Kg/ha) or

Plot area: 17'6" x 346' = 0.139 acre (0.056 ha)
Harvested: 9'4" x 346' = 0.074 acre (0.03 ha)

Rothamsted EXHAUSTION LAND


76/R/EX/4

21st Year

BARLEY

4

HOOSFIELD 1976



TREATMENTS per hectare (acre)
N to sub-plots - 1st Year, bare followed in 1975 (N rates to rotate every year)
 - 1, 2, 3 None, 48, 96, 144 Kg N as Nitro-Chalk 25 (- 43, 86, 129 lb N)

Basal manuring: None

Variety: Julia, seed dressed with ethirimol sown at 140 lb/acre (157 kg/ha) on 3 Mar.

Residuals of Manures applied to POTATOES

Annually 1876 - 1901

- N No manure since 1852
 x N 400 lb (448 kg/ha) sulphate of ammonia
 P 550 lb (617 kg) nitrate of soda
 K 400 lb (448 kg) superphosphate
 300 lb (336 kg) sulphate of potash } K with Na, Mg
 100 lb (112 kg) sulphate of soda.
 100 lb (112 kg) sulphate of magnesium }
 D 14 tons (35 tonnes) farmyard manure

Plot area: 17'6" x 345' = 0.139 acre (0.056 ha)
Sub-plot 17'6" x 84' = 0.034 acre (0.014 ha)
Harvested: 9'4" x 84' = 0.018 acre (0.0073 ha)

N sequence

1976	77	78	79	80	81
0	3	2	1	0	3
3	1	3	0	2	1


4
HOOSFIELD 1977

BARLEY

122nd Year

Rothamsted EXHAUSTION LAND

77/R/EX/4



TREATMENTS per hectare (acre)
N Co Sub-Plots - 2nd Year, bare followed in 1975 (N rates to rotate every year)
 0, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Chalk 25 (- 43, 86, 129 lb N)

Basal manuring: None

Variety: Julia, seed dressed with ethirimol sown at 140 lb/acre (157 kg/ha) on 9/3/77

Residuals of Manures applied to POTATOES
 Annually 1876-1901

NPK D
 No manure since 1852
 400 lb (448 kg/ha) Sulphate of ammonia
 550 lb (617 kg) Nitrate of soda
 400 lb (448 kg) Superphosphate
 300 lb (336 kg) Sulphate of potash } K with Na, Mg
 100 lb (112 kg) Sulphate of soda
 100 lb (112 kg) Sulphate of magnesia }
 14 tons (35 tonnes) farmyard manure


Plot area: 17'6" x 345' = 0.139 acre (0.056 ha)
Sub-plot 17'6" x 84' = 0.034 acre (0.014 ha)
Harvested: 9'4" x 84' = 0.018 acre (0.0073 ha)

N sequence

1976	77	78	79	80	81
0	3	2	1	0	3
1	0	3	0	2	1
2	1	0	3	1	0
3	2	1	0	3	2

W 151 9/18/77

Rothamsted EXHAUSTION LAND 123rd Year BARLEY HOOSFIELD 1978
 78/R/EX/4 (J.F.J., A.E.J., J.M.)



TREATMENTS per hectare (acre)

N to Sub-Plots - 3rd Year (N rates rotate every year) *

0, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Chalk 25 (- 43, 86, 129 lb N).

Basal manuring: None

Variety: Julia, seed dressed with ethirimol sown at 140 lb/acre (157 kg/ha) on 31 March

Residuals of Manures applied to POTATOES

Annually 1876-1901

- N manure since 1852
- 400 lb (448 kg/ha) sulphate of ammonia
- 550 lb (617 kg) nitrate of soda
- 400 lb (448 kg) superphosphate
- 300 lb (336 kg) sulphate of potash } K with
- 100 lb (112 kg) sulphate of soda. } Na, Mg
- 100 lb (112 kg) sulphate of soda. } Na, Mg
- 14 tons (35 tonnes) farmyard manure

Plot area: 17'6" x 345' = 0.139 acre (0.056 ha)
 Sub-plot 17'6" x 84' = 0.034 acre (0.014 ha)
 Harvested: 9'4" x 84' = 0.018 acre (0.0073 ha)

* N sequence

1976	1977	1978	79	80	81
0	3	2	1	0	3
1	0	3	3	2	0
2	1	0	0	1	2

Rothamsted EXHAUSTION LAND 12 4 Year BARLEY HOOSFIELD 1979

Sponsors: J.F. Jenkin, A.E. Johnston, J. McCuen

TREATMENTS per hectare (acre)

N to Sub-Plots

(N rates rotate every year)
 0, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Chalk
 (- 43, 86, 129 lb N)

Basal manuring: None

Variety: Julia, seed dressed with ethiri^{not}
 sown at 140 lb/acre (157 kg/ha) on 18 April


Residuals of Manures applied to POTATO

Annually 1876 - 1901

- NPK D
 No manure since 1852
 400 lb (448 kg/ha) sulphate of ammonia
 550 lb (617 kg) nitrate of soda
 400 lb (448 kg) superphosphate
 300 lb (336 kg) sulphate of potash } K²O
 100 lb (112 kg) sulphate of soda } Na₂
 100 lb (112 kg) sulphate of magnesia } Mg
 14 tons (35 tonnes) farmyard manure

Plot area: 17'6" x 34'5" = 0.139 acre (0.056)
 sub-plot 17'6" x 8'4" = 0.034 acre (0.014)
 Harvested: 9'4" x 8'4" = 0.018 acre (0.0073)

* N sequence 1976 77 78 79 80 81
 0 1 2 3 1 0 3
 1 3 0 3 3 0 2
 3 3 1 2 0 3 1 2



Rothamsted EXHAUSTION LAND 12.5 Year BARLEY HOOSFIELD 1980 4

Sponsors J.F. Jenkin, A.F. Johnston, J. MacEwen

TREATMENTS per hectare (acre)
N to Sub-Plots (Started 1976)

0, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Chalk 26
(N rates rotate every year)*

Basal manuring: None

Variety: Georgie, seed/dressed with ethirimol
Sown at 140 lb/acre (157 kg/ha) 9 April


Residuals of Manures applied to POTATOES

Annually 1876-1901

No manure since 1852
400 lb (448 kg/ha) Sulphate of ammonia
550 lb (617 kg) Nitrate of soda
400 lb (448 kg) Superphosphate
300 lb (336 kg) Sulphate of potash } K with
100 lb (112 kg) Sulphate of soda } Na, Mg
100 lb (112 kg) Sulphate of magnesia }
14 tons (35 tonnes) farmyard manure

Plot area: 17'6" x 34'5" = 0.139 acre (0.056 ha)
Sub-plot 17'6" x 8'4" = 0.034 acre (0.014 ha)
Harvested: 9'4" x 8'4" = 0.018 acre (0.0073 ha)


* N sequence 1976 77 78 79 80 81
0 3 2 1 0 3
1 0 3 3 0 3
2 3 0 1 2 3



NOTE: Northern sidebands to receive N at 96 kg/ha in 1980

4

81/R/EX/14 ROTHAMSTED EXHAUSTION LAND 126th YEAR BARLEY HOOSFIELD 1981



Sponsors: T. Fenby, A.E. Johnston, J. McEwen

TREATMENTS Per Acre (acre)

N to Sub-Plots (Started 1976)

(N rates rotate every year)

0, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Chalk, 26% N
(-, 43, 86, 129 lb N)

BASAL MANURING: None

VARIETY: Georgie, seed dressed with Ekaminol
Sown at 140 lb/acre (157 kg/ha) on 18 February

RESIDUALS OF MANURES APPLIED TO POTATOES

ANNUALLY 1876 - 1901

-	No Manure since 1852	
N	400 lb/AC (448 kg/ha)	Sulphate of ammonia
*N	550 " " (617 " ")	Nitrate of Soda
P	400 " " (448 " ")	Superphosphate
K	300 " " (336 " ")	Sulphate of potash
	100 " " (112 " ")	Sulphate of Soda
	100 " " (112 " ")	Sulphate of Magnesia
FYM	14 tons (35 tonnes)	Farmyard manure

PLOT AREA: 17'6" x 345' = 0.139 AC (0.056 ha)

Sub-Plot 17'6" x 84' = 0.034 " (0.014 ")

Harvested 9'4" x 84' = 0.018 " (0.0073 ")

T	N Sequence	1980	'81	'82	'83	'84	'85
		0	3	2	1	0	3
		1	0	3	2	1	0
		2	1	0	3	2	1
		3	2	1	0	3	2

82/R/EX/14 ROTHAMSTED EXHAUSTION LAND 12TH YEAR BARLEY HOOSFIELD 1982

Sponsors: J. F. Temlyn, A. E. Johnston, J. McEwen

TREATMENTS Per hectare (acre)

N to Sub-plots (Started 1976)
 (N rates rotate every year)
 -, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-chalk, 26% N
 (-, 43, 86, 129 lb N)

BASAL MANURING: None

VARIETY: Georgie, seed dressed with Ethirimol
 Sown at 140 lb/acre (157 kg/ha) on 30 March

RESIDUALS OF MANURES APPLIED TO POTATOES

ANNUALLY 1876-1901


-	No Manure since 1852	
N	400 lb/AC (448 kg/ha)	Sulphate of ammonia
*N	550 " " (617 " ")	Nitrate of Soda
P	400 " " (448 " ")	Superphosphate
K	300 " " (336 " ")	Sulphate of Potash
	100 " " (112 " ")	Sulphate of Soda
	100 " " (112 " ")	Sulphate of Magnesia
FYM	14 tons (35 tonnes)	Farmyard manure

PLOT AREA: 17'6" x 345' = 0.139 acre (0.056 ha)

Sub-Plot Harvested

17'6" x 84'	= 0.034 "	(0.014 ")
9'4" x 84'	= 0.018 "	(0.0073 ")

+ N Sequence	1980	'81	'82	'83	'84	'85
	0	3	2	1	0	3
	1	0	3	2	1	0
	2	1	0	3	2	1
	3	2	1	0	3	2



83/R/EX/4 ROTHAMSTED EXHAUSTION LAND

128th YEAR

BARLEY HOOSFIELD 1983



Sponsors: J. F. Tempy, A. E. Johnston, J. McEwen

TREATMENTS Per hectare (acre)

N to Sub-plots (Started 1976)

(N rates rotate every year)
 -, 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Cl (K, 26% N)
 (-, 43, 86, 129 lb N)

BASAL MANURING: None

VARIETY: Georgie, seed dressed with Ekurimol,
 Sown at 140 lb/acre (157 kg/ha) on 10/3/83


RESIDUALS OF MANURES APPLIED TO POTATOES

ANNUALLY 1876 - 1901

-	No Manure since 1852	Sulphate of ammonia	} K' with Na, Mg
N	400 lb/AC (448 kg/ha)	Nitrate of Soda	
*N	550 " " (617 " ")	Superphosphate	
P	400 " " (448 " ")	Sulphate of Soda	
K	300 " " (336 " ")	Sulphate of Magnesia	
FYM	14 tons (35 tonnes)	Farmyard manure	

PLOT AREA: 17'6" x 34'5" = 0.139 acre (0.056 ha)
 Sub-plot 17'6" x 8'4" = 0.034 " (0.014 ")
 Harvested 9'4" x 8'4" = 0.018 " (0.0073 ")


+ N Sequence	1980	'81	'82	'83	'84	'85
	-	3	2	1	-	3
	1	-	3	2	1	-
	2	1	-	3	2	1
	3	2	1	-	3	2



4

84/R/Ex/4 ROTHAMSTED EXHAUSTION LAND 129 H YEAR BARLEY HOOSFIELD 1984

Sponsors: J. F. Tembyn, A. E. Johnston, J. McEwen



TREATMENTS Per hectare (acre)

N to Sub-plots (Started 1976)

(N rates rotate every year)
 - 1, 2, 3 None, 48, 96, 144 kg N as Nitro-Clark, 26% N
 (- , 43, 86, 129 lb N)

BASAL MANURING: None

VARIETY: Triumph, seed dressed with Baytan, sown at 140 lb/acre (157 kg/ha) on 10 March

RESIDUALS OF MANURES APPLIED TO POTATOES

ANNUALLY 1876 - 1901

-	No Manure since 1852	
N	400 lb/AL (448 kg/ha)	Sulphate of ammonia
*N	550 " " (617 " ")	Nitrate of Soda
P	400 " " (448 " ")	Superphosphate
K	300 " " (336 " ")	Sulphate of potash
	100 " " (112 " ")	Sulphate of Soda
	100 " " (112 " ")	Sulphate of Magnesia
FYM	14 tons (35 tonnes)	Farmyard manure

PLOT AREA: 17'6" x 345' = 0.139 acre (0.056 ha)


Sub-Plot 17'6" x 84' = 0.034 " (0.014 ")

Harvested 9'4" x 84' = 0.018 " (0.0073 ")

+ N Sequence	1980	181	182	183	184	185	186
		3	2	1	-	3	2
		1	-	3	2	1	-
		2	1	-	3	2	1
		3	2	1	-	3	2

4

86/R/EX/4 ROTHAMSTED EXHAUSTION LAND 131 SE. YEAR S. BARLEY HOOSFIELD 1986



Sponsors: J.F. Jenkyn, A.E. Johnston, J. McEwen

TREATMENTS Per Aa. (Aa)

N to sub-plots (most half plots 07, 04, 06, 08, 10), in 1986. Started on all plots 1976.

1, 1, 2, 3. None, 48, 96, 144 kg N as Nitrate, 33.5% N
(, 43, 86, 129 lb N)

N sequence - to 3 to 2 to 1, rotating every year.

P to sub-plots (East half only plots 01, 03, 05, 07, 09) started 1986

None
P1 100 kg P₂O₅ (annually until index 1 achieved on most impoverished plots)
P2 200 " " (Annually until index 2 achieved)
P3 300 " " (" " 3 " ")

BASAL MANURING (Eastern half plots 01, 03, 05, 07, 09) started 1986

N at 144 kg N ha⁻¹ + K at 100 kg K₂O ha⁻¹

VARIETY Triumph, dressed 'Baytan', followed by fungicide strips as recommended by county miller

RESIDUALS OF MANURES APPLIED TO POTATOES ANNUALLY 1876-1901

No manure since 1852

N	400 lb/ac (448 kg/ha)	Sulphate of Ammonia
*N	550 " " (617 " ")	Nitrate of Soda
P	400 " " (448 " ")	Superphosphate
K	300 " " (336 " ")	Sulphate of Potash
	100 " " (112 " ")	Sulphate of Soda
	100 " " (112 " ")	Sulphate of Magnesia
FYM	14 tons (35 tonnes)	Farmyard manure

} K, with Na, Mg.

PLOT AREA

17'6" x 345'	= 0.139 ac (0.056 ha)
17'6" x 84'	= 0.034 " (0.014 ")
84' x 84'	= 0.018 " (0.0073 ")