

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)

The thumbnail image shows a table titled "WOBURN ARABLE AND LEY ROTATIONS" with columns for "STOCKING FIELD IN 1970", "TREATMENTS", and "CROPPING". The table lists various experimental treatments and their corresponding cropping patterns.

Woburn Ley-arable 1970b

Rothamsted Research


Rothamsted Research (1957-79) *Woburn Ley-arable 1970b* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 15 - 15

2018

WOBURN
ARABLE & LEY ROTATIONS
Potato Treatments Block 4

33rd year

STACKYARD D 1970



TREATMENTS per acre

- Whole plots
 - , d8 None, FYM residues applied 1963
Half plots - Fumigant
 - , F None, chloropicrin at 400lb
1/6 plots - Nitrogen rates
 2,3,4 1.0, 1.5, 2.0 cwt Nas 'Nitro-chalk' 21
1/2 plots - Nematicide
 - , T None, Temik at 100lb of 10% granules.

BASAL MANURING per acre

0.9 cwt P₂O₅, 1.8 cwt K₂O as compound fertiliser (0:14:28)
 336 lb Epsom salts

VARIETY: Maris Piper Planted: 22 Apr.

PLOT AREA 18' x 7' = 0.0029 acre
 harvested one row 2' 4" x 18' = 0.0010 acre