


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



**ROTHAMSTED
RESEARCH**

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)



WOBURN ARABLE AND LEY ROTATIONS

Treatment	1957-58	1958-59	1959-60	1960-61	1961-62	1962-63	1963-64	1964-65	1965-66	1966-67	1967-68	1968-69	1969-70	1970-71	1971-72	1972-73	1973-74	1974-75	1975-76	1976-77	1977-78	1978-79	
1. 1st Year																							
2. 2nd Year																							
3. 3rd Year																							
4. 4th Year																							
5. 5th Year																							
6. 6th Year																							
7. 7th Year																							
8. 8th Year																							
9. 9th Year																							
10. 10th Year																							
11. 11th Year																							
12. 12th Year																							
13. 13th Year																							
14. 14th Year																							
15. 15th Year																							
16. 16th Year																							
17. 17th Year																							
18. 18th Year																							
19. 19th Year																							
20. 20th Year																							

Woburn Ley-arable 1971b

Rothamsted Research


Rothamsted Research (1957-79) *Woburn Ley-arable 1971b* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 17 - 17

WOBURN 3171
71/W/RN/3

ARABLE AND LEY ROTATIONS
Potatoes Block 3

33rd year

STACKYARD D 1971 2018



TREATMENTS per acre (hectare)

Dung residues

- , d6 None, FYM residues - last applied 1966.

Chloropicrin residues

- , f None, 400lb (448kg) Chloropicrin in 1968.

Fresh chloropicrin and temik

- None

F 400lb (448kg) chloropicrin in autumn 1970
+ 100lb (112kg) of 10% granules of temik
in spring 1971.

Planted 30 March.

R resistant to *H. rostochianensis* Maris Piper

S susceptible to *H. rostochianensis* Pentland Crown

BASAL MANURING per acre (hectare)

1752lb (1964kg) compound fertiliser (13:13:20) broadcast 26 Mar.

PLOT AREA

Plots without chloropicrin residues 14' x 30' = 0.0096 acre (0.0039 ha)

Plots with chloropicrin residues 7' x 30' = 0.0048 acre (0.0048 ha)