

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)

WOBURN ARABLE AND LEY MIXTURE
STATIONARY FIELD EXPERIMENT
1957-79

Year	Plot	Area (sq ft)	Yield (lb/acre)	Notes
1957	1	1000	1000	
1958	1	1000	1000	
1959	1	1000	1000	
1960	1	1000	1000	
1961	1	1000	1000	
1962	1	1000	1000	
1963	1	1000	1000	
1964	1	1000	1000	
1965	1	1000	1000	
1966	1	1000	1000	
1967	1	1000	1000	
1968	1	1000	1000	
1969	1	1000	1000	
1970	1	1000	1000	
1971	1	1000	1000	
1972	1	1000	1000	
1973	1	1000	1000	
1974	1	1000	1000	
1975	1	1000	1000	
1976	1	1000	1000	
1977	1	1000	1000	
1978	1	1000	1000	
1979	1	1000	1000	

Woburn Ley-arable 1973c

Rothamsted Research



Rothamsted Research (1957-79) *Woburn Ley-arable 1973c* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 22 - 22

WOBURN 36th year STACKYARD D 1973 201c

→ N

Amended Plan 3d/1/73

ARABLE AND LEY ROTATION
Block 4
1st Test Crop



TREATMENTS per acre (hectare)

F.Y.M. residues

- , d II None, F.Y.M. residues - last applied 1963

chloropicrin residues

(-), (c) None, 400lb (448kg) chloropicrin for potatoes 1970

Fresh Fumigant

- None

F 400lb (448kg) chloropicrin in early spring 1973

+ 6.4lb (67 kg) of aldicarb

VARIETY Maris Piper

Planted: 9 Apr.

BASAL MANURING per acre (hectare)

1730 lb (1939 kg) compound fertiliser (13:13:20)

PLOT AREA

Plots with chloropicrin residues 14' x 30' = 0.0096 acre (0.0089ha)

Plots without chloropicrin residues 14' x 64' 6" = 0.0207 acre (0.0084ha)

Type
Rotation
FYM RES 6
FYM RES 7
FYM RES 13

A P P A