

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)

The thumbnail image shows a table titled "WOBURN ARABLE AND LEY ROTATIONS" with columns for "STOCKING FIELD IN 1957/58", "TREATMENTS", and "PLANTING". The table lists various experimental treatments and their corresponding stocking fields.

Woburn Ley-arable 1973c

Rothamsted Research

Rothamsted Research (1957-79) *Woburn Ley-arable 1973c* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 22 - 22



WOBURN 36th year STACKYARD D 1973 201c

→ N

Amended Plan 3d/1/73

ARABLE AND LEY ROTATION Block 4

1st Test Crop



Type Rotation
FYM RES 6
FYM RES 7
FYM RES 13

TREATMENTS per acre (hectare)

F.Y.M. residues

- , d II None, F.Y.M. residues - last applied 1963

chloropicrin residues

(-), (c) None, 400lb (448kg) chloropicrin for potatoes 1970

Fresh Fumigant

- None

F 400lb (448kg) chloropicrin in early spring 1973

+ 6.4lb (67 kg) of aldicarb

VARIETY Maris Piper

Planted: 9 Apr.

BASAL MANURING per acre (hectare)

1730 lb (1939 kg) compound fertiliser (13:13:20)

PLOT AREA

Plots with chloropicrin residues 14' x 30' = 0.0096 acre (0.0089ha)

Plots without chloropicrin residues 14' x 64' 6" = 0.0207 acre (0.0084ha)