

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)

The thumbnail image shows a table titled "WOBURN ARABLE AND LEY ROTATIONS" with columns for "STOCKING FIELD IN 1957/58" and "ARABLE ROTATIONS". The table contains several rows of data, including crop sequences like "1. 1957-58, 2. 1958-59, 3. 1959-60" and "1. 1957-58, 2. 1958-59, 3. 1959-60, 4. 1960-61".

Woburn Ley-arable 1975c

Rothamsted Research

Rothamsted Research (1957-79) *Woburn Ley-arable 1975c* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 29 - 29

201 C

STACKYARD D 1975

38th year

ARABLE AND LEY ROTATION

Winter Wheat Block 2


2nd Test Crop

FYM RES 74

WOBURN

FYM RES 64

N Sponsors: D.A. Boyd, J.M. Hirst, A.E. Dunston, J.G.M. Jones



Nitrogen

0, 1, 2, 3 None, 0.5, 1.0, 1.5 cut N (63, 126, 189 kg) as Nitro-chalk 2.5

BASAL MANURING per acre (hectare)

260 lb (290 kg) of (0:20:20) C.D.

VARIETY

Cappelle dressed diehldrin

PLOT AREA

14' x 30' = 0.0096 acre (0.0039 ha.)
Harvested: 9'4" x 30' = 0.0064 acre (0.0026 ha.)

TREATMENTS per acre (hectare)

FYM residues

- , d 12 None, FYM residues - last applied 1964

Fungicides - applied for potatoes 1974

(-) None

(F) Residues of 200 lb (224 kg) Telone + 6 lb (6.7 kg) addicarb.

PERM A/L 1