

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)


WOBURN ARABLE AND LEY ROTATIONS

Treatment	Area (Acre)	Soil	Year	Notes
1. 1st Year	1.0	1.0	1957	
2. 2nd Year	1.0	1.0	1958	
3. 3rd Year	1.0	1.0	1959	
4. 4th Year	1.0	1.0	1960	
5. 5th Year	1.0	1.0	1961	
6. 6th Year	1.0	1.0	1962	
7. 7th Year	1.0	1.0	1963	
8. 8th Year	1.0	1.0	1964	
9. 9th Year	1.0	1.0	1965	
10. 10th Year	1.0	1.0	1966	
11. 11th Year	1.0	1.0	1967	
12. 12th Year	1.0	1.0	1968	
13. 13th Year	1.0	1.0	1969	
14. 14th Year	1.0	1.0	1970	
15. 15th Year	1.0	1.0	1971	
16. 16th Year	1.0	1.0	1972	
17. 17th Year	1.0	1.0	1973	
18. 18th Year	1.0	1.0	1974	
19. 19th Year	1.0	1.0	1975	
20. 20th Year	1.0	1.0	1976	

Woburn Ley-arable 1976b

Rothamsted Research

Rothamsted Research (1957-79) *Woburn Ley-arable 1976b* ; Collection Of Plans For The Woburn Ley-Arable Experiment, pp 32 - 32



Nitrogen
 0, 1, 2, 3 None, 0.5, 1.0, 1.5 cat N (65, 126, 189 kg) as Nitro-chalk 25

BASAL MANURING per acre (hectare)
 270 lb (303 kg) of compound fertiliser (0:20:20) C.D.

VARIETY
 Cappelle

Plot AREA
 Sown: 15 Oct at : 190 lb (213 kg)
 14' x 30' = 0.0096 acre (0.0039 ha)
 Harvested 9'4" x 30' = 0.0064 acre (0.0026 ha)

TREATMENTS per acre (hectare)
FYM residues
 -, d 12 None, FYM residues - last applied 1965

FUMIGANTS - applied for potatoes 1975
 None.
 200 lb (22.4 kg) 'Telone' in autumn +
 10 lb (11.2 kg) aldicarb in spring