

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Collection of Plans for the Woburn Ley-arable Experiment

[Full Table of Content](#)



Woburn Ley-arable 1978a

Rothamsted Research

Rothamsted Research (1957-79) *Woburn Ley-arable 1978a ; Collection Of Plans For The Woburn Ley-Arable Experiment*, pp 37 - 37

WOBBURN ARABLE AND LEY ROTATION

BLK 78/W/RN/3
1st year 1978 STACKYARD D 1978 201A
Sponsors: A.E.Johnston, F.G.W.Jones, G.A.Salt

1.	01 d 14 (Ah) B 02	-	03 (C) Lc2 04	d14
05 d 14	(A) F 06	-	07 (Ah) LLc2* 08	d14
" 09	(LA) Lln2 10	d14	" d14 (AL) LLn2* 12	-
13 d 14	(L) Lln2 14	-	15 (SAh) Llc2 16	d14
17	(Ah) O 18	d15	19 (A) O 20	d15
21 d 15	(SA) LLc3 22	-	23 (AC) LLc3* 24	d15
25 d 15	(Ah) Lln3* 26	-	27 (LAh) Lln3 28	d15
29	(C) Lc3 30	d15	31 (L) Lln3 32	d15
600' 6"				
33 d 13	(C) Lc1 34	-	35 (AL) Lln6 36	-
37	(LAh) Lln1 38	d13	39 (A) F 40	-
3. 41	(SA) LLc1 42	d13	43 (L) Lln1 44	d13
45	(Ah) B 46	d13	47 (Ah) Llc6 48	d13
4. 49	Crop Winter Wheat 4' 6" → 64' 6" ←	↑ 2½"	4' 6" → 64' 6" ←	↑ 2½"
5. 65	Block 4 1st Test For Plot treatments	See Separate Plan 201B 64	Crop Barley See Separate plan 201C 80	17' ← 133' 6" → 284'

SYMBOLS
 From 1976 major revisions have been made:-
Previous Rotations: (A), (Ah), (C), (L), (S) not defined on this plan
 From 1976 AH: Barley B, Barley B, Barley B, Oats O.
 A : Fallow F, Fallow F, Oats O.
 LC : 3 years clover/grass ley (no N)
 LN : 3 years all grass ley (with N)
 LLC & LLN 8 year leys treated as above.
 * = ploughed up in less than 8 year leys treated as above.
 Followed by Test Crops Wheat W, Barley B.
 NOTE: All LC, LN, LLC and LLN 2 & 3 failed in 1976 and resown in autumn 1976.
 To all crops: - d in Residues of none visitors/acre (385/ha) FYM
BASAL MANURING per acre (hectare) to treatment crops:
 Barley - Porthos (dressed with ethirimol) 360lb (400kg) compound
 Oats - Manod
 All - Grass ley
 1st year: 75kg (67lb) N as Nitro-chalk 25, 75kg (67lb) P₂O₅ 150kg (134lb) K₂O as (0:14:28) in the seedbed, 75kg (67lb) N, 48kg (43lb) K₂O as (25:0:16) after each cut except the last.
 2nd + years: 75kg (67lb) P₂O₅ 150kg (134lb) K₂O as (0:14:28) in winter,
 75kg (67lb) N, 48kg (43lb) K₂O as (25:0:16) in spring,
 and after each cut except the last.
Clover/grass ley as LN, but omitting N, K per gal as Muriate of potash
 1st year: 75kg (67lb) P₂O₅ 150kg (134lb) K₂O as (0:14:28) in the seedbed, 48kg (43lb) K₂O as Muriate of potash after each cut except the last.
 2nd + years: 75kg (67lb) P₂O₅ 150kg (134lb) K₂O as (0:14:28) in winter, 48kg (43lb) K₂O as Muriate of potash in spring, and after each cut except the last.
 Barley sown at 145lb (162kg)
 Oats sown at 180lb (202kg)

Date: Redrilled 7 Apr
Date: 13 Mar