

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Collection of Plans for the Woburn Organic Manuring Experiment

[Full Table of Content](#)


W/RN/12 - Woburn Organic Manuring 1995 - 1999

Rothamsted Research

Rothamsted Research (1966-1988) *W/RN/12 - Woburn Organic Manuring 1995 - 1999* ; Collection Of Plans For The Woburn Organic Manuring Experiment, pp 30 - 34

WOBURN W. Wheat
 9.5/W/RN/12
 ORGANIC MANURING EXPERIMENT
 31st DEAR STACKYARD 3 1995
 Sponsor: P. R. Poulton

204



TREATMENTS

Residues of organic manures

- (Dg) FYM annually 1965-71 and 1981-85.
 - (St) Straw annually 1965-71 and 1981-85.
 - (Gm) Green manure annually 1965-71 then (Lc8) Ley (clover/grass) 1979-86 (Blks I-III) 1980-87 (Blks II-IV) (Pt) Peat annually 1985-71 then (Lc8) 1979-86 (Blks I-III) 1980-87 (Blks II-IV)
- Residues of Fertilizers
- (Fd) P K Mg Equivalent to (Dg) rate, annually 1965-71 & 1981-86 (Blks I-III) 7 (Blks II-IV)
 - (Fs) K Mg Equivalent to straw with P annually 1965-71 then
 - ½ Dg rate 1981-6 (Blks II-IV) 7 (Blks I-III)
 - (Lc) clover/grass ley 1965-71 then (Lc6) 1981-6 (Blks I-III) 7 (Blks II-IV)
 - (Lm) Grama ley with N - clover/grass ley
- Leys: (Lc) clover/grass ley }
 (Lm) Grama ley with N }

Subplot Residual nitrogen treatments

- (-), (1), (2), (3), (4), (5) 0, 50, 100, 150, 200, 250 Kg N ha⁻¹

BASAL MANURE

- P K None this year
- Mn foliar spray @ SFP in spring
- N at 100 Kg N

VARIETY

Mercia dressed Rappot @ 300 seeds/m² on 7 Oct 1994

SUB PLOT AREA

14' x 30' = 0.039 ha
 Harvested area = 2m x 9.16m = 0.0183 ha

230

32nd year STACKYARD 2 1996
 Sponsor: P. R. Poulton

WOBURN W. Wheat

Block I 6 H. wheat

01	02	03	04	05	06	07	08
(1)(5)	(4)(3)	(4)(4)	(3)(4)	(3)(4)	(5)(3)	(4)(1)	(5)(4)
(Lc)	(Lm)	(Sc)	(Gm)	(Dg)	(Fg)	(Lc)	(Fs)
(4)(1)	(5)(2)	(3)(1)	(3)(5)	(1)(5)	(4)(1)	(5)(2)	(1)(2)
(2)(3)	(1)(1)	(5)(2)	(1)(2)	(1)(2)	(2)(1)	(1)(3)	(3)(1)

Block II 5 H. wheat

09	10	11	12	13	14	15	16
(1)(5)	(2)(1)	(2)(5)	(2)(5)	(1)(5)	(2)(3)	(1)(4)	(5)(3)
(Fd)	(Gm)	(Dg)	(Lc)	(Lc)	(Fs)	(Sc)	(Lc)
(4)(2)	(4)(2)	(1)(4)	(3)(2)	(3)(2)	(5)(1)	(1)(5)	(1)(4)
(1)(3)	(1)(5)	(3)(4)	(1)(1)	(4)(1)	(1)(1)	(2)(3)	(1)(2)

Block III 6 H. wheat

17	18	19	20	21	22	23	24
(3)(1)	(1)(2)	(3)(1)	(1)(3)	(4)(3)	(1)(5)	(3)(4)	(1)(4)
(Sc)	(Fs)	(Lc)	(Lm)	(Pt)	(Lc)	(Dg)	(Lc)
(2)(1)	(1)(4)	(2)(2)	(1)(5)	(1)(1)	(2)(4)	(5)(3)	(3)(2)
(1)(4)	(4)(3)	(5)(1)	(2)(4)	(2)(5)	(1)(3)	(1)(1)	(1)(5)

Block IV 5 H. wheat

25	26	27	28	29	30	31	32
(5)(4)	(1)(4)	(2)(3)	(1)(4)	(1)(2)	(1)(3)	(3)(1)	(5)(1)
(Lm)	(Dg)	(Pt)	(Fs)	(Lc)	(Fd)	(Sc)	(Gm)
(1)(1)	(3)(1)	(4)(5)	(5)(1)	(4)(5)	(5)(2)	(2)(4)	(5)(1)
(3)(2)	(2)(5)	(1)(1)	(2)(3)	(1)(3)	(1)(4)	(1)(5)	(4)(2)

TREATMENTS

Residues of organic manures

- (Dg) F/M annually 1965-71 and 1981-85
 - (St) Straw annually 1965-71 and 1981-85
 - (Gm) Green manure annually 1965-71 then (Lc8) 1y (clover/grass) 1977-86 (Blks I+III) 1980-87 (Blks II+IV)
 - (Pt) Peat annually 1965-71 then (Lc8) 1977-86 (Blks I+III) 1980-87 (Blks II+IV)
- Residues of Fertilizers
- (Fd) P K Mg Equivalent to (Dg) rate, annually 1965-71 & 1981-6 (Blks I+III) 7 (Blks II+IV)
 - (Fs) K Mg Equivalent to straw with P annually 1965-71 then 1/2 Dg rate 1981-6 (Blks II+IV) 7 (Blks I+III)
- Lays: (Lc) clover/grass ley 1965-71 then (Lc6) 1981-6 (Blks I+III) 7 (Blks II+IV) - clover/grass ley
 (Lm) Great ley with N

Subplot Residual nitrogen treatments
 (-) (1)(2)(3)(4)(5) 0, 50, 100, 150, 200, 250 kg N ha⁻¹

BASAL MANURE


- P & K None this year
- Mn foliar spray @ SFP in spring
- N at 100 kg N

VARIETY

Mercia dressed Sibutol @ 375 seeds/m² on 3 Oct 95

SUB PLOT AREA


14' x 36' = 0.039 ha.
 harvested area: 2m x 9.16m = 0.0182 ha



WOBURN
33rd Year STACKYARD B 1997
Sponsor: P. R. Poulton

ORGANIC MANURING EXPERIMENT

W. Rye



TREATMENTS

- Residues of organic manures
 (Dg) FYM annually 1965-71 and 1981-85
 (St) Straw annually 1965-71 and 1981-85
 (Gm) Green manure annually 1965-71 then (Lc8) ley
 (Pt) Peat annually 1965-71 then (Lc8) 1979-86 (Blks I-III) 1980-87 (Blks II > IV)
 Residues of Fertilizers
 (Fd) P K Mg Equivalent to (Dg) rate, annually 1965-71 & 1981-86 (Blks I-III) 7 (Blks II > IV)
 (Fs) K Mg Equivalent to straw with P annually 1965-71 then
 1/2 Dg rate 1981-86 (Blks II > IV) 7 (Blks I-III)
 Lays: (Lc) clover/grass ley 1965-71 then (Lc6) 1981-86 (Blks I-III) 7 (Blks II > IV) -
 (Lm) Grass ley with N - clover/grass ley

Subplot Residual nitrogen treatments last applied 1994
 (-), (1), (2), (3), (4), (5) 0, 50, 100, 150, 200, 250. kg N ha⁻¹

BASAL MANURE
 P & K (0:20:32) @ 500 kg ha⁻¹
 No nitrogen


VARIETY
 Amando dressed Bataun @ 400 seeds/m² on 10 Oct 1996

SUB-PLOT AREA
 14' x 30' = 0.039 ha.
 harvested area = 2m x 9.16m = 0.00183 ha

Straw yields from (Fd) and (Dg) treatments

W. wheat

Sponsor P.R. Poulton



Sponsor P.R. Poulton

