

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Collection of Plans for the Woburn Organic Manuring Experiment

[Full Table of Content](#)



## W/RN/12 - Woburn Organic Manuring 2010 - 2014

### Rothamsted Research

Rothamsted Research (1966-1988) *W/RN/12 - Woburn Organic Manuring 2010 - 2014* ; Collection Of Plans For The Woburn Organic Manuring Experiment, pp 47 - 56

46th year Plan 203

Code: 2010/W/RN/12

Title: Organic Manuring

Field: Stackyard B

**Sponsors:**

A Macdonald K Goulding  
2479 2627

**Treatments:**

Organic Treatments since 2003

- F None
- Dg10 FYM at 10 t/ha annually
- Dg25 FYM at 25 t/ha annually
- St Chopped wheat straw at 7.5 t/ha annually
- CC Cover crop (white mustard) prior to spring sown crop
- Co Compost at 40 t/ha annually
- Lc Permanent grass/clover: Timothy, Fescue & white clover sown @ 30 kg/ha, 4-Oct-2002

**Basal PKS/ha (to be applied early spring)**

Basal P at 97.5 kg TSP (20 kg P) to all treatments except Dg25  
Basal K at 200 kg K<sub>2</sub>SO<sub>4</sub> (83 kg K + 36 kg S) to all treatments except Dg25

**No Nitrogen to W Beans for 2010**

**Nitrogen rotation**

N5>N4>N3>N2>N1>N0>N5

*Residues of previous treatments (see 'Yields' books 1965-2002 for details)*

- (Dg) FYM annually 1965-71, and 1981-85
- (St) Straw annually 1965-71, and 1981-85
- (Gm) Green manure annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)
- (Pt) Peat annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1980-87 (blocks 2 & 4)
- (Fd) PKMg equivalent to (Dg) rate, annually 1965-71, and 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)
- (Fs) KMg equivalent to (St) with P, annually 1965-71, then half (Fd) rate 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)
- (Ln) Grass ley with N 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)
- (Lc) Grass /clover ley 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)


**Crop rotation**

- 2004 S. Barley and Mustard
- 2005 W. Beans
- 2006 W. Wheat
- 2007 F Maize and Mustard
- 2008 W. Rye
- 2009 S. Barley and Mustard
- 2010 W. Beans
- 2011 W. Wheat

**Variety**

W Beans: Wizard

// From 1999 block 3 has a 3.0 m discard at the NE end following severe erosion in winter 1998-99; sub-plots are therefore shorter  
// Paths widened to 1.0 m in 2009, measurements corrected and detailed on layout sheet, including reference posts



47th year  
Plan 203

Code: 2011W/RN/12

Title: Organic Manuring

Field: Stackyard B

**Sponsors:**

A Macdonald  
2479  
K Goulding  
2627

**Treatments:**

Organic Treatments since 2003

F None

Dg10 FYM at 10 t/ha annually

Dg25 FYM at 25 t/ha annually

St Chopped wheat straw at 7.5 t/ha annually

CC Cover crop (white mustard) prior to spring sown crop

Co Compost at 40 t/ha annually

Lc Permanent grass/clover: Timothy, Fescue & white clover sown @ 30 kg/ha, 4-Oct-2002

**Basal PKS/ha (to be applied early spring)**

Basal P at 97.5 kg TSP (20 kg P) to all treatments except Dg25

Basal KS at 200 kg K2SO4 (83 kg K + 36 kg S) to all treatments except Dg25

**Nitrogen to W. Wheat for 2011, kg N/ha, N0-N5.**

0, 50, 100, 150, 200, 250 as Nitrochalk

Split 50 kg N late Feb/early March, remainder mid-April

**Nitrogen rotation**

N5>N4>N3>N2>N1>N0>N5

**Residues of previous treatments (see 'Yields' books 1965-2002 for details)**

(Dg) FYM annually 1965-71, and 1981-85

(St) Straw annually 1965-71, and 1981-85

(Gm) Green manure annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

(Pt) Peat annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

(Fd) PKMg equivalent to (Dg) rate, annually 1965-71, and 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

(Fs) KMg equivalent to (St) with P, annually 1965-71, then half (Fd) rate 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

(Ln) Grass ley with N 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

(Lc) Grass /clover ley 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

**Variety**

W. Wheat: Gallant

**Crop rotation**

2004 S. Barley and Mustard

2005 W. Beans

2006 W. Wheat

2007 F. Maize and Mustard


2008 W. Rye

2009 S. Barley and Mustard

2010 W. Beans

2011 W. Wheat

//Straw yields not required. Straw baled and removed  
// From 1999 block 3 has a 3.0 m discard at the NE end following severe erosion in winter 1998-99; sub-plots are therefore shorter  
// Paths widened to 1.0 m in 2009, measurements corrected and detailed on layout sheet, including reference posts



Code: 2012/W/RN/12  
 Title: Organic Manuring  
 Field: Stackyard B

48th year

Plan 203

**Sponsors:**

A Macdonald  
 2479  
 K Goulding  
 2627

**Treatments:**

Organic Treatments since 2003

F None  
 Dg10 FYM at 10 t/ha annually  
 Dg25 FYM at 25 t/ha annually  
 St Chopped wheat straw at 7.5 t/ha annually  
 CC Cover crop (white mustard) prior to spring sown crop  
 Co Compost at 40 t/ha annually  
 Lc Permanent grass/clover: Timothy, Fescue & white clover sown @ 30 kg/ha, 4-Oct-2002

**Basal PKS/ha (to be applied early spring)**

Basal P at 97.5 kg TSP (20 kg P) to all treatments except Dg25  
 Basal KS at 200 kg K2SO4 (83 kg K + 36 kg S) to all treatments except Dg25

**Nitrogen to F Maize for 2012, kg N/ha, N0-N5.**

0, 50, 100, 150, 200, 250 as Nitrochalk  
 Split 50 kg N to seedbed (except N0); remainder post emergent.

**Nitrogen rotation**

N5>N4>N3>N2>N1>N0>N5

*Residues of previous treatments (see 'Yields' books 1965-2002 for details)*

(Dg) FYM annually 1965-71, and 1981-85  
 (St) Straw annually 1965-71, and 1981-85  
 (Gm) Green manure annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
 (Pt) Peat annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1980-87 (blocks 2 & 4)  
 (Fd) PKMg equivalent to (Dg) rate, annually 1965-71, and 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
 (Fs) KMg equivalent to (St) with P, annually 1965-71, then half (Fd) rate 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
 (Ln) Grass ley with N 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
 (Lc) Grass /clover ley 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)


**Crop rotation**

2005 W. Beans  
 2006 W. Wheat  
 2007 F Maize and Mustard  
 2008 W. Rye  
 2009 S. Barley and Mustard  
 2010 W. Beans  
 2011 W. Wheat  
 2012 F Maize and Mustard

**Variety**

W. Mustard - var Rivona (farm to confirm)  
 F Maize - var Hudson (farm to confirm)

//Straw yields not required. Straw baled and removed  
 // From 1999 block 3 has a 3.0 m discard at the NE end following severe erosion in winter 1998-99; sub-plots are therefore shorter  
 // Paths widened to 1.0 m in 2009, measurements corrected and detailed on layout sheet, including reference posts



Code: 2013/W/RN/12

Title: Organic Manuring

Field: Stackyard B

49th year

Plan 203

**Sponsors:**

A Macdonald  
2479  
K Goulding  
2627

**Treatments:**

Organic Treatments since 2003

F None  
Dg10 FYM at 10 t/ha annually  
Dg25 FYM at 25 t/ha annually  
St Chopped wheat straw at 7.5 t/ha annually  
CC Cover crop (white mustard) prior to spring sown crop  
Co Compost at 40 t/ha annually  
Lc Permanent grass/clover: Timothy, Fescue & white clover sown @ 30 kg/ha, 4-Oct-2002

**Basal PKS/ha (to be applied early spring)**

Basal P at 97.5 kg TSP (20 kg P) to all treatments except Dg25  
Basal KS at 200 kg K<sub>2</sub>SO<sub>4</sub> (83 kg K + 36 kg S) to all treatments except Dg25

**Nitrogen to Rye for 2013, kg N/ha, N0-N5.**  
0, 30, 60, 90, 120, 150 as Nitrochalk

**Nitrogen rotation**  
N5>N4>N3>N2>N1>N0>N5

**Residues of previous treatments (see 'Yields' books 1965-2002 for details)**

(Dg) FYM annually 1965-71, and 1981-85  
(St) Straw annually 1965-71, and 1981-85  
(Gm) Green manure annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Pt) Peat annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Fd) PKMg equivalent to (Dg) rate, annually 1965-71, and 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Fs) KMg equivalent to (St) with P, annually 1965-71, then half (Fd) rate 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Ln) Grass ley with N 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Lc) Grass/clover ley 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

**Crop rotation**

2006 W. Wheat  
2007 F Maize and Mustard  
2008 W. Rye  
2009 S. Barley and Mustard  
2010 W. Beans  
2011 W. Wheat  
2012 F Maize and Mustard  
2013 W. Rye


**Variety**

Rye - What Variety? Is it Matador?

//Straw yields not required. Straw baled and removed  
// From 1999 block 3 has a 3.0 m discard at the NE end following severe erosion in winter 1998-99; sub-plots are therefore shorter  
// Paths widened to 1.0 m in 2009, measurements corrected and detailed on layout sheet, including reference posts



2013/W/RN/12 Stackyard B  
 Organic Manuring Winter Rye 49th year Stackyard B 203



Code: 2014/W/RN/12

Title: Organic Manuring

Field: Stackyard B

50th year

Plan 203

**Sponsors:**

A Macdonald  
2479

K Goulding  
2627

**Treatments:**

Organic Treatments since 2003

F None  
Dg10 FYM at 10 t/ha annually  
Dg25 FYM at 25 t/ha annually  
St Chopped wheat straw at 7.5 t/ha annually  
CC Cover crop (white mustard) prior to spring sown crop  
Co Compost at 40 t/ha annually  
Lc Permanent grass/clover: Timothy, Fescue & white clover sown @ 30 kg/ha, 4-Oct-2002

**Basal PKS/ha (to be applied early spring)**

Basal P at 97.5 kg TSP (20 kg P) to all treatments except Dg25

Basal KS at 200 kg K<sub>2</sub>SO<sub>4</sub> (83 kg K + 36 kg S) to all treatments except Dg25

**Nitrogen to Spring Barley for 2014, kg N/ha, N0-N5.**

0, 35, 70, 105, 140, 175 as Nitrochalk @ 27%N

**Nitrogen rotation**

N5>N4>N3>N2>N1>N0>N5

**Residues of previous treatments (see 'Yields' books 1965-2002 for details)**

(Dg) FYM annually 1965-71, and 1981-85  
(St) Straw annually 1965-71, and 1981-85  
(Gm) Green manure annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Pt) Peat annually 1965-71, then (Lc8) grass/clover ley 1979-86 (blocks 1 & 3), 1980-87 (blocks 2 & 4)  
(Fd) PKMg equivalent to (Dg) rate, annually 1965-71, and 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Fs) KMg equivalent to (St) with P, annually 1965-71, then half (Fd) rate 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Ln) Grass ley with N 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)  
(Lc) Grass /clover ley 1965-71, then (Lc6) grass/clover ley 1981-86 (blocks 1 & 3), 1981-87 (blocks 2 & 4)

**Crop rotation**

2008 W. Rye  
2009 S. Barley and Mustard  
2010 W. Beans  
2011 W. Wheat  
2012 F. Maize and Mustard  
2013 W. Rye  
2014 S. Barley and Mustard  
2015 W. Beans

**Variety**

Mustard - Zlata (or similar) to be sown as early as possible; preferably by mid-August.  
Sp. Barley - Tipple

//Straw yields not required. Straw baled and removed  
// From 1999 block 3 has a 3.0 m discard at the NE end following severe erosion in winter 1998-99; sub-plots are therefore shorter  
// Paths widened to 1.0 m in 2009, measurements corrected and detailed on layout sheet, including reference posts

