


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Rothamsted Report for 1908

[Full Table of Content](#)



Barley Hoos Field 1908 ; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1908

Rothamsted Research

Rothamsted Research (1909) *Barley Hoos Field 1908 ; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1908 ; Rothamsted Report For 1908*, pp 14 - 14 - DOI:

<https://doi.org/10.23637/ERADOC-1-100>

BARLEY. HOOS FIELD, 1908.

(Previous cropping : Potatoes, 1876-1901 ; Barley, 1902 and 1903 ;
Oats, 1904 ; Barley, 1905 and 1906.

(See " Guide," page 40, Table XIX).

Plot.	Manures applied to the Potatoes, 1876-1901. Unmanured since.	Dressed Grain.		Straw.	Total Produce.
		Yield.	Weight per Bushel.		
		Bushels.	lbs.		
1	Unmanured	6.8	54.6	3.4	800
2	Unmanured 1882 to 1901, previously Dung only ...	7.8	54.5	5.5	1097
3	Dung 1883-1901	20.2	55.2	11.4	2484
4	Dung 1883-1901	22.6	55.0	12.7	2771

WHEAT AFTER FALLOW (without manure 1851 and since).

HOOS FIELD, 1908.*

(See " Guide," page 41, Table 20).

Dressed Grain	Yield—7.2 bushels. Weight per bushel—63.6 lbs.
Straw	
Total Produce	1083 lbs.

* The young plants of wheat were much damaged by hares in the spring.