


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Annual Report for 1910 With the Supplement to the Guide to the Experimental Plots Containing Yields per Acre, Etc.



[Full Table of Content](#)

Barley Hoos Field 1910 ; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1910

Rothamsted Research

Rothamsted Research (1911) *Barley Hoos Field 1910 ; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1910 ; Annual Report For 1910 With The Supplement To The Guide To The Experimental Plots Containing Yields Per Acre, Etc.*, pp 13 - 13 - DOI:

<https://doi.org/10.23637/ERADOC-1-102>

BARLEY. HOOS FIELD, 1910.

(Previous Cropping: Potatoes, 1876-1901; Barley, 1902 and 1903;
Oats, 1904; Barley, 1905 and since.)

(See "Guide," page 40, Table XIX).

Plot.	Manures applied to the Potatoes, 1876-1901. Unmanured since.	Dressed Grain.		Straw.	Total Produce.
		Yield.	Weight per Bushel.		
		Bushels.	lbs.	Cwt.	lbs.
1	Unmanured	6.2	53.5	5.0	944
2	Unmanured 1882 to 1901, previously Dung only ...	11.4	53.1	8.3	1620
3	Dung 1883 to 1901	15.5	52.0	17.1	2895
4	Dung 1883 to 1901	15.6	52.2	15.4	2697

WHEAT AFTER FALLOW (without Manure 1851 and since).
HOOS FIELD, 1910.

(See "Guide," page 41, Table XX.)

Dressed Grain	{ Yield—9.3 Bushels. Weight per Bushel—61.2 lbs.
Straw	
Total Produce	1747 lbs.
