Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Annual Report for 1913 With the Supplements to the Guide to the Experimental Plots Containing the Yields per Acre, Etc.



Full Table of Content

Wheat Broadbalk Field 1913; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1913

Rothamsted Research

Rothamsted Research (1914) Wheat Broadbalk Field 1913; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1913; Annual Report For 1913 With The Supplements To The Guide To The Experimental Plots Containing The Yields Per Acre, Etc., pp 27 - 27 - DOI:

https://doi.org/10.23637/ERADOC-1-115

27

WHEAT. BROADBALK FIELD, 1913.

(See "Guide," 1913, page 30, Table XVI.)

Plot.		Dressed Grain.		Straw	Average for 61 years, 1852-1912.	
	Manuring.	Yield per Acre.	Weight per Bushel.	per Acre.	Dressed Grain per Acre.	Straw per Acre.
		Bushels	lb.	cwt.	Bushels.	cwt.
2	Farmyard Manure	25.7	64.9	29.6	35.2	34.8
3 5	Unmanured	5.8	63.3	4.2	12.6	10.3
5	Complete Mineral Manure	9.4	63.8	7.2	14.2	12.1
6	As 5, and single Amm. Salts	1412	64.7	13.8	23.2	21.4
7	As 5, and double Amm. Salts	20.8	65.8	26.7	32.1	32.9
8	As 5, and treble Amm. Salts	28.5	65.8	38.4	36.6	41.1
9	As 5, and single Nitrate Soda	23.3	61.6	24.2	_	_
10	Double Amm. Salts alone	11.3	64.2	11.7	20.0	18.4
11	As 10, and Superphosphate	13.4	63.9	14.6	22.9	22.3
12	As 10, and Super and Sulph. Soda	19.0	65.1	22.7	29.1	28.0
13	As 10, and Super and Sulph. Potash	21.6	65.6	36.3	31.0	31.2
14	As 10, and Super and Sulph. Mag.	19.5	65.4	25.7	28.8	28.0
15	Double Amm. Salts in Autumn, and					
	Minerals	22.4	65.6	22.3	29.9	29.7
16	Double Nitrate and Minerals	23.9	65.2	37.4		_
17	Minerals alone, or double Amm. Salts J	*9.8	*65.0	*9.2	*14.9	*13.0
18)	alone, in alternate years	+21.7	†65.7	+31.1	+29.9	+29.5
19	Rape Cake alone	19.8	65.2	24.8	25.4 (2)	25.7 (2)
20 (1)	As 7, but excluding Superphosphate	11.0	65.5	17.5	_	

^{*} Produce by Minerals.

Note.—As in the previous season (1912), owing to the foulness of the land on the upper half of the field, the produce here recorded was that obtained on the lower half of the field only. (See notes on the crop at p. 7).

WHEAT AFTER FALLOW (without manure 1851 and since). HOOS FIELD, 1913.

(See "Guide," 1913, page 44, Table XXI.)

			1913	Average 57 years 1856-1912
Dressed Grain Straw Total produce	•••	Yield—Bushels per Acre Weight per Bushel cwt. per Acre lb. per Acre	8°3 61°8 6°7 1284	16°0 59°4 13°7 2536

[†] Produce by Ammonium Salts.

⁽¹⁾ Commenced in 1906.

^{(2) 20} years, 1893—1912.