Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Annual Report for 1914 With the Supplements to the Guide to the Experimental Plots Containing the Yields per Acre, Etc.



Full Table of Content

# Wheat Broadbalk Field 1914; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1913 and 1914

#### **Rothamsted Research**

Rothamsted Research (1915) Wheat Broadbalk Field 1914; Wheat After Fallow (Without Manure 1851 and Since) Hoos Field 1913 and 1914; Annual Report For 1914 With The Supplements To The Guide To The Experimental Plots Containing The Yields Per Acre, Etc., pp 36 - 36 - DOI: https://doi.org/10.23637/ERADOC-1-107

### Wheat. Broadbalk Field, 1914. (See "Guide," 1913, page 30, Table XVI.)

|       |                                      | Dressed<br>Grain.     |                          | Straw         | Average<br>for 61 years,<br>1852-1912. |                    |
|-------|--------------------------------------|-----------------------|--------------------------|---------------|--|--------------------|
| Plot. | Manuring.                            | Yield<br>per<br>Acre. | Weight<br>per<br>Bushel. | per<br>Acre.  | Dressed<br>Grain<br>per Acre.          | Straw<br>per Acre. |
|       |                                      | Bushels               |                          | cwt.          | Bushels.                               | cwt.               |
| 2     | Farmyard Manure                      | 30.7                  | 62.2                     | 36.6          | 35.2                                   | 34.8               |
| 3     | Unmanured                            | 5.8                   | 60.6                     | 5.1           | 12.6                                   | 10.3               |
| 5     | Complete Mineral Manure              | 9.8                   | 61.4                     | 7.7           | 14.2                                   | 12.1               |
| 6     | As 5, and single Amm. Salts          | 16.9                  | 62.6                     | 16.0          | 23.2                                   | 21.4               |
| 7     | As 5, and double Amm. Salts          | 23.4                  | 62.2                     | 27.3          | 32.1                                   | 32.9               |
| 8     | As 5, and treble Amm. Salts          | 27.7                  | 62.1                     | 41.9          | 36.6                                   | 41.1               |
| 9     | As 5, and single Nitrate Soda        | 18.7                  | 62.0                     | 19.8          |  |                    |
| 10    | Double Amm. Salts alone              | 12.0                  | 62.1                     | 12.6          | 20.0                                   | 18.4               |
| 11    | As 10, and Superphosphate            | 14.9                  | 61.8                     | 16.1          | 22.9                                   | 22.3               |
| 12    | As 10, and Super. and Sulph. Soda    | 22.7                  | 62.7                     | 22.0          | 29.1                                   | 28.0               |
| 13    | As 10, and Super. and Sulph. Potash  | 19.2                  | 62.4                     | 21.6          | 31.0                                   | 31.5               |
| 14    | As 10, and Super. and Sulph. Mag.    | 14.6                  | 61.9                     | 17.0          | 28.8                                   | 28.0               |
| 15    | Double Amm. Salts in Autumn, and     |                       |                          |               | 20.0                                   |                    |
|       | Minerals                             | 19.5                  | 62.3                     | 22.6          | 29.9                                   | 29.7               |
| 16    | Double Nitrate and Minerals          | 19.2                  | 62.3                     | 26.1          |  |                    |
| 17)   | Minerals alone, or double Amm. Salts | *15.9                 | *62.2                    | *19.6         | *29.9                                  | *29.5              |
| 18 ∫  | alone, in alternate years            | †5.3                  | †62.4                    | †4.2          | †14.9                                  | †13.0              |
| 19    | Rape Cake alone                      | 9.5                   | 62.2                     | 12.0          | 25.4 (2)                               | 25.7 (2)           |
| 1     |                                      |                       |                          | - Compression |  |                    |

<sup>\*</sup> Produce by Ammonium Salts. † Produce by Minerals.

Note.—As in the two previous seasons, (1912 and 1913), owing to the foulness of the land on the upper half of the field, the produce here recorded was that obtained on the lower half of the field only. (See notes on the crop at p. 7.)

## Wheat after Fallow (without Manure 1851 and since). Hoos Field, 1913 and 1914.

(See "Guide," 1913, page 44, Table XXI.)

|               |       |       |   | 1913 | 1914                                   | Average<br>57 years<br>1856-191: |
|---------------|-------|-------|---|------|--|----------------------------------|
| Dressed Grain | • • • | • • • | ( Yield—Bushels per Acre<br>( Weight per Bushel | 8.3  | 14 <sup>.</sup> 9<br>62 <sup>.</sup> 7 | 16.0<br>59.4                     |
| Straw         |       |       | (Weight per Bushel cwt. per Acre                | 6.7  | 11.9                                   | 13.7                             |
| Total produce | • • • |       | lb. per.Acre                                    | 1284 | 2351                                   | 2536                             |

Digitized by Microsoft @

<sup>(1)</sup> Commenced in 1906. (2) 20 years, 1893—1912.