


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



## Report 1915-17 With the Supplement to the Guide to the Experimental Plots Containing the Yields per Acre Etc.



[Full Table of Content](#)

### Table of Results - the Classical Experiments

#### Rothamsted Research

Rothamsted Research (1917) *Table of Results - the Classical Experiments* ; Report 1915-17 With The Supplement To The Guide To The Experimental Plots Containing The Yields Per Acre Etc., pp 51 - 69  
- DOI: <https://doi.org/10.23637/ERADOC-1-108>

DATES OF SOWING AND HARVESTING, 1915

Field.	Crop.	Variety.	Sowing began.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, west	Oats	Grey Winter	Oct. 15, '14	Aug. 6	Aug. 21	Aug. 21	4 qrs.
" " east	Swedes	Monarch	May 22, '15	...	...	...	...
" " "	Brussels Sprouts	Sutton's Matchless	* June 30, '15	...	...	...	...
" " "	Savoys	...	...	...	...	...	...
Little Knott Wood	Oats	Grey Winter	Oct. 20, '14	Aug. 11	Aug. 24	Aug. 24	4½ qrs.
Sawpit	Clover	Red	Apr. 9, '14	{ June 19 Sept. 7	July 2 Sept. 14	July 3 Sept. 17	...
Great Harpenden, west	Wheat	Squareheads Master	Nov. 6, '14	Aug. 17	Aug. 25	Aug. 26	3¼ qrs.
" " east	Barley	Plumage Cross	Mar. 29, '15	Aug. 19	Aug. 26	Aug. 28	5 "
New Zealand	Grass	(Cattle Grazing)	...	...	...	...	...
Stackyard	Barley	Plumage Cross	Mar. 31, '15	Aug. 27	Sept. 6	Sept. 6	4½ qrs.
Long Hoos	"	"	Mar. 22, '15	Aug. 30	Sept. 9	Sept. 13	3½ "
West Barn	Potatoes	{ King Edward Arran Chief, Scottish Farmer	Apr. 4, '15	...	Sept. 21	Sept. 27	5½ tons
Foster's, west	Wheat	Dalhousie	Apr. 16, '15	...	Oct. 18	Oct. 23	...
" " east	Oats	Squareheads Master	Nov. 4, '14	Aug. 23	Aug. 28	Aug. 28	2½ qrs.
Broadbalk	Wheat	Grey Winter	Oct. 17, '14	Aug. 7	Aug. 23	Aug. 23	3½ "
Little Hoos	Mangolds	Squareheads Master	Nov. 3, '14	Aug. 16	Aug. 24	Aug. 25	see p. 60
Hoos	Barley	Yellow Globe	May 15, '15	...	Nov. 15	Nov. 26	" 63
Barnfield	Mangolds	Archer's Stiff Straw	Apr. 3, '15	Aug. 28	Sept. 3	Sept. 4	" 61
Agdell	Wheat	Sutton's Yellow Globe	Apr. 22, '15	...	Oct. 30	Nov. 10	" 56
Greatfield	Grass	Squareheads Master	Nov. 10, '14	Aug. 17	Aug. 26	Aug. 27	" 54
Park	"	(Cattle Grazing)	...	...	...	...	...
" " "	"	...	...	June 21	June 23	June 26	see p. 57
" " "	"	...	...	Sept. 16	Sept. 20	Sept. 21	" "

\* Setting out.

DATES OF SOWING AND HARVESTING, 1916.

Field.	Crop.	Variety.	Sowing began.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, east	Barley	Burton Brewing	Apr. 28	Sept. 4	Sept. 8	Sept. 12	3 qrs.
" " west	Swedes	Magnum Bonum	June 14	...	Jan. 1, '17	Jan. 15, '17	
	Potatoes	King Edward	May 24	...	Oct. 20	Oct. 24	3½ tons
	Savoys	"	June 23*	...	Dec., sold	...	
	Mangolds	Yellow Globe	June 6	...	Dec. 1	Dec. 4	
Little Knott Wood	Barley	Burton Brewing	Apr. 8	Aug. 28	Sept. 14	Sept. 16	3½ qrs.
Sawpit, north	Wheat	White Chaff	Nov. 9, '15	Aug. 24	Sept. 12	Sept. 13	3½ "
" south	Oats	Grey Winter	Oct. 15, '15	Aug. 9	Sept. 13	Sept. 14	
Great Harpenden, east	Wheat	Wilhelmina	Nov. 20, '15	Aug. 21	Sept. 11	Sept. 12	3½ qrs.
" " west	Wheat	Rivetts	Nov. 16, '15	Aug. 21	Sept. 11	Sept. 12	
" " "	Oats	Grey Winter	Oct. 13, '15	Aug. 8	Aug. 24	Sept. 26	4½ "
	Beans	"	Apr. 15	Sept. 27	Oct. 9	...	
	Winter Oats	Grey Winter	Apr. 8	Aug. 12	Aug. 28	...	
	Wheat	Squareheads	Apr. 5	Aug. 26	Sept. 9	...	
	Spring Oats	Master	Apr. 14	Sept. 7	Sept. 14	...	see
New Zealand	Barley	"	Apr. 15	Sept. 6	Sept. 14	...	p. 69
	Potatoes	Plumage	May 30	...	Oct. 27	Oct. 28	
	Mangolds	"	May 31	Nov. 9	...	...	
Stackyard	Oats	Yellow Globe	Oct. 10, '15	Aug. 7	Aug. 22	Sept. 24	3½ qrs.
Long Hoos, east	Clover	Grey Winter	Apr. 24, '15	June 22	July 5	July 21	
" " west	"	Red	Apr. 24, '15	June 22	July 5	July 21	
West Barn	Wheat	Rivetts	Nov. 8, '15	Sept. 8	Sept. 21	Sept. 22	5 qrs.†
Foster's, east	Potatoes	(King Edward, Arran Chief and Dalhousie)	May 11	...	Oct. 6	Oct. 20	3½ tons
" " west	Barley	Burton Brewing	Apr. 11	Sept. 1	Sept. 18	Sept. 20	6½ qrs.
Broadbalk	Wheat	Squareheads	Nov. 4, '15	Aug. 22	Sept. 11	Sept. 12	see p. 60
Little Hoos	"	"	Nov. 24, '15	Aug. 26	Sept. 13	Sept. 14	" 63
Hoos	Barley	Plumage Cross	Apr. 4	Sept. 1	Sept. 14	Sept. 16	" 61
Barnfield	Mangolds	Sutton's Yellow Globe	May 11	...	Nov. 1	Nov. 29	" 56
Agdell	Swedes	Magnum Bonum	June 16	...	Feb. 26, '17	Feb. 28, '17	" 54
Greatfield	Grass	Cattle Grazing	...	...	...	...	
Park	"	"	...	June 20	July 3	July 5	see p. 57
"	"	"	...	Mar. 9, '17	Mr. 14, '17	Mr. 14, '17	" "

\* Setting out. † Measured Plot.

DATES OF SOWING AND HARVESTING, 1917.

Field.	Crop.	Variety.	Sowing began.	Cutting began.	Carting began.	Carting finished.	Yield per Acre.
Great Knott Wood, east	Oats	Grey Winter	Oct. 12, '16	Aug. 16	Sept. 1	Sept. 4	3½ qrs.
" " west	Barley	Burton Brewing	Apr. 25	Sept. 8	Sept. 14	Sept. 15	30 8bus.
Little Knott Wood	Potatoes	(King Edward, Arran Chief, ) (Scottish Farmer	May 2	...	Sept. 27	Oct. 15	5 tons
Sawpit, north	Oats	Grey Winter	Oct. 16, '16	Aug. 14	Sept. 3	Sept. 3	27 2bus.
" south	(Fallowed)	...	...	...	...	...	...
Great Harpenden	Wheat	(Squareheads Master (10 acres), ) Red Standard (10 acres), ) (Red Marvel (4 acres) ... )	Feb. 28, '17	Aug. 23	Sept. 5	Sept. 6	21 0bus. 25 6bus. 26 7bus.
New Zealand	Barley	Burton Brewing	Apr. 21	Sept. 11	Sept. 24	Sept. 25	24 8bus.
Stackyard	Wheat	Rivetts	Nov. 1, '16	Sept. 13	Sept. 25	Sept. 26	10 9bus.
Long Hoos	"	"	Nov. 1, '16	Sept. 12	Sept. 26	Sept. 28	½ qrs.
West Barn	Oats	Grey Winter	Oct. 16, '16	Aug. 17	Sept. 7	Sept. 8	2½ qrs.
Foster's, west	Clover	Red	May 1, '16	June 14	June 21	June 22	½ ton
" east	Wheat	Red Standard	Nov. 15, '16	Aug. 21	Sept. 4	Sept. 5	21 4bus.
Broadbalk	"	"	Dec. 1, '16	Aug. 22	Sept. 3	Sept. 4	see p. 60
Little Hoos	Clover	Red	Apr. 26, '16	June 15	June 22	June 23	" 63
Hoos	Barley	Plumage Cross	Apr. 16	Sept. 6	Sept. 10	Sept. 11	" 61
Barnfield	Mangolds	Sutton's Yellow Globe	May 16	...	Nov. 16	Dec. 3	" 56
Agdell	Barley	Plumage Cross	Apr. 23	Sept. 13	Sept. 22	Sept. 23	" 54
Greatfield	Grass	...	...	June 26	July 3	July 7	" 57
Park	"	...	...	July 6	July 12	July 13	" 57
"	"	...	...	Dec. 14	Dec. 15	Dec. 28	" "

+ Measured Plots.



## CROP YIELDS ON THE EXPERIMENTAL PLOTS.

1 acre	...	...	...	...	=	0.404 Hectare
1 bushel	...	...	...	...	=	0.364 Hectolitre.
1 lb. (pound avoird.)	...	...	...	...	=	0.453 Kilogramme.
1 cwt. (hundredweight)	...	...	...	...	=	50.8 Kilogrammes.
1 metric quintal	...	...	...	...	=	(100.0 Kilogrammes. 220.46 lb.)
1 bushel per acre	...	...	...	...	=	0.9 Hectolitre per Hectare.
1 lb. per acre	...	...	...	...	=	1.12 Kilogramme per Hectare.
1 cwt. per acre	...	...	...	...	=	125.6 Kilogrammes per Hectare or 1.256 metric Quintals per Hectare.

### Crops Grown in Rotation. Agdell Field.

#### PRODUCE PER ACRE.

Year.	CROP.	O.		M.		C.		
		Unmanured.		Mineral Manure.		Complete Mineral and Nitrogenous Manure.		
		5.	6.	3.	4.	1.	2.	
		Fallow.	Beans or Clover.	Fallow.	Beans or Clover.	Fallow.	Beans or Clover.	
<b>SEVENTEENTH COURSE, 1912-15.</b>								
1912	Roots (Swedes) Cwt.	8.2	2.3	151.7	251.9	586.6	463.0	
1913	Barley Grain Bush.	18.5	24.6	24.7	33.2	22.0	32.5	
	Barley Straw ... Cwt.	8.2	13.0	10.6	14.5	9.0	15.0	
1914	Clover Hay ... Cwt. (1 crop)	—	4.1	—	6.5	—	3.9	
1915	Wheat Grain Bush.	3.2	6.3	13.2	15.2	13.3	10.5	
	Wheat Straw ... Cwt.	11.2	13.2	19.9	19.8	17.1	10.8	
<b>PRESENT COURSE (18th), 1916-17.</b>								
1916	Roots (Swedes) Cwt.	12.5	1.4	125.2	145.3	285.2	37.8	
1917	Barley Grain Bush.	9.4	2.5	14.2	15.2	13.1	15.0	
	Barley Straw ... Cwt.	11.6	5.1	16.8	15.6	13.1	19.8	

### METEOROLOGICAL RECORDS, 1915-17

	Rain.			Drainage through soil.			Bright Sun- shine.	Temperature. (Means).		
	Total Fall.		No. of Rainy Days.	20 ins. deep.	40 ins. deep.	60 ins. deep.		Hours.	Max.	Min.
	5-inch Funnel Gauge.	$\frac{1}{1000}$ Acre Gauge.								
	Inches.	Inches.	No.	Inches.	Inches.	Inches.		°F.	°F.	
<b>1915</b>										
Jan. ...	3.783	4.114	19	3.926	3.943	3.918	44.4	42.1	33.9	
Feb. ...	4.198	4.540	20	3.942	3.845	3.830	82.0	44.3	32.7	
Mar. ...	1.194	1.384	13	0.624	0.789	0.791	87.9	45.8	34.2	
April ...	1.088	1.222	9	0.129	0.196	0.187	161.6	53.3	35.8	
May ...	2.337	2.477	8	1.222	1.279	1.279	236.9	61.3	42.2	
June ...	1.727	1.793	9	0.360	0.358	0.341	242.0	67.5	47.3	
July ...	4.390	4.717	16	1.841	2.010	1.827	188.7	66.2	50.7	
Aug. ...	2.385	2.587	14	1.166	1.235	1.154	173.7	67.2	52.1	
Sept. ...	2.300	2.491	8	0.825	0.781	0.743	187.7	63.7	47.3	
Oct. ...	2.375	2.597	13	1.453	1.301	1.204	63.4	53.8	41.8	
Nov. ...	2.162	2.376	15	1.932	2.115	1.915	85.0	42.9	31.7	
Dec. ...	5.149	5.561	25	5.316	5.381	5.232	33.5	46.6	35.6	
<b>Total or Mean</b>	<b>33.088</b>	<b>35.859</b>	<b>169</b>	<b>22.736</b>	<b>23.233</b>	<b>22.421</b>	<b>1586.8</b>	<b>54.6</b>	<b>40.4</b>	
<b>1916</b>										
Jan. ...	2.067	2.237	14	1.826	2.001	1.897	49.1	49.0	38.6	
Feb. ...	3.279	3.974	23	3.387	3.337	3.273	76.5	42.4	32.5	
Mar. ...	3.841	4.919	23	5.550	6.052	5.608	63.2	42.4	32.3	
April ...	1.338	1.430	12	0.212	0.365	0.308	197.5	55.1	37.8	
May ...	1.819	1.970	15	0.633	0.738	0.713	185.0	61.9	44.0	
June ...	2.558	2.711	17	0.057	0.154	0.128	136.7	58.7	44.9	
July ...	1.610	1.771	10	0.281	0.397	0.347	161.2	66.5	50.0	
Aug. ...	3.319	3.576	16	1.111	1.209	1.156	174.4	69.6	53.0	
Sept. ...	1.497	1.673	11	0.446	0.510	0.452	106.2	61.2	47.4	
Oct. ...	3.399	3.696	24	2.104	2.095	2.022	88.5	56.5	44.3	
Nov. ...	4.193	4.491	17	4.260	4.353	4.452	73.8	48.4	36.8	
Dec. ...	3.065	3.386	16	3.088	3.162	3.023	24.8	39.7	29.7	
<b>Total or Mean</b>	<b>31.985</b>	<b>35.834</b>	<b>198</b>	<b>22.955</b>	<b>24.373</b>	<b>23.379</b>	<b>1336.9</b>	<b>54.3</b>	<b>40.9</b>	
<b>1917</b>										
Jan. ...	1.598	1.795	17	1.501	1.693	1.662	22.9	35.7	30.1	
Feb. ...	0.787	0.927	11	0.758	0.566	0.694	49.8	37.7	28.1	
Mar. ...	1.497	1.826	17	0.818	0.874	0.830	72.3	42.3	30.0	
April ...	1.935	2.154	16	1.226	1.306	1.061	138.8	48.3	32.6	
May ...	1.819	1.980	12	0.670	0.775	0.747	223.7	65.6	45.2	
June ...	1.960	2.152	12	0.365	0.428	0.443	207.1	69.9	50.3	
July ...	4.200	4.567	10	2.236	2.336	2.250	212.1	68.6	51.7	
Aug. ...	6.049	6.514	22	4.378	4.424	4.250	147.9	65.3	53.4	
Sept. ...	1.829	2.076	13	0.686	0.704	0.669	155.4	51.1	49.4	
Oct. ...	4.636	5.097	22	3.403	3.242	3.133	155.3	52.3	38.1	
Nov. ...	1.108	1.414	16	0.910	0.981	0.927	50.6	50.0	39.6	
Dec. ...	0.651	0.761	14	0.182	0.189	0.201	70.4	39.1	29.0	
<b>Total or Mean</b>	<b>28.069</b>	<b>31.263</b>	<b>182</b>	<b>17.133</b>	<b>17.518</b>	<b>16.867</b>	<b>1506.3</b>	<b>52.2</b>	<b>39.8</b>	

Mangolds, Barn Field, 1915, 1916, 1917.

Strip.	Strip Manures.	Cross Dressings.				
		O.	N.	A.	A.C.	C.
		None.	Nitrate of Soda	Ammon. Salts.	Ammon. Salts and Rape Cake.	Rape Cake.
<b>1915</b>						
1	Dung only ... ..	Tons. { R. 16 36 L. 3 15	Tons. { 26 85 4 42	Tons. { 21 42 4 69	Tons. { 22 56 6 00	Tons. { 21 97 4 80
2	Dung, Super., Potash ...	{ R. 15 00 L. 2 82	{ 21 69 4 24	{ 23 07 4 96	{ 29 59 6 62	{ 22 39 4 57
4	Complete Minerals ...	{ R. 1 91 L. 0 67	{ R. 8 65 L. 2 36 R. 6 35 L. 1 96	2 11	4 63	3 08
5	Superphosphate only ...	{ R. 1 29 L. 0 70	{ 2 22 0 82	{ 0 94 0 76	{ 4 72 2 63	{ 2 86 1 60
6	Super. and Potash ...	{ R. 1 71 L. 0 73	{ 7 54 2 06	{ 4 82 1 81	{ 10 13 3 27	{ 6 77 2 13
7	Super., Sulphate of Mag., and Sodium Chloride	{ R. 1 97 L. 0 83	{ 9 20 2 56	{ 6 71 2 17	{ 10 55 3 38	{ 10 62 2 54
8	None ... ..	{ R. 1 19 L. 0 75	{ 3 30 1 52	{ 0 69 0 59	{ 3 88 2 42	{ 4 70 1 95
<b>1916</b>						
1	Dung only ... ..	{ R. 19 37 L. 3 09	{ 31 93 4 54	{ 27 68 5 97	{ 28 04 5 37	{ 26 45 4 81
2	Dung, Super., Potash ...	{ R. 23 59 L. 3 90	{ 33 91 5 80	{ 34 17 7 95	{ 36 78 9 05	{ 32 45 6 41
4	Complete Minerals ...	{ R. 3 24 L. 0 65	{ R. 21 42 L. 3 77 R. 20 68 L. 4 13	19 65	34 28	27 37
5	Superphosphate only ...	{ R. 3 54 L. 0 66	{ 18 80 2 92	{ 9 63 3 51	{ 12 16 3 06	{ 14 64 2 97
6	Super. and Potash ...	{ R. 3 03 L. 0 62	{ 19 22 2 43	{ 20 34 2 78	{ 32 02 5 68	{ 25 00 2 33
7	Super., Sulphate of Mag., and Sodium Chloride	{ R. 3 54 L. 0 76	{ 20 25 3 46	{ 20 99 3 45	{ 30 10 5 57	{ 27 15 3 14
8	None ... ..	{ R. 2 32 L. 0 67	{ 10 28 2 62	{ 6 85 3 18	{ 10 66 3 03	{ 11 59 2 77
9	Sulphate of Mag. Sodium Chloride and Nitrate of Soda ... ..	{ R. 20 44 L. 2 70				
<b>1917</b>						
1	Dung only ... ..	{ R. 23 16 L. 2 88	{ 31 97 3 34	{ 24 02 2 52	{ 23 72 2 78	{ 25 09 3 46
2	Dung & Superphosphate (Potash omitted) ...	{ R. 27 71 L. 2 71	{ 32 68 3 39	{ 32 45 3 81	{ 33 44 4 41	{ 31 05 3 57
4	Sodium Chloride and Super. (Potash & Mag. omitted) ... ..	{ R. 3 92 L. 0 44	{ R. 17 93 L. 1 79 R. 17 12 L. 1 94	19 21	22 57	19 15
5	Superphosphate only ...	{ R. 3 28 L. 0 46	{ 15 03 1 61	{ 6 85 1 57	{ 8 36 1 84	{ 9 90 1 77
6	Superphosphate only (Potash omitted) ...	{ R. 2 38 L. 0 36	{ 13 87 1 21	{ 15 11 1 25	{ 20 60 2 41	{ 15 46 1 13
7	Sodium Chloride, and Superphosphate ...	{ R. 2 56 L. 0 41	{ 18 17 1 56	{ 19 69 1 47	{ 22 79 2 40	{ 19 27 1 38
8	None ... ..	{ R. 1 89 L. 6 38	{ 10 22 1 51	{ 5 89 1 46	{ 8 45 1 54	{ 7 48 1 23
9	Sodium Chloride, Nitrate of Soda ... ..	{ R. 19 56 L. 1 65				

R. = roots. L. = leaves. Tons per acre in all cases.



Hay. The Park Grass Plots. 1915, 1916, 1917.

Plot.	Manuring.	Yield of Hay per acre.			Yield of Hay per acre.			Yield of Hay per acre.			Average for 57 years 1856-1912 (1st and 2nd crops).	Plot.
		1915.			1916.			1917.				
		1st Crop.	2nd Crop.	Total.	1st Crop.	2nd Crop.	cwt.	1st Crop.	2nd Crop.	cwt.		
1	Amm. Salts alone ; with Dung 8 years, 1856-63 ...	{ not limed lined ...	8.5 17.0	25.5	15.1	1.5	7.0	9.1	35.9	cwt.	1	
2	Unmanured ; Dung 8 years, 1856-63 ...	{ not limed lined ...	7.6 17.7	25.3	15.9	1.5	15.1	1.6	28.6	cwt.	2	
3	Unmanured ...	{ not limed lined ...	5.9 13.4	19.3	10.8	.67	14.3	1.6	20.9	cwt.	3	
4-1	Superphosphate of Lime ...	{ not limed lined ...	5.3 10.9	16.2	7.4	.36	11.2	1.2	21.6	cwt.	4-1	
4-2	Superphosphate of Lime and Amm. Salts ...	{ not limed lined ...	6.7 16.1	22.8	15.1	.39	21.3	2.7	33.5	cwt.	4-2	
5-1	(N. half) Unmanured ; following Amm. Salts alone, 1856-97 ...	{ not limed lined ...	4.5 14.4	18.8	10.8	.96	3.7	8.5	14.4a	cwt.	5-1	
5-2	(S. half) Complete Minerals ; following Amm. Salts alone, 1856-97 ...	{ not limed lined ...	3.3 6.6	9.9	18.3	.37	7.3	3.7	23.2a	cwt.	5-2	
6	Complete Mineral Manure as plot 7 ; following Amm. Salts alone, 1856-68 ...	{ not limed lined ...	17.0 23.5	40.4	30.0	1.3	20.5	9.1	37.2	cwt.	6	
7	Complete Mineral Manure ...	{ not limed lined ...	2.1 5.0	7.0	8.4	2.7	29.2	21.0	40.9	cwt.	7	
8	Mineral Manure without Potash ...	{ not limed lined ...	18.7 28.3	47.0	20.8	2.4	39.5	6.8	54.3	cwt.	8	
9	Complete Mineral Manure and Amm. Salts ...	{ not limed lined ...	16.3 29.2	45.4	21.5	3.3	40.4	17.4	47.7	cwt.	9	
10	Mineral Manure (without Potash) and Amm. Salts ...	{ not limed lined ...	18.8 28.7	47.5	31.0	2.4	26.0	20.9	66.5	cwt.	10	
11-1	Complete Mineral Manure and extra Amm. Salts ...	{ not limed lined ...	8.5 16.7	25.2	12.4	3.9	40.3	15.5	73.3	cwt.	11-1	
11-2	As plot 11-1 and Silicate of Soda ...	{ not limed lined ...	5.0 16.8	21.8	10.9	4.2	21.4	16.0	23.9	cwt.	11-2	
12	Unmanured ...	{ not limed lined ...	9.3 29.9	39.2	40.7	5.3	14.1	4.5	—	cwt.	12	
13	Dung and Fish Guano, once in 4 years ...	{ not limed lined ...	26.8 24.4	51.1	46.0	3.9	30.4	15.5	56.9	cwt.	13	
14	Complete Mineral Manure and Nitrate of Soda = 86 lbs. N. ...	{ not limed lined ...	8.5 15.8	24.3	27.7	3.9	27.7	10.4	36.8	cwt.	14	
15	Complete Mineral Manure as plot 7 ; following Nitrate of Soda alone, 1858-75 ...	{ not limed lined ...	24.9 39.0	63.8	52.7	3.5	39.2	15.0	46.3	cwt.	15	
16	Complete Mineral Manure and Nitrate of Soda = 43 lbs. N. ...	{ not limed lined ...	45.0 30.2	75.2	54.8	2.2	33.7	6.7	33.7	cwt.	16	
17	Nitrate of Soda alone ...	{ not limed lined ...	50.0 36.9	86.9	54.1	1.3	24.6	14.2	—	cwt.	17	
18	Potash, Sulphate of Soda, Magnesia, and Sulphate of Amm. ...	{ not limed lined ...	67.1 34.6	101.7	52.3	1.7	25.0	12.7	—	cwt.	18	
19	Farmyard Dung ...	{ not limed lined ...	4.6 10.9	15.5	9.1	1.7	23.1	8.1	—	cwt.	19	
20	Farmyard Dung ...	{ not limed lined ...	30.5 27.4	57.9	35.0	1.7	25.0	13.3	—	cwt.	20	

Ground lime was applied to the Southern portion (lined) of the plots at the rate of 2,000 lb. to the acre in 1903, 1907, and March, 1915.  
 In 1917 all Potash and Magnesia were omitted from the Mineral Manures in plots 5-2, 6, 7, 9, 11-1, 11-2, 14, 15, 16, and 18; and Dung was omitted from plots 19 and 20.  
 Up to 1914 the limed and unlimed plot results were not separately given in the Annual Report, but the mean of the two was given. From 1915 onwards the separate figures are given.  
 a - Average for 15 years 1893-1912.



The Park  
BOTANICAL COMPO-

Plot.	Manuring.	Liming.	Crop.	1915.		
				Gram-ineae.	Legu-minosae.	Other Orders.
3	Unmanured ... ..	Not limed	1st	52.52	5.29	42.19
		"	2nd	42.08	11.49	46.43
4-1	Superphosphate of Lime ... ..	Whole plot	1st	44.91	17.45	37.64
		"	2nd	45.39	8.87	45.73
4-2	Super. of Lime and Amm. Salts ...	Not limed	1st	93.63	—	6.37
		"	2nd	98.98	0.24	0.77
4-2	Super. of Lime and Amm. Salts ...	Limed ...	1st	98.81	—	1.19
		"	2nd	98.83	—	1.17
5-2	(S. half) Complete Minerals; follow- ing Amm. Salts alone, 1856-97 ...	Not limed	1st	79.07	6.02	14.91
		Whole Plot	*2nd	59.70	19.32	20.99
6	Complete Mineral Manure as plot 7; following Amm. Salts alone, 1856-68	Not limed	1st	51.38	33.79	14.83
		"	2nd	60.94	27.92	11.14
7	Complete Mineral Manure ... ..	Not limed	1st	60.85	24.76	14.39
		"	2nd	50.19	36.73	13.08
7	Complete Mineral Manure ... ..	Limed ...	1st	53.76	36.90	9.34
		"	2nd	55.44	34.44	10.12
8	Mineral Manure without Potash ...	Not limed	1st	52.73	11.00	36.27
		"	2nd	44.24	13.88	41.88
8	Mineral Manure without Potash ...	Limed ...	1st	50.54	22.43	27.03
		"	2nd	40.51	22.21	37.27
9	Complete Mineral Manure and Amm. Salts ... ..	Not limed	1st	89.39	—	10.61
		"	2nd	85.98	—	14.02
9	Complete Mineral Manure and Amm. Salts ... ..	Limed ...	1st	98.40	0.14	1.46
		"	2nd	97.72	0.32	1.96
10	Mineral Manure (without Potash) and Amm. Salts ... ..	Not limed	1st	98.38	—	1.62
		"	2nd	96.20	—	3.80
10	Mineral Manure (without Potash) and Amm. Salts ... ..	Limed ...	1st	99.64	—	0.36
		"	2nd	99.00	—	1.00
11-2	Complete Mineral Manure and extra Amm. Salts and Silicate of Soda	Not limed	1st	99.71	—	0.29
		"	2nd	99.59	—	0.41
11-2	Complete Mineral Manure and extra Amm. Salts and Silicate of Soda	Limed ...	1st	100.0	—	—
		"	2nd	99.65	—	0.35
14	Complete Mineral Manure and Ni- trate of Soda = 86 lb. N.	Not limed	1st	88.41	4.41	7.18
		"	2nd	80.71	11.73	7.56
15	Complete Mineral Manure as plot 7; following Nitrate of Soda alone, 1858-75 ... ..	Not limed	1st	49.77	38.94	11.29
		"	2nd	54.84	32.74	12.42
19	Farmyard Dung ... ..	Not limed	1st	68.91	19.65	11.44
		"	2nd	58.80	33.67	7.52
20	Farmyard Dung ... ..	Not limed	1st	76.95	11.91	11.14
		"	2nd	73.81	14.98	11.20

\* 2nd Crop was sampled from whole of plot 5 (i.e. 5-1 and 5-2).

Grass Plots.

SITATION, PER CENT.

1916.			1917.			"Other Orders" consist largely of	Plot.
Gram-ineæ.	Legu-minosæ.	Other Orders.	Gram-ineæ.	Legu-minosæ.	Other Orders.		
66·04	8·79	25·17	43·96	5·53	50·51	Leontodon hispidus and Centaurea nigra (very varied herbage)	3
64·12	7·49	28·39	51·44	4·98	43·58	Leontodon hispidus, Centaurea nigra, and Plantago lanceolata	4-1
99·63	—	0·37	91·8	—	8·2	Rumex acetosa ... ..	4-2
98·33	—	1·67	98·60	—	1·40	Rumex acetosa and Galium verum	4-2
85·34	2·33	12·32	72·56	10·96	16·48	Centaurea nigra and Rumex acetosa	5-2
74·08	17·42	8·50	61·64	25·87	12·48	Centaurea nigra and Achillea millefolium	6
74·84	15·14	10·02	59·11	11·36	29·52	Centaurea nigra and Achillea millefolium	7
69·94	26·31	3·75	70·96	18·21	10·83	Centaurea nigra ... ..	7
69·00	8·27	22·73	48·31	2·69	49·01	Centaurea nigra and Plantago lanceolata	8
71·34	7·50	21·16	58·70	4·70	36·60	Centaurea nigra and Plantago lanceolata	8
—	—	—	85·90	0·06	14·01	Rumex acetosa ... ..	9
—	—	—	98·28	—	1·72	Rumex acetosa and Achillea millefolium	9
—	—	—	93·18	—	6·82	Rumex acetosa ... ..	10
—	—	—	99·9	—	0·1	Rumex acetosa ... ..	10
100·0	—	—	—	—	—	... ..	11-2
99·36	—	0·64	—	—	—	Heracleum sphondylium and Rumex acetosa	11-2
83·75	6·68	9·57	—	—	—	Taraxacum vulgare, Anthriscus sylvestris	14
—	—	—	70·18	15·74	14·07	Achillea millefolium ... ..	15
74·46	19·20	6·34	68·7	21·38	9·92	Anthriscus sylvestris, Rumex acetosa, Centaurea nigra, Achillea millefolium	19
81·83	12·03	6·14	66·13	24·95	8·92	Anthriscus sylvestris, Centaurea nigra, Achillea millefolium	20

2nd Crop, 1916, was very small and was not sampled for Botanical Analysis.  
2nd Crop, 1917, results not yet available.

Wheat. Broadbalk Field, 1915-17. Produce.

Plot.	1915. (Top portion).			1916. (Bottom portion).			1917. (Bottom portion).			Average for 61 years, 1852-1912.	
	Dressed Grain.		Straw per Acre.	Dressed Grain.		Straw per Acre.	Dressed Grain.		Straw per Acre.	Dressed Grain per Acre.	Straw per Acre.
	Yield per Acre.	Weight per Bushel.		Yield per Acre.	Weight per Bushel.		Yield per Acre.	Weight per Bushel.			
	Bushels	lb.	cwt.	Bushels	lb.	cwt.	Bushels	lb.	cwt.	Bushels	cwt.
2	32.2	62.0	37.8	33.3	61.0	41.3	16.1	57.6	14.8	35.2	34.8
3	12.1	62.3	12.4	16.4	61.0	15.8	8.2	59.9	5.5	12.6	10.3
5	15.8	62.6	15.8	18.5	60.2	20.8	9.9	60.8	7.1	14.5	12.1
6	26.7	62.7	27.4	25.4	60.8	24.1	18.1	61.3	13.4	23.2	21.4
7	33.9	62.4	34.5	31.3	60.3	40.9	23.3	60.3	18.3	32.1	32.9
8	37.5	61.5	40.9	31.7	60.5	42.4	30.3	59.7	25.5	36.6	41.1
9	30.3	62.5	32.2	29.2	60.6	35.5	20.6	57.7	18.8	—	—
10	19.3	62.4	20.4	18.5	60.3	26.6	13.8	57.1	9.9	20.0	18.4
11	28.2	61.6	27.5	13.6	59.3	24.9	14.6	57.7	11.3	22.9	22.3
12	33.0	61.5	32.3	22.5	59.7	33.6	19.0	58.5	13.7	29.1	28.0
13	33.2	60.6	38.0	25.1	60.2	35.8	29.8	60.3	22.9	31.0	31.5
14	30.5	60.6	31.0	21.4	59.6	32.5	21.2	59.7	15.6	28.8	28.0
15	19.0	60.6	23.6	21.8	60.6	27.8	27.0	60.5	20.1	29.9	29.7
16	31.8	61.1	42.5	26.0	60.1	36.1	25.7	58.7	22.5	—	—
17	16.1	61.3	18.8	21.7	61.1	33.2	11.1	59.8	7.9	29.9	29.5
18	24.1	61.8	28.0	19.6	61.0	20.4	23.0	60.8	17.1	14.9	13.0
19	25.5	61.9	27.3	20.3	61.0	23.4	11.1	57.1	9.5	*25.4	*25.7
20	16.9	62.0	23.1	—	—	—	—	—	—	—	—

NOTE.—The top portion (western half) was fallow in 1914 owing to the weedy condition of the field. The bottom portion (eastern half) was fallow in 1915.  
\* 20 years, 1893-1912.

Wheat. Broadbalk Field, 1915-17. Manures.

Plot.	1915 and 1916.	1917.
2	Farmyard Manure ... ..	Farmyard Manure
3	Unmanured ... ..	Unmanured
5	Complete Mineral Manure ... ..	Complete Minerals (Potash omitted)
6	As 5, and single Amm. Salts ... ..	As 5, and single Amm. Salts
7	As 5, and double Amm. Salts ... ..	As 5, and double Amm. Salts
8	As 5, and treble Amm. Salts .. ..	As 5, and treble Amm. Salts
9	As 5, and single Nitrate Soda ... ..	As 5, and single Nitrate Soda
10	Double Amm. Salts alone ... ..	Double Amm. Salts alone
11	As 10, and Superphosphate ... ..	As 10, and Superphosphate
12	As 10, and Super. and Sulph. Soda ... ..	As 10, and Super. (Sulphate of Soda omitted)
13	As 10, and Super. and Sulph. Potash ... ..	As 10, and Super. (Potash omitted)
14	As 10, and Super. and Sulph. Magnesia ... ..	As 10, and Super. (Magnesia omitted)
15	Double Amm. Salts in Autumn and Minerals ... ..	Double Amm. Salts in Autumn, and Minerals (Potash omitted)
16	Double Nitrate and Minerals ... ..	Double Nitrate and Minerals (Potash omitted)
17 } 18 }	Minerals alone, or double Amm. Salts alone, in alternate years ... ..	(Minerals alone (Potash omitted), or double Amm. Salts alone, in alternate years
19	Rape Cake alone ... ..	Rape Cake alone
20	Mineral Manure (without Super.) and Amm. Salts ... ..	Sulphate of Soda, Sulphate of Magnesia, and Sulphate of Ammonium

NOTE.—No Autumn manures were applied for the 1915 crop: dressings were given in the Autumn of 1913, but not in 1914 as that half of the field was then left fallow. In 1916, Sulphate of Potash being short, the dressing was in each plot made up with the required amount of woodash.

ERRATA.—Plot 19, 1917, for "Rape Cake only" read "Rape Cake omitted," there being none applied in this year.



Permanent Barley Plots. Hoos Field, 1915, 1916, 1917.  
MANURING PRODUCE PER ACRE

Plot.	1915 and 1916.		1917.		1915		1916		1917		Average 60 years, 1852-1911.	
					Dressed Grain.	Weight per Bushel.	Straw.	Dressed Grain.	Weight per Bushel.	Straw.	Dressed Grain.	Straw.
1 O	Unmanured		Unmanured		Bush.	lb.	cwt.	Bush.	lb.	cwt.	Bush.	cwt.
2 O	Superphosphate only		Superphosphate only		8.5	51.8	5.1	23.4	55.8	11.4	12.7	8.4
3 O	Alkali Salts only		Sulphate of Soda		10.9	52.3	7.7	36.5	55.3	16.8	19.7	10.0
4 O	Complete Minerals		Super. and Sulphate of Soda		10.8	52.8	10.4	24.6	55.7	15.7	15.2	8.8
5 O	Potash and Superphosphate		Superphosphate		12.7	53.5	13.1	37.4	55.3	20.9	19.7	11.1
1 A	Ammonium Salts only		Ammonium Salts only		12.0	53.3	10.9	18.6	56.4	13.0		
2 A	Superphosphate and Amm. Salts		Super. and Amm. Salts		21.5	51.6	8.5	34.8	54.6	17.4	25.5	14.7
3 A	Alkali Salts and Ammonium Salts		Sulphate of Soda and Amm. Salts		20.2	51.5	10.1	44.7	53.8	23.7	38.2	22.0
4 A	Complete Minerals and Amm. Salts		Super., Sulphate of Soda and Amm. Salts		29.2	51.9	13.5	*	54.2	21.9	28.0	16.9
5 A	Potash, Super. and Amm. Salts		Super. and Amm. Salts		38.1	53.0	18.5	46.8	54.3	27.5	41.5	25.0
1 AA	Nitrate of Soda only		Nitrate of Soda only		36.2	52.8	20.2	40.1	55.6	24.9		
2 AA	Super. and Nitrate of Soda		Super. and Nitrate of Soda		24.8	52.3	12.4	34.8	55.2	20.2	29.3	17.8
3 AA	Alkali Salts and Nitrate of Soda		Sulphate of Soda and Nitrate of Soda		32.9	53.1	16.9	47.4	55.5	28.5	43.1	26.3
4 AA	Complete Minerals and Nitrate of Soda		Sulphate of Soda, Super. and Nitrate of Soda		29.9	52.9	14.6	34.7	55.5	22.5	30.0	19.3
1 AAS	As Plot 1 AA and Silicate of Soda		As Plot 1 AA and Silicate of Soda		35.2	53.6	17.1	45.1	55.1	28.8	42.7	27.3
2 AAS	" " 2 AA		" " 2 AA		31.7	53.0	15.0	41.1	55.3	22.1	32.8 (1)	19.7 (1)
3 AAS	" " 3 AA		" " 3 AA		36.2	53.0	18.6	50.6	55.3	29.2	42.3 (1)	26.0 (1)
4 AAS	" " 4 AA		" " 4 AA		31.8	53.2	16.0	39.2	55.3	23.3	35.2 (1)	21.7 (1)
1 C	Rape Cake only		Rape Cake omitted		36.2	53.6	17.9	46.2	54.9	29.0	43.6 (1)	27.7 (1)
2 C	Superphosphate and Rape Cake		Super. (Rape Cake omitted)		34.0	53.2	14.7	45.4	55.2	23.9	38.3	22.1
3 C	Alkali Salts and Rape Cake		Sulph. of Soda (Rape Cake omitted)		34.0	53.0	14.8	45.7	55.3	25.9	40.5	23.6
4 C	Complete Minerals and Rape Cake		Sulphate of Soda and Super. (Rape Cake omitted)		32.1	53.1	14.2	46.1	55.4	23.4	36.9	22.3
7-1	Unmanured (after dung 20 years, 1852-71)		Unmanured (after dung 20 years, 1852-71)		33.0	53.1	15.9	46.8	55.6	26.8	40.5	24.5
7-2	Farmyard Manure		Farmyard Manure		16.3	52.8	12.9	29.7	55.7	16.7	24.8 (2)	14.8 (2)
6-1	Unmanured		Unmanured		31.4	53.7	19.0	31.9	55.2	30.9	47.1	29.6
6-2	Ashes		Sifted Ashes from the Lab. furnace		9.7	52.0	6.2	25.8	55.0	14.3		
1 N	Nitrate of Soda only		Nitrate of Soda only		7.3	52.5	6.0	29.4	54.4	16.7		
2 N	" " " "		" " " "		22.3	52.1	11.5	35.8	53.8	20.4		
	" " " "		" " " "		29.3	53.2	14.3	43.4	55.2	25.2		

(1) 48 years, 1864-1911. (2) 40 years, 1872-1911. \* Plot 3A figures lost.

Hoos Field (formerly Potato Plots). No Manure since 1901.

Plot.	Manure given prior to 1901.	1915. OATS.				1916. BARLEY.				1917. BARLEY.			
		Dressed Grain.		Total Produce per Acre.	Straw per Acre.	Dressed Grain.		Total Produce per Acre.	Straw per Acre.	Dressed Grain.		Total Produce per Acre.	Straw per Acre.
		Yield per Acre.	Weight per Bush.			Yield per Acre.	Weight per Bush.			Yield per Acre.	Weight per Bush.		
Previous Cropping: Potatoes, 1876-1901; Barley, 1902 and 1903; Oats, 1904; Barley, 1905-1911; Oats, 1912; Barley, 1913 and 1914.													
1	Unmanured ...	Bush. 9.7	lb. 32.3	cwt. 8.3	lb. 1276	Bush. 10.3	lb. 55.6	cwt. 6.9	lb. 1402	Bush. 5.1	lb. 49.5	cwt. 3.7	lb. 707
2	Unmanured 1882 to 1901, previously Dung only ...	19.4	32.6	17.3	2618	17.8	56.1	10.8	2270	9.9	48.3	6.4	1244
3	Dung 1883 to 1901 ...	23.6	30.2	20.8	3097	17.7	56.5	13.5	2579	8.0	49.1	6.3	1144
4	Dung 1883 to 1901 ...	26.5	32.1	26.3	3842	21.0	56.2	14.7	2902	10.7	49.3	7.3	1408
Previous Cropping: Potatoes, 1876-1901; Barley, 1902-1903; Oats, 1904; Plots 5, 7, 9, Cow Peas (failed), 1905; Plots 6, 8, 10, Red Clover, 1905; Red Clover, 1906-1911; Oats, 1912; Barley, 1913 and 1914.													
5	Ammonium Salts ...	13.3	34.0	14.4	2103	8.9	55.4	7.1	1335	5.6	49.5	3.7	725
6	Nitrate of Soda ...	11.6	32.8	14.3	2012	11.2	56.5	8.6	1647	7.7	49.5	4.6	947
7	{ Ammonium Salts and Mixed Minerals ... }	25.8	33.0	25.2	3709	23.1	55.5	13.8	2896	12.9	48.7	7.3	1506
8	{ Nitrate of Soda and Mixed Minerals ... }	28.7	33.5	27.4	4077	21.0	55.8	11.5	2508	11.9	50.3	6.9	1421
9	Superphosphate ...	22.0	32.6	19.3	2923	19.8	55.2	10.3	2296	6.9	49.3	3.9	812
10	Mixed Minerals ...	23.5	33.3	24.4	3572	22.5	55.7	13.3	2810	7.1	50.0	4.8	932



## Little Hoos Field PLAN OF ROTATION PLOTS

Arranged to test the RESIDUAL VALUE of VARIOUS MANURES in one, two, three, and four years after their application. Produce per acre.

		1915 (12th Season). Mangolds.			1916 (13th Season). Wheat.			1917 (14th Season). Clover.			
		Roots.	Leav's	Total Pr'd'ce.	Dress- ed Grain.	Straw.	Total Pr'd'ce.	1st Crop.	2nd Crop.	Total.	
		Tons.	Tons.	Tons.	Bush.	cwt.	lb.	cwt.	cwt.	cwt.	
A	1	Control ...	5.36	2.03	7.39	20.7	19.1	3492	19.7	22.2	41.9
	2	Dung (ordinary) 16 tons per acre	<i>a</i> 9.44	2.16	11.60	<b>28.4</b>	<b>30.4</b>	<b>5292</b>	38.2	27.4	65.6
	3		<i>b</i> 11.66	2.25	13.91	26.8	26.6	4766	31.7	27.7	59.4
	4		<i>c</i> 9.37	1.97	11.34	25.7	25.1	4520	26.2	26.6	52.8
	5		<i>d</i> <b>13.53</b>	<b>2.92</b>	<b>16.45</b>	27.6	28.9	5040	31.9	26.1	58.0
B	1	Dung (cake fed)	9.52	2.44	11.96	<b>34.0</b>	<b>34.1</b>	<b>6003</b>	38.4	27.2	65.6
	2	Control ...	7.18	2.20	9.38	21.3	17.4	3324	19.1	21.7	40.8
	3	Dung (cake fed) 16 tons per acre	<i>b</i> 13.00	2.58	15.58	27.2	28.3	4952	34.3	26.9	61.2
	4		<i>c</i> 11.81	2.47	14.28	26.6	27.1	4772	35.8	26.1	61.9
	5		<i>d</i> <b>15.27</b>	<b>3.02</b>	<b>18.29</b>	29.5	29.6	5135	32.7	26.9	59.6
C	1	Shoddy 956 lb. per	5.17	1.99	7.16	<b>13.6</b>	<b>14.7</b>	<b>2668</b>	17.6	24.0	41.6
	2	acre ...	<i>b</i> 6.42	2.15	8.57	11.4	10.6	1961	18.6	21.7	40.3
	3	Control ...	6.87	2.39	9.26	16.5	12.6	2523	20.7	24.6	45.3
	4	Shoddy 956 lb. per	<i>c</i> 7.23	2.56	9.79	18.6	13.9	2800	19.7	25.1	44.8
	5	acre ...	<i>d</i> <b>8.71</b>	<b>2.79</b>	<b>11.50</b>	20.3	14.1	2923	17.6	26.6	44.2
D	1	Guano 776 lb. per	4.90	1.60	6.50	<b>22.0</b>	<b>20.4</b>	<b>3801</b>	19.4	27.4	46.8
	2	acre ...	<i>b</i> 7.37	1.96	9.33	17.1	13.0	2605	18.6	24.0	42.6
	3	Control ...	7.35	2.06	9.41	18.2	11.1	2452	21.7	25.6	47.3
	4	Guano ...	<i>c</i> 6.64	2.20	8.84	17.8	13.6	2734	20.4	27.2	47.6
	5	Control ...	<i>d</i> <b>8.39</b>	<b>2.69</b>	<b>11.08</b>	13.3	12.8	2348	19.7	29.0	48.7
E	1	Rape Cake 1036	5.30	1.81	7.11	<b>18.4</b>	<b>15.5</b>	<b>2986</b>	18.6	25.6	44.2
	2	lb. per acre ...	<i>b</i> 7.80	2.03	9.83	18.6	13.8	2768	18.9	26.1	45.0
	3	Control ...	<i>c</i> 7.84	2.19	10.03	20.9	12.3	2749	19.7	24.3	44.0
	4	Rape Cake ...	<i>d</i> <b>8.71</b>	<b>2.95</b>	<b>11.66</b>	14.6	13.0	2466	20.2	28.2	48.4
	5	Control ...	4.70	1.62	6.32	10.1	12.0	2084	21.1	28.5	52.6
F	1	Control ...	4.80	2.09	6.89	11.7	11.4	2119	14.2	26.9	41.1
	2	Superphosphate 600 lb. per acre	<i>a</i> 7.90	2.29	10.19	<b>19.9</b>	<b>14.2</b>	<b>2906</b>	18.1	23.8	41.9
	3		<i>b</i> 8.51	2.35	10.86	17.7	13.9	2729	16.5	25.9	42.4
	4		<i>c</i> 7.36	2.42	9.78	20.3	16.2	3165	19.9	29.3	49.2
	5		<i>d</i> <b>7.23</b>	<b>2.11</b>	<b>9.34</b>	19.6	16.0	3098	23.0	30.8	53.8
G	1	Bone Meal 430 lb.	6.24	2.23	8.47	<b>19.7</b>	<b>15.1</b>	<b>2980</b>	14.7	29.0	43.7
	2	per acre ...	<i>b</i> 6.70	2.32	9.02	22.0	16.5	3235	15.2	29.3	44.5
	3	Control ...	5.90	2.08	7.98	20.4	16.3	3172	14.7	29.0	43.7
	4	Bone Meal 430 lb.	<i>c</i> 5.87	2.06	7.93	22.0	17.6	3430	17.8	29.5	47.3
	5	per acre ...	<i>d</i> <b>4.41</b>	<b>1.86</b>	<b>6.27</b>	23.6	18.7	3680	18.9	29.3	48.2
H	1	Basic Slag 600 lb.	9.01	1.99	11.00	<b>24.1</b>	<b>17.6</b>	<b>3528</b>	24.3	24.6	48.9
	2	per acre ...	<i>b</i> 10.34	2.25	12.59	26.5	19.5	3867	21.7	23.8	45.5
	3	Control ...	<i>c</i> 10.05	2.13	12.18	27.7	19.9	3921	23.3	22.5	45.8
	4	Basic Slag ...	<i>d</i> <b>8.56</b>	<b>1.93</b>	<b>10.49</b>	26.5	19.7	3874	21.0	22.8	43.8
	5	Control ...	6.23	2.11	8.34	24.6	19.3	3734	18.0	23.6	41.6

*a* received its dressings in 1912, 1916.  
*b* " " " 1913.

*c* received its dressings in 1910, 1914.  
*d* " " " 1911, 1915.

### NOTES AS TO MANURES.

The five plots of Series A to E which deal with nitrogenous manures received cross dressings as under:—

- 1904 3 cwt. Superphosphate per acre.
- 1906 3 cwt. Sulphate of Potash ditto.
- 1907, 1908, 1909 3 cwt. Superphosphate, each year.
- 1911 ditto plus 200 lbs. Sulphate of Potash, but no cross dressings have been applied since.

The five plots of Series F to H dealing with Phosphatic Manures received dressings as under:—

- 1904 1 cwt. Sulphate of Ammonia.
- 1905 ditto.
- 1906 2 cwt. ditto plus 3 cwt. Sulphate of Potash.
- 1907 1 cwt. Sulphate of Ammonia.
- 1908-10 ditto.
- 1911 ditto plus 200 lbs. Sulphate of Potash.
- 1912 1 cwt. Nitrate of Soda.
- 1914 1 cwt. Sulphate of Ammonia.
- 1915-16 ditto.

Clover was grown over the whole field in 1917; no manures applied in Autumn 1916 nor in Spring 1917. Thirteen tons of dung per acre was used on A and B for 1916 crop.

Figures in italics denote the unmanured plots.

The yields on the plots to which the manure was applied in a given year are printed in heavy type. In 13th season plots A and B were sown Nov. 24, 1915. C to H were sown Feb. 17, 1916.



Long Hoos Field. Green Manuring. 1914-15.  
WHEAT—Produce per Acre.

	No Treatment.		Dung 10 tons per Acre.		Mustard ploughed in	
	Grain.	Straw.	Grain.	Straw.	Grain.	Straw.
No artificials ... ..	Bushels. 14.4	cwt. 11.6	Bushels. 16.4	cwt. 20.5	Bushels. 20.8	cwt. 30.5
Superphosphate 3 cwt. per acre ... ..	—	—	18.2	18.8	26.9	26.0
Superphosphate 3 cwt. and Nitrate of Soda 1½ cwt. per acre ... ..	21.2	18.5	22.0	24.5	26.6	27.9
	—	—	21.6	24.7	27.2	32.5

SWEDES—Tons per Acre.

	No treatment.	Dung, 10 tons per Acre.	Winter Barley ploughed in.	Clover ploughed in.	Trifolium ploughed in.
No artificials ... ..	9.2 11.3 7.6	— — —	8.4	8.2	10.3
Superphosphate 3 cwt. per acre ... ..	— — — — —	12.0 14.0 12.8 12.9 12.7	10.6 10.5	10.7 8.3	11.0 10.4
Superphosphate 3 cwt., Sulphate of Ammonia 1½ cwt. per acre ... ..	11.4 13.3 13.5	7.4 9.3 12.6	7.0 12.9	6.0 11.0	8.9 12.3

No treatment } ploughed in January: fairly free from growth of weeds.  
Dung }

Winter Barley.—Ground well covered but not much bulk.

Clover.—Ground well covered but not much growth.

Trifolium.—Most of this died during winter, but there was a dense growth of annual weeds; not much bulk.

Barley (after Swedes). Long Hoos Field, 1916.

	No Treatment		Dung 10 Tons per Acre		Mustard ploughed in		Red Clover ploughed in		Rape ploughed in		Trifolium	
	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw	Grain	Straw
No artificials	Bush. 25.7 32.1 37.4 36.9 33.7	cwt. 14.7 17.4 19.4 18.2 19.6	Bush. 41.7 45.2 41.5 40.5	cwt. 21.1 24.2 21.9 21.0	bush. 34.9 30.0	cwt. 19.1 14.1	bush. 36.3 37.3	cwt. 19.7 18.4	bush. 34.7 32.7	cwt. 17.2 15.6	bush. 33.7 36.5	cwt. 17.0 19.5
Superphosphate 3 cwt. per acre	— — —	— — —	43.1 46.7 33.3 38.7	24.8 26.4 21.6 21.3	37.7	19.3	30.5	20.6	25.7	18.3	31.8	20.4
Superphosphate 3 cwt. per acre and Sulphate of Ammonia 1½ cwt. per acre	48.7 49.8 55.0 57.3 55.1	26.6 27.5 29.6 31.8 30.8	47.0 46.1 47.0 36.1	32.2 33.9 31.7 31.0	52.6 51.6	29.1 29.8	38.4 53.7	28.8 29.6	40.0 51.7	27.5 28.8	45.5 53.3	29.4 28.9

## CROP YIELDS FROM DUNG STORED IN DIFFERENT WAYS.

Yield of Potatoes manured with bullock dung, 10 tons of stored manure per acre, stored in heaps for three months.

West Barnfield, 1915.					
Yield of Potatoes in Tons per Acre.	No Manure.	Loose heap under cover.	Compact heap under cover.	Loose heap in the open.	Compact heap in the open.
Plot 1 ... ..	5·21	9·29	9·23	8·18	7·61
2 ... ..	4·95	8·36	8·82	8·00	7·18
3 ... ..	5·18	—	8·93	7·89	7·32
Mean	5·11	8·82	9·00	8·02	7·38
Percentage increase over the unmanured plots ... ..	—	73·00	76·00	57·00	44·00
Weight of original dung, tons per acre	—	15·52	12·96	13·68	12·05

## Great Harpenden Field, 1916.

Yield of Rivetts Wheat, manured at same rate from same heaps as above, six months later.

Dressed Grain per acre in bushels :—					
Plot 1 ... ..	34·6	37·3	40·0	35·6	31·7
2 ... ..	32·4	36·2	37·8	35·9	35·7
3 ... ..	34·6				
4 ... ..	31·1				
Mean	33·2	36·7	38·9	35·8	33·7
Weight of Grain per bushel in lb. :—					
Plot 1 ... ..	56·0	55·1	55·3	55·0	56·1
2 ... ..	54·3	55·3	55·0	56·9	55·5
3 ... ..	56·0				
4 ... ..	55·0				
Mean	55·3	55·2	55·2	56·0	55·8
Straw per acre in cwt. :—					
Plot 1 ... ..	26·1	32·0	31·6	24·1	29·3
2 ... ..	24·8	30·9	31·4	30·7	31·1
3 ... ..	28·9				
4 ... ..	27·7				
Mean	26·9	31·4	31·5	27·4	30·2
Total Produce per acre in lb. :—					
Plot 1 ... ..	4975	5748	5893	4760	5168
2 ... ..	4635	5550	5690	5610	5560
3 ... ..	5275				
4 ... ..	4885				
Mean	4943	5649	5792	5185	5364
Percentage increase in grain over the unmanured plots ... ..					
	—	10·5	17·2	7·8	1·5

### Residual effect of Dung stored in different ways.

Manure applied at the rate of 20 tons of original manure per acre for Potatoes in 1916, followed in 1917 by Wheat without manure.

#### Foster's Field.

POTATOES.	No Manure.	Compact heap in the open.	Compact heap in the open, covered with soil.	Compact heap under cover.
Weight of Potatoes. Tons per Acre	2.63	3.65	3.91	4.00
Percentage increase over Unmanured Plots ... ..	—	39	48	52
WHEAT.				
Weight of Grain per Acre ... lb.	1349	1637	1572	1752
Weight of Straw per Acre ... lb.	1870	2135	1965	2315
Weight of total Produce per Acre lb.	3219.	3772	3537	4067
Bushels of Grain per Acre ... ..	19.95	24.55	23.25	25.55
Percentage increase over Unmanured	—	23	16	28

### EXPERIMENTS WITH VARIOUS NITROGENOUS MANURES.

#### Potatoes. Great Knott Wood Field, 1916.

Plot.	All plots received per Acre : Dung, 10 tons ; Superphosphate, 2 cwt. ; Bone Flour, 2½ cwt.	Weight of Potatoes per Acre.
	Additional Manure per Acre :—	Tons.
1	} Nitrolim, 1 cwt. ... ..	5.45
2		5.20
3	No additional manure ... ..	4.80
4	} Sulphate of Ammonia, 1 cwt. ... ..	5.00
5		4.54

#### Savoys. Great Knott Wood Field, 1916.

Plot.	All plots received per Acre : Dung, 10 tons ; Superphosphate, 2 cwt. ; Bone Flour, 1½ cwt. ; Salt, 1½ cwt.	No. of plants per Acre.	Weight of produce per Acre.
1	} Received an additional dressing of Nitrolim at {	10600	Tons. 13.04
2		12300	14.64
3		11800	11.52

#### Savoys. Little Knott Wood Field, 1917.

Plot.	All plots received per Acre : Dung, 10 tons, and Superphosphate, 2½ cwt.	No. of plants per Acre.	Weight per Acre.
	Additional Manure per Acre :—		Tons.
1	Sulphate of Ammonia, 2 cwt. ... ..	10380	15.24
2	Nitre Cake Sulphate of Ammonia, 2 cwt. ... ..	10450	14.59
3	No additional manure ... ..	10160	11.56
4	Decomposed Cordite, 275 lb. ... ..	11350	12.32



EXPERIMENTS IN SOIL MANAGEMENT.  
CHALKING.  
Sawpit Field.

	Chalked in 1913.		Not Chalked.	
	20 loads per Acre Carted (1).	50 loads per Acre Dug (2).		
1914, OATS (Grey Winter—Yield per acre bush.	37·3	41·1	—	44·6
1915, CLOVER—Yield per acre as Hay cwt.	35·8	39·2	20·2	18·6
1916, WHEAT—Yield per acre Dressed Grain bush.	33·8	30·2	24·2	31·3
Weight per bushel ... lb.	62·0	63·3	62·4	63·0
Straw per acre ... cwt.	40·3	35·0	30·5	35·5
Total Produce per acre ... lb.	6878	6130	5163	6246
1917, OATS—Yield per acre Dressed Grain bush.	29·7	27·1	23·6	28·3
Weight per bushel ... lb.	33·3	36·4	36·8	35·3
Straw per acre ... cwt.	22·8	22·9	23·2	23·6
Total Produce per acre ... lb.	3842	3804	3675	3895

1, Chalk carted from Harpenden New Sewage beds, February, 1913.  
2, Chalk dug on Sawpit Field, November, 1912, to March, 1913, and spread as dug.  
Journal of Board of Agriculture, October 1916 (Vol. XXIII, No. 7, page 625) gives a detailed account of the method and cost of Chalking.

Great Harpenden Field.

		1914.	1915.							
		POTATOES (Dalhousie). Tons per Acre.	BARLEY. (Plumage Cross).				WHEAT. (Squareheads Master).			
			Dressed Grain.		Straw per Acre.	Total produce per Acre.	Dressed Grain.		Straw per Acre.	Total produce per Acre.
			Yield per Acre. Bush.	Weight per Bushel. lb.			Yield per Acre. Bush.	Weight per Bushel. lb.		
Unchalked ...	9·3	}	W 31·9	55·6	17·0	3788	20·2	62·0	19·2	3584
Chalked in 1913 ...	8·8		E 40·5	55·9	21·2	4784	21·6	61·5	23·7	4228
(about 20 loads per Acre)			W 31·9	55·5	18·8	4025	21·7	62·0	20·8	3859
			E 35·9	54·9	19·0	4212	17·6	62·0	20·1	3525
1916.			WHEAT. (Wilhelmina).				WINTER OATS.			
Unchalked ...	...		31·7	59·5	39·7	6631	29·4	42·2	20·5	3661
Chalked in 1913 ...	...		27·3	59·0	37·3	6100	33·3	45·0	19·5	3850
							40·7	43·0	26·3	4825
							36·3	44·4	25·0	4553
1917.			WHEAT. (Red Standard).				WHEAT. (Squarehead's Master).			
Unchalked ...	...		24·2	61·2	18·7	3809	22·2	59·7	17·9	3539
Chalked in 1913 ...	...		24·9	61·2	18·4	3801	19·1	58·8	15·9	3079
			27·7	60·8	20·1	4138	23·2	60·0	18·0	3643

W = West portion of plot. E = East portion.

## EXPERIMENTS IN SUBSOILING.

### Potatoes. Great Knott Wood Field, 1916.

Plot.	All plots received per Acre : Dung, 10 tons Superphosphate, 2 cwt. Bone Flour, 2½ cwt. Sulphate of Ammonia, 1 cwt.	Weight of produce per Acre.
4	} Not subsoiled ... .. {	Tons.
5		5·00
6		4·54
7		5·27
	} Subsoiled for this crop ... .. {	5·50

### Great Harpenden Field.

1914.		1915.			1916.			1917.		
POTA- TOES. (King Ed- ward VII)		WHEAT. (Squarehead's Master).			WINTER OATS.			WHEAT. (Square- heads Master).		
Tons per Acre.		1	2	3	1	2	3	1	2	
Sub- soiled in 1914	7·4 (mean of 4 plots)	Dressed Grain per Acre ... Bus.	20·3	19·4	16·4	30·5	30·9	29·3	19·0	21·5
		Weight per Bus. lb.	62·0	61·8	61·3	42·9	43·6	43·8	58·5	59·1
		Straw per Acre cwt.	20·8	18·8	21·2	20·9	21·1	23·0	16·2	17·9
		Total produce per Acre ... lb.	3775	3478	3541	3808	3863	4075	3123	3450
Not sub- soiled	6·9 (mean)	Dressed Grain per Acre ... Bus.	19·1	15·5	13·7	29·4	33·3	22·2	19·1	
		Weight per Bus. lb.	61·8	62·3	62·0	42·2	45·0	59·7	58·8	
		Straw per Acre cwt.	21·0	16·3	15·8	20·5	19·5	17·9	15·9	
		Total produce per Acre ... lb.	3709	2938	2788	3661	3850	3539	3079	

### Wheat after Fallow (without Manure, 1851, and since).

#### Hoos Field, 1915, 1916 and 1917.

			1915.	1916.	1917.	Average 61 years, 1856-1916.
Dressed Grain	...	(Yield—Bush. per Acre	7·1	8·8	6·6	15·6
		Weight per Bushel lb.	59·8	60·2	59·4	59·5
Straw ...	...	cwt. per Acre ...	8·4	7·3	7·8	13·4
Total produce	...	lb. per Acre ...	1462	1475	1346	2477

### COMPARISON IN VARIETIES OF WHEAT, 1917. Great Harpenden Field.

	Red Standard.		Squareheads Master.		Red Marvel.	
Dressed Grain per Acre Bush.	24.2	24.9	22.2	19.1	25.2	28.3
Weight per Bushel ... lb.	61.2	61.2	59.7	58.8	59.3	60.6
Straw per Acre ... cwt.	18.7	18.4	17.9	15.9	19.5	22.2
Total produce per Acre lb.	3809	3801	3539	3079	3830	4354

### METHODS of SOWING WHEAT after POTATOES.

	Produce per Acre.	Land ploughed Wheat then sown in usual way		Wheat ploughed in after being		Drilled on potato tilth, not plough'd
		Seed drilled.	Seed broad- casted.	De- posited by drill.	Broad- casted.	
Great Harpenden Field 1915. WHEAT. Squareheads Master.	Dressed Grain Bush.	24.6	25.0	24.0	24.6	23.4
	Weight per Bushel ... lb.	62.9	61.9	62.6	62.8	62.6
	Straw per Acre cwt.	21.1	23.4	19.7	21.4	20.3
	Total produce ... lb.	4084	4329	3855	4121	3898
West Barn Field, 1916. RIVETTS WHEAT.	Dressed Grain Bush.	46.1	42.9	36.9	40.4	37.9
	Weight per Bushel ... lb.	59.5	59.4	59.1	59.4	59.4
	Straw per Acre cwt.	34.4	35.7	28.7	32.1	29.8
	Total produce ... lb.	6722	6634	5479	6088	5690
Foster's Field, 1917. WHEAT. Red Standard.	Dressed Grain Bush.	23.9	—	13.8	—	—
	Weight per Bushel lb.	25.1	—	11.0	—	—
	Straw per Acre cwt.	60.5	—	60.0	—	—
	Total produce lb.	60.8	—	59.8	—	—
		23.8	—	14.1	—	—
	22.0	—	7.6	—	—	
	4286	—	2525	—	—	
	4124	—	1588	—	—	

### PLOUGHED UP GRASSLAND.

#### New Zealand Field, 1916.

This field had been pasture land for 8 years and was ploughed up in autumn of 1915.  
No manure was given.

	Produce per Acre.		Produce per Acre.
Potatoes—"King Edward"	Tons. 1.34	Mangolds	Tons. 12.25
"Dalhousie"	1.07	"	8.98
Beans—Crop failed, the seed being taken by birds.			
	Yield per Acre. Grain.	Weight per Bush. Grain.	Weight per Acre. Straw.
Wheat* ...	14.41 bush.	56.3 lb.	26.1 cwt.
Barley ...	27.6 bush.	55.0 lb.	20.9 cwt.
Winter Oats ...	45.7 bush.	40.2 lb.	30.9 cwt.
Spring Oats ...	37.9 bush.	37.0 lb.	21.5 cwt.
Total Produce per Acre.			
	4011 lb.		3981 lb.
			5553 lb.
			3957 lb.

1917.—Barley was sown over the whole field and gave a yield of 24.8 bush. per acre.

\* Crop attacked by birds.