


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



## Report 1921-22 With the Supplement to the Guide to the Experimental Plots Containing the Yields per Acre Etc.



[Full Table of Content](#)

---

### Table of Results - Later Experiments

#### Rothamsted Research

Rothamsted Research (1923) *Table of Results - Later Experiments* ; Report 1921-22 With The Supplement To The Guide To The Experimental Plots Containing The Yields Per Acre Etc., pp 90 - 104 - DOI: <https://doi.org/10.23637/ERADOC-1-110>

### ROTATION PLOTS.

Little Hoos Field, 1921 and 1922.

Arranged to test the RESIDUAL VALUE of VARIOUS MANURES in year of application and one, two, and three years after.  
Produce per acre.

Plot.	Manure per Acre from 1919 onwards.	Year of Dressing.	1921 (18th Season), Barley.					1922 (19th Season), Barley.				
			Dressed Grain.		Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.	Dressed Grain.		Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.		
			Yield per Acre.	Weight per Bush.			Yield per Acre.	Weight per Bush.				
A 1	Control	—	Bush. 19.6	lb. 57.1	64.0	10.0	107.7	Bush. 20.9	lb. 51.6	82.0	10.6	96.1
A 2	...	1920	39.4	58.8	1512	19.9	107.4	[38.4]	52.9	1404	[19.2]	[98.9]
A 3	Ordinary Dung, 16 tons	1921	<b>37.7</b>	<b>58.9</b>	<b>1388</b>	<b>18.5</b>	<b>110.0</b>	35.7	53.0	1284	16.5	106.5
A 4	...	1922	25.9	58.0	844	13.5	103.8	<b>38.2</b>	<b>52.7</b>	<b>1372</b>	<b>17.8</b>	<b>105.9</b>
A 5	...	1915	31.2	57.4	1020	16.6	101.2	32.9	53.1	1232	15.3	106.7
B 1	Cake fed dung, 16 tons	1920	42.5	58.2	1804	21.3	107.7	38.1	52.7	1384	18.8	99.6
B 2	Control ...	—	18.8	56.6	688	10.4	97.3	26.9	52.8	1001	13.1	101.6
B 3	...	1921	<b>40.4</b>	<b>58.6</b>	<b>1628</b>	<b>20.0</b>	<b>108.9</b>	[39.7]	52.8	1484	[20.2]	[97.0]
B 4	Cake fed dung, 16 tons	1922	32.2	57.4	1236	16.3	106.2	[44.2]	<b>53.2</b>	<b>1660</b>	[22.5]	[97.5]
B 5	...	1915	30.9	57.8	1196	15.7	105.8	35.7	53.0	1364	17.3	102.0
C 1	Shoddy; Superphosphate; Sulphate of Potash ...	1920	22.0	56.0	940	12.1	99.9	16.6	51.9	696	9.9	86.8
C 2	...	1921	<b>30.8</b>	<b>56.3</b>	<b>1276</b>	<b>16.0</b>	<b>106.0</b>	19.2	52.5	816	11.2	87.0
C 3	Control ...	—	19.7	56.0	880	11.2	96.9	22.8	52.5	918	11.8	96.1
C 4	...	1922	21.1	56.0	1164	13.8	84.1	<b>39.9</b>	<b>52.5</b>	<b>1448</b>	<b>18.4</b>	<b>105.3</b>
C 5	Shoddy; Superphosphate; Sulphate of Potash ...	1919	23.2	56.4	1068	13.2	95.4	28.4	52.9	1132	14.5	97.6

D	1	Guano; Sulphate of Ammonia; Sulphate of Potash	1920	28.4	56.0	118	1252	15.4	99.3	17.9	52.5	80	732	10.7	85.0
	2		1921	<b>27.9</b>	<b>55.7</b>	<b>188</b>	<b>1348</b>	<b>17.1</b>	<b>90.8</b>	21.0	52.6	71	924	12.1	86.9
	3		1922	18.7	56.3	112	888	11.6	89.9	<b>35.6</b>	<b>53.0</b>	<b>113</b>	<b>1408</b>	<b>19.0</b>	<b>93.9</b>
	4		—	14.7	54.5	122	772	9.9	83.5	17.9	51.8	87	788	12.2	74.5
	5		1919	19.1	54.9	131	992	12.6	83.7	18.4	52.1	84	824	11.9	78.0
E	1	Rape Dust; Superphosphate; Sulphate of Potash	1920	25.5	55.8	111	1092	13.4	102.5	19.4	50.6	83	872	11.0	86.6
	2		1921	<b>36.1</b>	<b>54.7</b>	<b>152</b>	<b>1524</b>	<b>17.3</b>	<b>109.8</b>	25.5	53.6	70	1068	13.8	93.1
	3		1922	13.1	56.0	123	588	7.8	97.6	<b>34.8</b>	<b>52.6</b>	<b>85</b>	<b>1328</b>	<b>17.0</b>	<b>100.8</b>
	4		1919	15.4	55.8	127	688	9.4	93.3	16.5	50.9	75	764	10.2	79.9
	5		—	21.4	55.9	134	872	11.9	99.8	19.7	53.1	63	824	11.8	84.2
F	1	Control	—	10.3	52.3	84	476	7.1	78.1	20.3	52.3	87	1048	13.1	77.8
	2		1920	22.1	55.5	94	896	11.9	98.5	23.6	52.6	73	960	12.1	96.9
	3		1921	<b>33.5</b>	<b>55.9</b>	<b>177</b>	<b>1480</b>	<b>17.8</b>	<b>102.9</b>	23.4	53.8	68	992	12.8	92.5
	4		1922	17.6	53.0	115	808	10.5	89.5	<b>37.8</b>	<b>52.5</b>	<b>97</b>	<b>1616</b>	<b>19.7</b>	<b>94.5</b>
	5		1919	16.4	55.5	122	896	11.7	78.7	19.6	52.0	70	892	12.0	81.3
G	1	Bone Meal; Sulphate of Ammonia; Sulphate of Potash	1920	20.4	53.7	115	844	11.6	93.5	22.6	53.7	96	1076	14.3	81.9
	2		1921	<b>24.1</b>	<b>54.6</b>	<b>123</b>	<b>1208</b>	<b>14.1</b>	<b>90.8</b>	28.0	52.8	76	1228	15.1	91.7
	3		—	22.9	54.9	93	1068	13.9	86.4	29.2	52.8	69	1292	15.3	93.9
	4		1922	22.8	54.5	91	1004	12.6	94.1	<b>36.2</b>	<b>53.8</b>	<b>103</b>	<b>1696</b>	<b>20.6</b>	<b>88.6</b>
	5		1919	20.8	53.8	100	944	13.2	82.3	25.5	52.3	79	1208	14.6	86.3
H	1	Basic Slag; Sulphate of Ammonia; Sulphate of Potash	1920	30.8	56.8	88	1184	14.8	111.2	28.1	52.7	87	1148	14.4	97.1
	2		1921	<b>34.1</b>	<b>55.8</b>	<b>150</b>	<b>1672</b>	<b>20.1</b>	<b>91.0</b>	29.5	53.2	82	1292	15.6	94.4
	3		1922	24.6	56.5	91	1040	13.9	94.9	<b>40.3</b>	<b>53.2</b>	<b>100</b>	<b>1844</b>	<b>22.0</b>	<b>90.8</b>
	4		1919	30.3	55.9	88	1236	15.2	104.5	30.4	53.8	64	1252	15.1	100.7
	5		—	27.8	55.4	106	1301	16.3	90.3	20.8	53.4	82	1020	12.5	85.6

NOTES.—Since 1919 the manure for each plot (except of series A and B) has been rationed at 40 lb. Nitrogen, 100 lb. Calcium Phosphate, and 50 lb. Potash per acre. Each plot has been supplied with as much of its particular manure (shoddy, guano, &c.) as possible without exceeding the receipt in any of the three rationed ingredients. Any deficit in either of these three has then been made good by adding the necessary quantity of Sulphate of Ammonia, Superphosphate, or Sulphate of Potash. Figures in italics denote unmanured plots. The yields on the plots to which the manure was applied in a given year are printed in heavy type. Figures in square brackets are estimated yields.

### STRAW EXPERIMENT, 1921. Potatoes (Arran Chief). Sawpit Field.

Manure per Acre.	Yield per Acre.		
	1st Plot	2nd Plot	3rd Plot
8 tons Rotted Straw Manure—Single Nitrogen ...	Tons 2.30	Tons 2.18	Tons 1.96
16 " " " " " " " " ...	2.48	2.63	2.16
32 " " " " " " " " ...	1.73	2.39	2.29
2 cwt. Sulphate of Ammonia ... ..	1.20	1.48	1.13
4 " " " " " " " " ...	1.66	1.57	1.48
8 " " " " " " " " ...	1.52	1.71	1.38
16 " " " " " " " " ...	1.41	1.55	1.27
8 tons Rotted Straw Manure—Double Nitrogen ...	2.09	2.20	1.86
16 " " " " " " " " ...	3.32	2.59	2.50
32 " " " " " " " " ...	2.16	2.68	2.04
Control—No Manure ... ..	1.39	1.61	1.41
" " " " " " " " ...	1.52	1.45	1.39

Single Nitrogen represents 1 cwt. Sulphate of Ammonia added to 1 ton of straw.  
Double Nitrogen represents 2 cwt. Sulphate of Ammonia added to 1 ton of straw.

### RESIDUAL VALUE OF SLUDGE, 1921. Long Hoos Field.

Treatment of Plots in 1920. Manure per Acre.	Dressed Grain.				Offal Grain per Acre. lb.		Straw per Acre.				Proportion of Total Grain to 100 of Total Straw.	
	Yield per Acre. Bush.		Weight per Bushel. lb.				Straw. lb.		Total Straw. cwt.			
	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.	1st Plot.	2nd Plot.
<b>1921, Wheat (Red Standard) after Potatoes (1920).*</b>												
Activated Sewage Sludge, 13.3 tons	29.8	27.9	64.0	64.1	371	406	2925	2624	32.7	30.8	62.2	63.6
Farmyard Dung 15 tons ...	34.8	31.6	64.0	64.1	296	371	2461	2600	30.3	29.7	74.5	72.0
Control ... ..	26.0	26.9	63.3	63.0	342	325	2299	1997	26.2	26.6	67.6	67.7
<b>1921, Wheat (Red Standard) after Barley (1920).†</b>												
Sulph. of Ammonia 1.45 cwt. ...	24.1		63.0		387		2738		31.1		54.6	
Activated Sewage Sludge, 2.7 tons	30.1		63.0		351		2857		31.4		64.1	
Control ... ..	27.2		62.5		405		2738		29.4		63.9	
Control ... ..	27.4		63.0		435		2333		30.3		63.7	

\* In 1920 this set received a basal dressing of 6 cwt. Super. and 1 cwt. Nitrate of Ammonia per acre. No manure was given in 1921.

† In 1921 this set was manured as farm, viz., 1 cwt. Sulphate of Ammonia and 1 cwt. Superphosphate per acre.

TOP DRESSING EXPERIMENTS.

Treatment of Plots and Quantities per Acre.	Dressed Grain									Offal Grain per Acre.						Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.							
	Yield per Acre Bushels.			Weight per Bushel lb.			Plot			Plot			Plot			Plot			Plot			Plot			Plot				
	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot	Plot		
	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3		
<b>Oats (Grey Winter)*</b>																													
<b>Great Knott Field, 1921.</b>																													
Super. 1½ cwt. ...	52.1	57.4	54.7	44.1	43.0	43.6	42.0	49.0	46.0	19.00	26.00	25.40	31.8	37.5	40.2	76	71	63	...	...	...	...	...	...	...	...	...	...	
Super. 1½ cwt.; Sul./Amm. 1½ cwt., applied March 4 ...	45.0	43.6	44.2	43.3	45.0	43.0	38.0	48.5	38.0	20.20	19.20	26.20	31.6	35.5	37.9	66	61	54	...	...	...	...	...	...	...	...	...	...	
Super. 1½ cwt.; Mur./Amm. 1.44 lb., applied March 4 ...	48.4	57.3	44.0	44.0	44.0	44.6	43.5	59.0	48.5	22.80	29.80	24.80	37.1	47.0	36.4	62	59	60	...	...	...	...	...	...	...	...	...	...	
No Manure ...	47.1	48.6	47.9	43.6	44.0	43.5	41.0	47.5	42.0	20.60	23.40	22.00	31.1	36.1	33.4	71	65	67	...	...	...	...	...	...	...	...	...	...	
<b>Wheat (Red Standard).</b>																													
<b>Great Harpenden Field, 1921.</b>																													
Super. 200 lb.; Sul./Amm. 100 lb., applied March 2-3 ...	19.4	16.9	—	62.5	62.5	—	56.5	43.0	—	23.50	16.60	—	30.0	22.9	—	53	58	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Sul./Amm. 200 lb., applied March 2-3 ...	20.3	15.4	—	62.5	62.5	—	54.0	44.3	—	20.60	19.00	—	27.1	25.6	—	60	49	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Sul./Amm. 100 lb., applied March 2-3 and Sul./Amm. 100 lb., applied May 2 ...	21.3	13.8	—	62.5	62.5	—	49.5	35.0	—	20.80	14.50	—	27.2	20.7	—	60	52	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Mur./Amm. 86 lb., applied March 2-3 ...	17.7	15.7	—	62.0	62.0	—	43.0	37.0	—	19.60	17.00	—	24.4	22.8	—	56	53	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; N./Soda 140 lb., applied March 2-3 ...	15.2	16.9	—	62.5	62.0	—	35.5	41.5	—	17.50	19.80	—	21.4	24.0	—	54	55	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Sul./Amm. 100 lb., applied May 2 ...	23.1	16.1	—	62.3	62.0	—	50.0	38.5	—	26.60	18.00	—	30.4	21.9	—	57	56	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Sul./Amm. 200 lb., applied May 2 ...	23.8	17.4	—	62.5	62.5	—	50.0	48.5	—	26.20	19.00	—	30.6	22.6	—	58	62	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Mur./Amm. 86 lb., applied May 2 ...	17.5	18.2	23.6	62.0	62.0	63.0	33.8	40.5	46.5	17.00	17.10	25.00	21.3	22.6	27.8	60	61	63	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb.; Sul./Amm. 100 lb., applied April 4 ...	17.1	—	—	62.5	—	—	36.5	—	—	17.80	—	—	24.0	—	—	53	—	—	...	...	...	...	...	...	...	...	...	...	
No Manure ...	13.5	17.6	—	63.5	61.5	—	37.5	36.8	—	13.20	18.80	—	17.6	22.2	—	63	58	—	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb. ...	19.0	12.2	22.0	62.5	62.5	62.5	41.5	38.5	52.5	18.30	12.60	25.40	22.7	17.1	28.1	63	60	60	...	...	...	...	...	...	...	...	...	...	
Super. 200 lb. ...	—	16.6	—	—	61.0	—	—	36.0	—	—	18.40	—	—	22.6	—	—	54	—	—	...	...	...	...	...	...	...	...	...	
<b>Wheat (Red Standard)†</b>																													
<b>Foster's Field, 1922.</b>																													
1 cwt. Sul./Amm. applied March 18 ...	14.4	18.4	18.5	59.5	60.0	60.0	42.5	39.1	40.0	14.25	16.50	16.50	17.0	19.2	19.9	67	70	68	...	...	...	...	...	...	...	...	...	...	...
1 cwt. Sul./Amm., applied April 20 ...	18.3	21.3	15.8	59.5	60.3	60.0	39.4	39.4	35.3	17.50	18.50	14.75	21.2	23.0	17.6	62	65	66	...	...	...	...	...	...	...	...	...	...	...
1 cwt. Sul./Amm., applied May 18 ...	17.2	21.9	16.4	60.0	60.5	60.3	37.2	40.9	40.6	16.25	17.50	14.25	19.4	21.4	17.9	65	72	70	...	...	...	...	...	...	...	...	...	...	...
93 lb. Mur./Amm., applied March 18 ...	17.8	19.6	16.4	60.0	60.3	60.5	36.2	40.0	30.0	15.50	18.75	13.75	19.2	22.8	18.0	66	62	64	...	...	...	...	...	...	...	...	...	...	...
186 lb. Mur./Amm., applied March 18 ...	19.7	21.2	20.2	59.5	60.0	60.3	35.0	40.9	36.2	17.00	16.50	18.00	21.0	20.8	21.7	65	72	65	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Sul./Amm., applied March 18 ...	20.0	20.8	18.3	60.3	60.0	60.0	43.8	50.3	36.9	18.00	19.50	17.25	22.5	25.2	22.1	65	62	59	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Sul./Amm., applied April 20 ...	20.1	18.4	13.2	60.8	60.0	55.5	39.4	52.5	36.9	17.00	17.75	14.75	21.4	24.3	19.0	67	60	52	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Sul./Amm., applied May 18 ...	19.4	17.9	17.2	60.0	60.5	60.0	42.5	50.6	44.1	17.00	16.75	15.25	21.2	23.0	20.1	67	62	65	...	...	...	...	...	...	...	...	...	...	...
Basal Manuring only ...	13.4	15.0	11.9	59.8	60.0	60.0	32.8	39.1	29.7	12.50	13.25	12.25	15.2	17.4	14.7	67	66	61	...	...	...	...	...	...	...	...	...	...	...
No Manure ...	12.9	16.7	13.2	60.0	60.3	59.5	37.8	32.8	33.1	13.25	15.25	13.50	15.6	18.5	16.5	66	64	60	...	...	...	...	...	...	...	...	...	...	...
<b>Barley (Plumage Archer).</b>																													
<b>Long Hoos, 1922.</b>																													
2 cwt. Super.; 1 cwt. Sul./Amm., applied May 9 ...	34.5	30.1	—	51.3	51.8	—	28.1	23.8	—	16.50	13.25	—	22.5	18.0	—	81	89	—	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Super.; 2 cwt. Sul./Amm., applied May 9 ...	31.2	35.3	—	51.8	51.6	—	36.9	28.7	—	16.00	17.75	—	21.4	23.0	—	83	82	—	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Super.; 98 lb. Mur./Amm., applied May 9 ...	27.7	30.4	—	51.0	51.3	—	25.6	26.3	—	14.75	15.00	—	18.5	23.0	—	77	82	—	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Super.; 1 cwt. Mur./Amm., applied May 9 ...	25.7	32.3	—	52.0	54.3	—	25.0	35.0	—	11.75	15.75	—	21.0	22.5	—	83	86	—	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Super.; 51½ lb. Urea, applied May 9 ...	32.1	34.7	—	51.8	52.3	—	28.7	35.6	—	15.75	16.75	—	16.3	17.2	—	77	83	—	...	...	...	...	...	...	...	...	...	...	...
2 cwt. Super. ...	23.8	26.6	—	51.0	52.3	—	18.4	20.0	—	11.75	12.50	—	16.5	18.5	—	77	83	—	...	...	...	...	...	...	...	...	...	...	...
No Manure ...	23.6	31.2	—	52.0	51.3	—	20.0	24.4	—	11.00	16.50	—	16.5	22.1	—	77	75	—	...	...	...	...	...	...	...	...	...	...	...

\* A luxuriant crop of mustard was ploughed in previous to the sowing of the Oats; hence the high yields on these plots.  
 † All plots (except the unmanured), received a basal dressing of 2 cwt. Super.; 1 cwt. S./Pot. per Acre.

Top Dressing Experiments—*contd.*

Root Crops. Great Harpenden Field, 1922.

Manuring per Acre.	Yield per Acre.	
	1st Plot. Tons.	2nd Plot. Tons.
<b>Potatoes (Kerr's Pink).</b>		
<b>Dunged Series: 15 tons Farmyard Dung per Acre—</b>		
Super. 4 cwt., Sul./Pot. 1½ cwt. ... ..	6·73	5·41
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 3 cwt. (half as Top Dressing) ... ..	7·92	9·17
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 1½ cwt. ...	7·91	8·06
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul. Amm. 4½ cwt.		
Super. 4 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 3 cwt. ...	10·54	9·62
(1½ cwt. as Top Dressing) ... ..	10·08	9·37
Super. 4 cwt., Sul./Pot. 1½ cwt., Mur./Amm. 290 lb.	10·66	10·74
<b>Undunged Series:</b>		
Super. 6 cwt., Sul./Pot. 2 cwt. ... ..	6·10	4·90
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 3 cwt. (half as Top Dressing) ... ..	7·99	7·89
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 1½ cwt. ...	6·98	7·75
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 4½ cwt. (1½ cwt. as Top Dressing) ... ..	9·60	8·36
Super. 6 cwt., Sul./Pot. 2 cwt., Sul./Amm. 3 cwt. ...	8·72	9·22
Super. 6 cwt., Sul./Pot. 2 cwt., Mur./Amm. 290 lb. ...	9·21	8·50
<b>Swedes (Hurst's Monarch).</b>		
589 lb. Slag,* 1 cwt. Sul./Pot. ... ..	{ R <b>25·13</b>	<b>28·24</b>
	{ L 3·04	4·29
589 lb. Slag,* 1 cwt. Sul./Pot., 2 cwt. Sul./Amm. (as Top Dressing) ... ..	{ R <b>27·48</b>	<b>30·65</b>
	{ L 3·82	4·87
589 lb. Slag,* 1 cwt. Sul./Pot., 10 tons Farmyard Dung ... ..	{ R <b>28·75</b>	<b>32·37</b>
	{ L 4·22	4·12
589 lb. Slag,* 1 cwt. Sul./Pot., 10 tons Farmyard Dung, 2 cwt. Sul./Amm. (as Top Dressing)	{ R <b>32·61</b>	<b>32·43</b>
	{ L 4·60	4·71

\* Equivalent to 5 cwt. Super. R = Roots. L = Leaves.



**Slag Experiments—contd.**  
**Clover. West Barnfield, 1921 and 1922.**

No. of Plot.	Treatment of Plots and Quantities per Acre.	1921				1922			
		Yield per Acre.		Dry Matter per Acre.		Yield per Acre.		Dry Matter per Acre.	
		Series A	Series B	Series A	Series B	Series A	Series B	Series A	Series B
1	High Grade Slag No. 12, 1170 lb.	cwt. 40.8	cwt. 40.4	lb. 3521	lb. 3567	cwt. 10.5	cwt. 16.1	lb. 941	lb. 1418
2	Open Hearth, High Soluble Slag No. 13, 1925 lb.	43.0	38.7	3629	3470	18.5	13.7	1644	1183
3	Open Hearth, Low Soluble Slag No. 14, 1930 lb.	42.4	40.4	3720	3567	18.6	15.6	1679	1374
4	Gafsa Phosphate, 750 lb.	41.7	39.4	3654	3502	17.6	18.3	1604	1681
C	No Manure	43.6	39.4	3812	3593	16.3	18.3	1486	1630
C	"	40.8	40.8	3563	3563	17.1	17.1	1490	1490

**Barley (Plumage Archer). Long Hoos Field, 1922.**

Treatment of Plots.	Dressed Grain.				Offal Grain per Acre.				Straw per Acre.		Proportion of Total Grain to Total Straw.				
	Yield per Acre in Bushels.		Weight per Bushel in lb.		per Acre.		lb.		Straw.		Total Straw.				
	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 1.	Slag No. 20.		
Basal Manuring, Slag, full quantity	36.0	26.0	51.4	51.3	197	162	231	1375	1175	1250	18.5	16.9	19.3	99	80
Basal Manuring, Slag, half quantity	35.1	31.7	52.0	52.8	200	172	213	1375	1238	1238	19.3	18.0	19.1	95	71
Gafsa Phosphate, 87 lb.	29.9	25.5	51.8	51.9	194	169	241	1238	1113	1438	18.5	16.9	20.3	84	79
	26.2	36.4	51.4	51.3	163	181	200	*875	1500	1275	*11.7	19.6	18.3	*115	93
Basal Manuring, Gafsa Phosphate, 174 lb.	34.8	25.2	52.3	51.6	231	162	209	1488	1088	1213	19.6	16.5	18.2	92	80
Basal Manuring only	25.7	33.7	52.0	52.0	178	178	203	1063	1363	1388	15.8	18.3	18.9	85	94
No Manure	30.1	24.7	50.8	51.1	188	203	228	1363	1100	1263	19.2	15.9	18.8	81	82

Basal Manuring is 1 cwt. Sulphate of Potash; 1 cwt. Sulphate of Ammonia and 436 lb. Slag No. 1 per acre. Full Quantity Slag represents 636 lb. Slag No. 20, 602 lb. Slag No. 2. \*There was a high wind blowing when this plot was threshed, hence the low figure for the yield of straw.



Slag Experiments—*contd.*

Swedes (Hurst's Monarch) Produce per Acre.  
Great Harpenden Field, 1922.

Manuring per Acre.	Roots.			Leaves.		
	Slag No. 20.	Slag No. 2.	Slag No. 1.	Slag No. 20.	Slag No. 2.	Slag No. 1.
	Tons.	Tons.	Tons.	Tons.	Tons.	Tons.
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag full quantity ...	25·92	27·92	30·40	4·89	3·82	4·16
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag full quantity ...	32·08	30·31	30·40	4·01	5·04	4·20
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag half quantity, Gafsa Phosphate, 175 lb. ...	27·19	28·04	31·88	4·18	3·53	4·10
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., Slag half quantity, Gafsa Phosphate, 175 lb. ...	28·21	29·78	28·82	4·28	4·16	4·27
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., No. 7 Nauru Phosphate, 262½ lb. ...	30·96	26·43	26·50	4·49	4·00	3·98
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt., No. 3 Gafsa Phosphate, 350 lb. ...	27·83	31·12	28·46	3·95	4·58	4·66
Sulphate Ammonia 2 cwt., Sulphate Potash 1 cwt. ...	27·21	31·45	25·74	4·16	5·02	3·99
No Manure ...	25·67	27·23	22·70	3·54	3·67	3·19

NOTE.—“Full Quantity” Slag is No. 20, 1275 lb. per Acre.  
No. 2, 1225 „ „  
No. 1, 875 „ „

Description of Slags Used.

No.	Type.	Total Phosphate as Ca <sub>3</sub> (PO <sub>4</sub> ) <sub>2</sub>	Solubility %
1	Open Hearth, L.G., H.S. ...	25·0	90·4
2	„ „ L.G., L.S. ...	18·0	35·7
8	Phosphate, Slag Mixture ...	53·1	25·5
12	Talbot Process, H.G., H.S. ...	37·3	80·7
13	Open Hearth, L.G., H.S. ...	22·7	91·5
14	„ „ L.G., L.S. ...	22·6	29·0
15	Talbot Process, H.G., H.S. ...	40·0	72·5
16	Open Hearth, L.G., H.S. ...	21·3	88·3
17	Bessemer, H.G., H.S. ...	42·5	77·2
18	Open Hearth, L.G., H.S. ...	20·8	67·0
19	„ „ L.G., L.S. ...	20·2	21·0
20	„ „ L.G., H.S. ...	17·2	78·8

L.G. = Low Grade. L.S. = Low Soluble.  
H.G. = High Grade. H.S. = High Soluble.

### POTASH EXPERIMENTS.

Manuring per Acre.	Dry Matter per Acre.			Yield per acre.		
	1st Plot	2nd Plot	3rd Plot	1st Plot	2nd Plot	3rd Plot
<b>Clover. West Barn Field, 1922.</b>						
Control ... ..	lb. 1369	lb. 1273	lb. 1507	cwt. 15·2	cwt. 15·7	cwt. 18·6
Sulphate of Potash, 210 lb. ... ..	1533	1929	2123	18·6	25·0	26·4
Cement Works' Dust, 511 lb. ... ..	1381	1710	1729	17·5	21·8	21·4

<b>Potatoes (Arran Chief). Sawpit Field, 1921.</b>						
With Dung, 12 tons per Acre.						
				Tons.	Tons.	Tons.
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 470 lb. Sylvénite ...				3·57	*3·15	3·71
3 cwt. Super., 1½ cwt. Sulphate Ammonia, ... ..				3·55	*3·18	3·72
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 1½ cwt. Sulphate Potash				3·67	4·27	3·88
3 cwt. Super., 1½ cwt. Sulphate Ammonia, 1½ cwt. Sulphate Potash, 95 lb. Sulphate Magnesium ... ..				*3·07	3·92	3·87
No Manure. Control ... ..				*2·28	3·48	3·18
3 cwt. Super., 1½ cwt. Sul. Amm., 1½ cwt. Muriate Potash ... ..				*2·31	4·24	3·97
3 cwt. Super., 1½ cwt. Sul. Amm., 1½ cwt. Muriate Potash, 84 lb. Sul. Magnesium ... ..				*2·43	3·90	4·15

Without Dung.						
4 cwt. Super., 2 cwt. Sul. Amm., 625 lb. Sylvénite ... ..				3·49	4·04	3·11
4 cwt. Super., 2 cwt. Sul. Amm. ... ..				1·43	1·48	1·15
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Sul. Potash ... ..				3·48	4·28	3·52
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Sul. Pot., 127 lb. Sul. Mag.				3·85	4·26	3·25
No Manure. Control ... ..				1·24	1·72	1·65
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Muriate Potash ... ..				4·15	4·20	4·00
4 cwt. Super., 2 cwt. Sul. Amm., 2 cwt. Muriate Potash, 111 lb. Sul. Magnesium ... ..				4·27	3·95	3·63

<b>Potatoes (Arran Chief). Sawpit Field, 1921.</b>						
4 cwt. Super., 2 cwt. Sulphate Ammonia, 232 lb. Sul. Potash ...				3·00	2·46	2·82
4 cwt. Super., 2 cwt. Sulphate Ammonia ... ..				1·16	0·98	0·89
4 cwt. Super., 2 cwt. Sulphate Ammonia, 5·4 cwt. Sylvénite ...				*1·93	3·36	3·04
Control. No Manure. ... ..				*0·73	1·10	1·16

<b>Potatoes (Kerr's Pink). Great Harpenden Field, 1922.</b>						
With Dung 15 tons per Acre.						
Basal Manuring (= Super. 4 cwt., Sul. Amm. 1·5 cwt. per Acre) ...				8·78	7·72	7·60
Sulphate Potash 183 lb. + Basal Manuring ... ..				9·49	9·72	9·45
Muriate Potash 148 lb. + Basal Manuring ... ..				9·22	9·60	8·82
Muriate Potash 148 lb. + Salt 497 lb. + Basal Manuring ... ..				9·84	9·49	9·14
Without Dung.						
Basal (= Super. 6 cwt., Sulphate Ammonia 2 cwt. per Acre) ...				2·11	2·75	2·57
Sulphate Potash 244 lb. + Basal ... ..				7·88	8·96	8·06
Muriate Potash 197 lb. + Basal ... ..				8·62	8·73	7·62
Muriate Potash 197 lb. + Salt 662 lb. + Basal ... ..				8·45	8·27	8·43
Muriate Potash 197 lb. Sulphate Magnesium, 344 lb. + Basal ...				8·68	8·90	7·62
Muriate Potash 197 lb. Salt 662 lb. + Basal ... ..				8·66	8·02	7·51
No Manure ... ..				3·23	2·87	2·83
Sulphate Potash 244 lb. Sulphate Magnesium 344 lb. + Basal ...				9·25	8·79	7·11
Cement Works' Dust 614 lb. + Basal ... ..				7·47	6·66	6·38
Sylvénite 541 lb. + Basal ... ..				8·38	7·92	6·90

\* On these plots the bouts were badly broken down due to extra hoeing on account of the growth of Wheatbind.

### Mangolds (Prizewinner Yellow Globe). Great Harpenden Field, 1922.

Produce per Acre.

Manuring per Acre.	Roots.		Leaves.	
	1st Plot Tons.	2nd Plot Tons.	1st Plot Tons.	2nd Plot Tons.
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt., Sulphate Potash 2 cwt. ... ..	17·64	14·12	5·57	5·13
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt. ... ..	10·45	11·61	4·73	4·94
No. 9 Slag 4 cwt., Sulphate Ammonia 2 cwt., Cement Works' Dust ... ..	18·75	18·25	5·61	5·96
No Manure ... ..	10·88		4·25	

## POTATOES.

Relative Effects of Sulphates and Chlorides on different varieties.

Great Harpenden Field, 1922.

Variety.	Dunged Series.						Undunged Series.					
	Actual Weight of Potatoes.			Average Weight per Plant.			Actual Weight of Potatoes.			Average Weight per Plant.		
	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.
	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.
Ajax ...	16	12 $\frac{3}{4}$	19 $\frac{3}{4}$	3'20	2'55	2'82	13 $\frac{1}{2}$	17 $\frac{3}{4}$	4	2'25	2'54	1'00
	24	21 $\frac{1}{4}$	12 $\frac{1}{4}$	4'00	3'04	1'75	17 $\frac{1}{4}$	18	4 $\frac{1}{2}$	2'46	3'00	0'64
	27	16 $\frac{3}{4}$	33	3'86	4'13	4'71	7 $\frac{1}{4}$	23	4	2'42	3'29	0'67
Arran Comrade	15 $\frac{3}{4}$	11 $\frac{3}{4}$	7 $\frac{1}{4}$	2'25	1'96	2'42	11 $\frac{1}{4}$	8 $\frac{3}{4}$	3 $\frac{1}{4}$	2'25	1'46	0'65
	10 $\frac{1}{4}$	10 $\frac{3}{4}$	13	2'56	2'15	2'17	11 $\frac{1}{4}$	11 $\frac{1}{4}$	1 $\frac{1}{2}$	1'88	2'25	0'30
	15 $\frac{1}{2}$	10 $\frac{1}{2}$	13	2'58	2'10	2'17	15	18	1 $\frac{3}{4}$	2'14	2'57	0'29
British Queen	19 $\frac{1}{4}$	19	19 $\frac{1}{4}$	3'21	2'71	2'75	10 $\frac{1}{4}$	13 $\frac{3}{4}$	7 $\frac{3}{4}$	1'46	1'96	1'11
	19 $\frac{3}{4}$	18 $\frac{3}{4}$	19 $\frac{1}{4}$	2'82	2'68	2'75	16 $\frac{3}{4}$	11 $\frac{1}{4}$	8 $\frac{1}{4}$	2'36	1'92	1'18
	26 $\frac{3}{4}$	25	23 $\frac{1}{4}$	3'82	4'17	3'32	11 $\frac{3}{4}$	15 $\frac{3}{4}$	3 $\frac{3}{4}$	1'96	2'63	0'75
Duke of York ...	7 $\frac{3}{4}$	11	11 $\frac{1}{4}$	1'11	1'57	1'61	9	8	1	1'80	1'60	0'33
	8 $\frac{3}{4}$	14	14	1'25	2'00	2'00	9 $\frac{1}{4}$	6 $\frac{1}{2}$	2 $\frac{1}{2}$	1'54	0'93	0'63
	13 $\frac{1}{2}$	10 $\frac{1}{2}$	14 $\frac{3}{4}$	2'25	1'75	2'46	6 $\frac{1}{4}$	10 $\frac{1}{4}$	1 $\frac{1}{4}$	1'04	1'46	0'42
Epicure ...	16 $\frac{1}{2}$	14 $\frac{3}{4}$	10	2'36	2'11	1'43	12 $\frac{1}{4}$	9 $\frac{3}{4}$	1 $\frac{3}{4}$	2'04	1'63	0'35
	11 $\frac{1}{2}$	13 $\frac{1}{2}$	13 $\frac{1}{2}$	1'64	1'93	2'25	13 $\frac{1}{2}$	13 $\frac{3}{4}$	3 $\frac{3}{4}$	1'93	1'96	0'54
	16	18 $\frac{1}{2}$	19 $\frac{1}{2}$	2'29	2'64	2'79	11 $\frac{3}{4}$	12 $\frac{1}{4}$	1	1'68	1'79	0'25
Great Scott ...	13 $\frac{1}{2}$	19 $\frac{1}{2}$	21 $\frac{1}{2}$	3'38	2'79	3'07	21 $\frac{1}{2}$	17 $\frac{1}{4}$	4 $\frac{1}{2}$	3'07	2'46	0'75
	21 $\frac{1}{2}$	24 $\frac{1}{4}$	19 $\frac{1}{2}$	3'07	3'54	3'25	11 $\frac{3}{4}$	12 $\frac{3}{4}$	1 $\frac{1}{4}$	1'96	2'13	0'42
	27 $\frac{1}{4}$	29	24 $\frac{1}{2}$	3'89	4'14	3'50	14 $\frac{1}{4}$	13 $\frac{1}{4}$	1	2'38	2'65	0'50
Iron Duke ...	24	20	21	3'43	3'33	3'50	16 $\frac{3}{4}$	19 $\frac{1}{4}$	4 $\frac{3}{4}$	2'79	2'75	0'68
	21	18 $\frac{1}{2}$	16 $\frac{1}{4}$	3'00	3'08	2'32	10 $\frac{3}{4}$	20 $\frac{1}{4}$	4	1'79	2'89	1'00
	23 $\frac{3}{4}$	23 $\frac{1}{4}$	23	3'96	3'32	3'29	20	13	4 $\frac{1}{2}$	2'86	3'25	0'64
K. of K. ...	26	23 $\frac{3}{4}$	20 $\frac{1}{4}$	3'71	3'39	2'89	21 $\frac{1}{2}$	18 $\frac{1}{2}$	7 $\frac{1}{4}$	3'07	3'08	1'04
	28 $\frac{1}{2}$	27 $\frac{3}{4}$	21	4'07	4'63	4'20	—	—	—	—	—	—
	29 $\frac{1}{2}$	29 $\frac{1}{2}$	30 $\frac{1}{4}$	4'21	4'21	4'32	19 $\frac{1}{4}$	15 $\frac{1}{2}$	5 $\frac{3}{4}$	2'75	3'10	0'82
Kerr's Pink ...	18 $\frac{1}{4}$	20 $\frac{3}{4}$	12	3'04	2'96	2'00	18 $\frac{3}{4}$	20 $\frac{3}{4}$	6 $\frac{1}{2}$	2'68	2'96	0'93
	25	22 $\frac{1}{4}$	15	3'57	3'18	3'00	11 $\frac{1}{2}$	19 $\frac{1}{2}$	3 $\frac{1}{4}$	1'92	3'90	0'46
	26 $\frac{3}{4}$	30 $\frac{1}{4}$	15 $\frac{1}{2}$	3'82	4'32	3'88	24 $\frac{1}{4}$	22	5	3'46	3'14	0'71
Nithsdale ...	18	14 $\frac{1}{4}$	30 $\frac{3}{4}$	2'57	2'04	1'96	9	9 $\frac{1}{2}$	1 $\frac{1}{4}$	1'29	1'58	0'42
	15 $\frac{1}{2}$	20 $\frac{1}{2}$	20	2'21	2'93	2'86	12	15	1	2'00	2'14	0'33
	21 $\frac{1}{2}$	26	14 $\frac{1}{4}$	3'58	3'71	3'56	14 $\frac{1}{4}$	14 $\frac{1}{2}$	2 $\frac{1}{2}$	2'04	2'07	0'63
Tin Perfection	20 $\frac{3}{4}$	17	12 $\frac{3}{4}$	3'46	2'83	2'55	20	19 $\frac{1}{2}$	7 $\frac{1}{2}$	2'86	2'79	1'07
	21 $\frac{3}{4}$	20 $\frac{3}{4}$	23 $\frac{3}{4}$	3'11	2'96	3'39	18 $\frac{3}{4}$	17 $\frac{3}{4}$	8 $\frac{3}{4}$	2'68	2'54	1'21
	17 $\frac{1}{2}$	19 $\frac{1}{4}$	23 $\frac{1}{2}$	2'50	3'21	3'36	21 $\frac{3}{4}$	17	7	3'11	2'83	1'00
Up-to-Date ...	25 $\frac{3}{4}$	23 $\frac{3}{4}$	25 $\frac{1}{4}$	4'29	3'39	4'21	26 $\frac{3}{4}$	20 $\frac{1}{4}$	9 $\frac{1}{4}$	3'82	2'89	1'32
	20 $\frac{1}{2}$	25 $\frac{3}{4}$	25 $\frac{1}{2}$	2'93	3'68	3'64	20 $\frac{1}{2}$	14 $\frac{1}{4}$	8 $\frac{1}{4}$	2'93	2'38	1'18
	29 $\frac{3}{4}$	28 $\frac{3}{4}$	28 $\frac{3}{4}$	4'25	4'07	4'11	21 $\frac{3}{4}$	21	11	3'11	3'00	1'83

NOTE.—7 Plants were set in each Row.

Manures were:—Dunged Series: Basal Row: Super. 4 cwt.; Sulphate of Ammonia 1 $\frac{1}{2}$  cwt.;  
 Dung 15 tons per Acre.  
 Sulphate Row: Basal Manuring; Sulphate of Potash 184 lb. per Acre.  
 Chloride Row: Basal Manuring; Muriate of Potash 147 lb. per Acre.  
 Undunged Series: Basal Row: Super. 6 cwt.; Sulphate of Ammonia 2 cwt. per Acre.  
 Sulphate Row: Basal Manuring; Sulphate of Potash 244 lb. per Acre.  
 Chloride Row: Basal Manuring; Muriate of Potash 197 lb. per Acre.

Potatoes. Great Harpenden Field, 1922.  
Comparison of Varieties.

	Ajax.	Arran Comrade.	British Queen.	Duke of York.	Epicure.	Great Scott.	Iron Duke.	K. of K.	Kerr's Pink.	Nithsdale.	Tin Perfection.	Up-to-Date
	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.	lb.
Average weight of Potatoes lifted per row	16.21	10.54	16.08	8.86	11.88	16.58	16.89	21.62	17.63	13.49	17.49	21.47
Average weight per plant ...	2.70	1.90	2.43	1.50	1.81	2.79	2.67	3.28	2.73	2.23	2.62	3.17

Comparison of Manurial Treatment.

	Dunged Series.			Undunged Series.		
	Sulphate Row.	Chloride Row.	Basal Row.	Sulphate Row.	Chloride Row.	Basal Row.
	lb.	lb.	lb.	lb.	lb.	lb.
Average weight of Potatoes lifted per row	20.12	19.82	18.62	15.17	15.41	4.40
Average weight per plant ...	3.08	3.03	2.94	2.36	2.37	0.80

PROFESSOR BLACKMAN'S  
ELECTRO CULTURE EXPERIMENTS.

Clover. Great Knott Field, 1921.

Plots.										Yield per Acre
Electro-Culture	...	...	...	...	...	...	...	...	...	cwt.
Control	...	...	...	...	...	...	...	...	...	42.0
										41.2

Cereal Crops.

Plots.	Dressed Grain.		Offal Grain per Acre.	Straw per Acre.		Proportion of Total Grain to 100 of Total Straw.
	Yield per Acre.	Weight per Bush.		Straw.	Total Straw.	
	Bushels.	lb.	lb.	lb.	cwt.	

Oats (Grey Winter). Foster's Field, 1921.

Electro-Culture	...	...	40.7	43.4	241	1543	19.3	93.0
Control I.	...	...	33.1	42.0	298	1220	14.9	101.4
Control II.	...	...	31.6	42.2	234	1102	14.6	96.0

Wheat (Red Standard). Foster's Field, 1922.

Electro-Culture	...	...	15.4	61.4	234	1229	15.8	66.9
Control, North East	...	...	16.5	60.6	249	1272	15.5	72.1
Control, South East	...	...	17.2	61.8	231	1196	14.2	81.5

Barley (Plumage Archer). Great Knott Field, 1922.

Electro-Culture	...	...	34.1	49.1	273	1808	22.2	78.2
Control	...	...	32.4	48.6	244	1840	22.3	72.8

## BORON EXPERIMENT

Barley (Plumage Archer). Little Hoos, 1922.

Treatment of Plots.	Dressed Grain.						Offal Grain per Acre.			Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre. Bushels.			Weight per Bushel. lb.						Straw. lb.			Total Straw. cwt.					
	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3	Series 1	Series 2	Series 3
Boric Acid 20 lb. per acre ...	37.9	40.8	30.8	51.1	51.8	52.0	191	138	84	2025	1875	1850	24.6	23.2	22.1	77.4	86.5	68.2
Boric Acid 8 lb. per acre ...	36.5	40.0	41.3	51.5	52.0	52.0	169	113	150	1825	1800	1850	23.4	22.8	23.0	78.1	86.0	89.2
Control ...	34.9	40.8	38.6	50.9	52.4	52.5	156	134	119	1725	1775	1850	21.4	22.5	23.4	80.5	89.9	81.7

All plots received a basal dressing of Superphosphate 3 cwt.; Sulphate of Potash 1 cwt.; Sulphate of Ammonia 1½ cwt.

## EXPERIMENTS WITH NITROGENOUS MANURES

Potatoes (Arran Chief). Sawpit Field, 1921.

Manure per Acre.	Yield per Acre.		
	1st Plot.	2nd Plot.	3rd Plot.
	Tons.	Tons.	Tons.
4 cwt. Super., 1 cwt. Sulphate Potash, 2 cwt. Sulphate Ammonia	2.27	2.24	2.43
4 cwt. Super., 1 cwt. Sulphate Potash ... ..	1.84	2.13	1.99
4 cwt. Super., 1 cwt. Sulphate Potash, 193 lb. Muriate Ammonia	2.18	2.67	2.61
Control ... ..	1.33	1.41	1.49
4 cwt. Super., 1 cwt. Sulphate Potash, 102 lb. Urea ... ..	*1.72	2.69	2.57

\* The bouts on this plot were badly broken down due to extra hoeing on account of growth of Wheatbind.

Barley (Plumage Archer). Stackyard Field, 1921.

Manures per Acre.	Dressed Grain.						Offal Grain per Acre.			Straw per Acre.						Proportion of Total Grain to 100 of Total Straw.		
	Yield per Acre. Bushels.			Weight per Bushel. lb.						Straw. lb.			Total Straw. cwt.					
	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3	Plot 1	Plot 2	Plot 3
1½ cwt. Super., 145 lb. M./Amm.	40.4	34.8	—	54.7	55.5	—	197	135	—	2000	2000	—	23.5	25.9	—	91	71	—
1½ cwt. Super....	27.2	27.1	24.1	56.0	55.5	55.0	153	103	109	1325	1475	1350	17.1	18.4	17.2	88	78	75
1½ cwt. Super., 1½ cwt. S./Amm.	38.3	36.5	30.2	55.7	55.2	54.2	144	175	194	1900	2050	1825	23.6	24.9	22.1	87	79	74
1½ cwt. Super., 76½ lb. Urea ...	38.2	34.6	29.2	55.0	54.5	54.2	150	150	169	2000	2025	1775	24.2	24.7	21.3	83	74	73
No Manure ...	27.5	24.7	—	55.0	54.5	—	103	97	—	1400	1450	—	17.3	17.9	—	83	72	—

### MALTING BARLEY EXPERIMENT.

Plumage Archer. Long Hoos Field, 1922.

Manures per Acre.	Dressed Grain.		Offal Grain per Acre.	Straw per Acre		Proportion of Total Grain to 100 of Total Straw.
	Yield per Acre.	Weight per Bushel.		Straw.	Total Straw	
	Bushels	lb.	lb.	lb.	cwt.	
Super. 3 cwt., Sul./Pot. 1½ cwt., Sul./Amm. 1 cwt. ... ..	36·0	50·8	163	1213	17·1	104
Super. 3 cwt., Sul./Pot. 1½ cwt., Mur./Amm. 93 lb. ... ..	35·7	51·0	169	1388	18·5	96
Super. 3 cwt., Sul./Pot. 1½ cwt. ...	31·0	50·8	188	1263	17·0	93
Super. 3 cwt., Sul./Amm. 1 cwt. ...	30·0	50·3	175	975	14·1	107
Super. 3 cwt., Sul./Amm. 1 cwt., Mur./Pot. 1½ cwt.* ... ..	34·8	50·0	206	not	reco-	ded.
Sul./Amm. 1 cwt., Sul./Pot. 1½ cwt.	36·8	50·3	191	1438	19·9	92
No Manure ... ..	28·6	50·5	184	1125	15·5	94

\*Muriate of Potash applied on April 3rd. Other Manures on March 24th.

### MISCELLANEOUS EXPERIMENTS.

Clover. Hoos Field, 1921 and 1922.

(Formerly Barley after Alsike).

Plot.	Manures per Acre.	Yield per Acre.	
		1921.	1922.
		cwt.	cwt.
1	Slag 8 cwt., Lime 10 cwt. ... ..	45·3	17·4
2	Farmyard Manure 14 tons, Super. 5 cwt., Lime 10 cwt.	53·8	17·9
3	Lime 10 cwt. ... ..	35·9	17·6
4	Super. 5 cwt., Lime 10 cwt., Sulph. Potash 1½ cwt. ...	40·6	19·6
5	Super. 5 cwt., Lime 10 cwt. ... ..	45·3	13·0
6	Lime 10 cwt. ... ..	41·1	13·0
7	Farmyard Manure 14 tons, Lime 10 cwt. ... ..	54·5	16·7
8	Slag 8 cwt. ... ..	42·9	11·4
9	Farmyard Manure 14 tons, Super. 5 cwt. ... ..	50·5	17·2
10	Control ... ..	36·8	14·1
11	Super. 5 cwt., Sulph. Potash 1½ cwt. ... ..	45·1	20·3
12	Super. 5 cwt. ... ..	49·1	14·3
13	Control ... ..	36·6	9·4
14	Farmyard Manure 14 tons ... ..	46·2	10·7
15	Horse Dung 14 tons, Lime 10 cwt. ... ..	35·3	6·7
16	Control ... ..	35·5	7·1
17	Horse Dung 14 tons ... ..	54·9	11·6
18	Super. 5 cwt. ... ..	39·7	6·3
19	Cattle Dung 14 tons, Lime 10 cwt. ... ..	50·5	13·0
20	Control ... ..	33·3	3·6
21	Cattle Dung 14 tons ... ..	41·5	5·8

Manures applied and Clover sown in 1920.

Barley. Hoos Field. Leguminous Strips, 1921, 1922.

Description of Plot.	Manurial Treatment	1921.						1922.					
		Dressed Grain.		Offal Grain per Acre.	Straw per Acre.	Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.	Dressed Grain.		Offal Grain per Acre.	Straw per Acre.	Total Straw per Acre.	Proportion of Total Grain to 100 of Total Straw.
		Yield per Acre.	Weight per Bushel.					Yield per Acre.	Weight per Bushel.				
lb.	lb.	lb.	lb.	lb.	lb.	Bush.	lb.	Bush.	lb.	lb.	lb.	cwt.	
After Lucerne ...	Sulphate Amm. 1½ cwt. ...	14.8	56.5	134	688	10.2	85.5	27.2	51.4	188	1921	22.7	62.3
	S. Amm. 1½ cwt. Super. 3 cwt.	37.8	57.3	154	1310	18.0	115.4	41.2	52.0	161	1884	22.2	92.6
After Red Clover	Sulphate Amm. 1½ cwt. ...	12.1	56.4	122	555	8.4	85.2	25.2	50.6	134	1556	18.1	69.3
	S. Amm. 1½ cwt. Super. 3 cwt.	31.1	57.4	134	1037	15.7	109.0	35.4	51.5	109	1579	18.0	96.0
After Alsike ...	Sulphate Amm. 1½ cwt. ...	11.0	56.3	100	557	8.3	77.6	25.9	50.9	125	1481	17.2	74.8
	S. Amm. 1½ cwt. Super. 3 cwt.	28.7	57.8	137	871	15.2	105.4	33.6	52.0	92	1421	16.5	99.6

Leguminous crops ploughed in November, 1911.

