


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Report 1925-26 With the Supplement to the Guide to the Experimental Plots



[Full Table of Content](#)

The Classical Experiments

Rothamsted Research

Rothamsted Research (1926) *The Classical Experiments* ; Report 1925-26 With The Supplement To The Guide To The Experimental Plots, pp 123 - 137 - DOI:

<https://doi.org/10.23637/ERADOC-1-84>

DATES OF SOWING AND HARVESTING (Harvest 1925).

| Field. | Crop. | Variety. | Sowing began. | Sowing finished. | Cutting began. | *Carting began. | *Carting finished. | Yield per Acre. |
|-------------------|----------------|--|---------------|------------------|----------------|-----------------|--------------------|-----------------|
| Great Knott, east | Forage Mixture | Beans, Peas, Vetches, Oats, Wheat | Mar. 12, '25 | Mar. 16, '25 | June 29 | July 5 | July 6 | 21 cwt. |
| " west | Fallow | — | — | — | — | — | — | — |
| Little Knott | Grass | Mixture† | May 19, '25 | May 20, '25 | — | — | — | — § |
| Foster's, east | Barley | Plumage Archer | Mar. 19, '25 | Mar. 20, '25 | Aug. 18 | Aug. 28 | Aug. 31 | 48 bush. |
| " west | Swedes | Webb's Purple | June 3, '25 | June 8, '25 | — | Nov. 11 | Nov. 16 | 11 tons |
| West Barnfield | Potatoes | Kerr's Pink, King Edward, Great Scott | April 29, '25 | May 4, '25 | — | Oct. 6 | Oct. 29 | see p. 139 |
| | Mangolds | Red Intermediate | May 11, '25 | May 11, '25 | — | Oct. 16 | Oct. 24 | see p. 14† |
| Long Hoos, east | Oats | Grey Winter | Oct. 3, '24 | Oct. 4, '24 | July 20 | Aug. 8 | Aug. 17 | 68 bush. |
| " west | Wheat | Red Standard | Oct. 17, '24 | Oct. 17, '24 | Aug. 11 | Aug. 17 | Aug. 18 | 40 bush. |
| New Zealand | Mangolds | Sutton's Prizewinner, Red Intermediate | May 14, '25 | May 15, '25 | — | Sept. 17 | Oct. 15 | 25 tons |
| Stackyard | Turnips | Mammoth Green Top | June 2, '25 | June 2, '25 | — | July 16 | July 20 | 17 tons |
| Great Harpenden | Oats | Giant Eliza | Mar. 6, '25 | Mar. 6, '25 | Aug. 1 | Aug. 15 | Aug. 15 | 40 bush. |
| | Clover | Broad Red | Mar. 18, '24 | Mar. 21, '24 | Aug. 15 | Aug. 28 | Oct. 15 | Failed |
| | Beans | Spring | Feb. 19, '25 | Feb. 21, '25 | June 23 | June 24 | June 26 | 2 tons† |
| Sawpit | Clover | Broad Red | April 4, '24 | April 5, '24 | June 15 | June 19 | June 20 | 30 cwt. |
| | Wheat | Red Standard | Nov. 10, '24 | Nov. 11, '24 | Sept. 26 | Oct. 10 | Oct. 15 | — |
| Sawyers | Oats | Grey Winter | Nov. 24, '24 | Nov. 26, '24 | Aug. 14 | Aug. 31 | Sept. 2 | 32 bush. |
| Broadbalk | Wheat | Svalof Victory | Mar. 30, '25 | Mar. 30, '25 | July 20 | Aug. 6 | Aug. 8 | 48 bush. |
| Little Hoos | Fallow | Red Standard | Oct. 24, '24 | Oct. 24, '24 | Aug. 26 | Sept. 7 | Sept. 8 | 40 bush. |
| Great Hoos | Barley | Plumage Archer | Mar. 19, '25 | Mar. 19, '25 | Aug. 17 | Aug. 29 | Aug. 29 | see p. 132 |
| Barnfield | Mangolds | Sutton's Prizewinner | May 15, '25 | May 15, '25 | — | — | — | see p. 135 |
| Agdell | Barley | Plumage Archer | April 3, '25 | April 3, '25 | Sept. 3 | Sept. 22 | Sept. 22 | see p. 127 |
| Great Field | Hay | — | — | — | June 18 | June 22 | June 23 | see p. 125 |
| Park | Grass | — | — | — | June 12 | June 17 | June 18 | 25 cwt. |
| | Grass | — | — | — | — | — | — | see p. 128 |

* In the case of roots, the dates given are those on which lifting began and finished.
 † Crop cut green for silage.
 ‡ The mixture consisted of Broad Red Clover; Wild White Clover; Indigenous Cocksfoot; Meadow Fescue; Timothy; Perennial Rye; Wild Perennial Rye; Rough-stalked Meadow Grass.
 § No yield. First year of permanent grass.

DATES OF SOWING AND HARVESTING (Harvest 1926).

| Field. | Crop. | Variety. | Sowing began. | Sowing finished. | Cutting began. | *Carting began. | *Carting finished. | †Yield per acre. |
|-----------------------|------------------|---|---------------|------------------|----------------|--------------------|--------------------|------------------|
| Great Knott, west ... | Wheat | Red Standard Cambridge, Bro-wick, Little Joss, Midlothian III | Oct. 26, '25 | Oct. 29, '25 | Aug. 20 | Aug. 26 | — | — |
| " east | Fallow | — | — | — | — | — | — | — |
| Little Knott... | Oats | Svalof Victory | — | — | June 21 | June 29 | July 1 | 40 cwt. |
| Foster's, east | Grass | Permanent Grass | — | — | June 22 | July 1 | July 2 | 37 cwt. |
| " west | Clover | Broad Red, late flowering | — | — | — | — | — | — |
| West Barnfield | Clover | — | — | — | — | — | — | — |
| Long Hoos, east | Barley | Plumage Archer | Mar. 17, '26 | Mar. 18, '26 | Aug. 30 | Sept. 11 | — | — |
| " west | Winter Oats | Grey Winter | Oct. 9, '25 | Oct. 10, '25 | July 28 | Aug. 16 | Aug. 18 | 8 qrs. |
| Stackyard ... | Forage Crop | Beans, Peas, Vetches and Cereals | Oct. 14, '25 | Oct. 15, '25 | Sept. 9 | Sept. 13 | — | 4 qrs. |
| New Zealand | Potatoes | Kerr's Pink | April 23, '26 | — | Sept. 21 | — | — | Av. 10 tn. |
| Great Harpenden ... | Mangolds | Cannells QQ | April 14, '26 | April 24, '26 | Oct. 6 | Oct. 13 | Oct. 19 | Av. 22 tn. |
| Sawpit ... | Barley | Plumage Archer | Mar. 16, '26 | Mar. 16, '26 | Aug. 23 | Aug. 31 | Sept. 10 | — |
| Sawyers ... | Rye | Swedish | Oct. 29, '25 | — | Aug. 11 | Aug. 19 | Aug. 20 | 5 qrs. |
| Broadbalk ... | Wheat | Red Standard | Oct. 30, '25 | Nov. 11, '25 | Aug. 16 | Aug. 26 | — | 2½ qrs. |
| Little Hoos | Grass | Permanent seeding | April 17, '25 | April 19, '25 | June 30 | July 8 | July 9 | 15 cwt. |
| Hoos | Swedes | Dreadnought | June 14, '26 | — | Dec. 1 | Dec. 1 | Dec. 20 | 19 tons |
| Agdell | Fallow | (White Mustard ploughed down) | July 2, '26 | — | Ploughed down | Sept. 2 to Sept. 2 | Sept. 2 | 20 |
| Great Field | Wheat | Red Standard | Nov. 25, '25 | — | — | Sept. 1 | Sept. 2 | — |
| " | Fallow | — | — | — | — | — | — | — |
| " | Swedes | Purple King | June 3, '26 | — | Oct. 20 | — | — | — |
| " | Barley | Plumage Archer | April 7, '26 | April 8, '26 | Aug. 25 | Sept. 10 | Sept. 11 | — |
| " | Oats | Svalof Victory | — | — | Aug. 16 | Aug. 30 | — | 6 qrs. |
| " | Wheat and Fallow | Red Standard | — | — | — | — | — | — |
| " | Clover (failed) | Broad Red | — | — | June 28 | July 6 | — | 10 cwt. |
| " | Mixed Legumes | Vetches, Oats, Italian Clover, etc. | April 16, '26 | — | Ploughed down | down after math. | — | — |
| " | Grazing Plots | — | — | — | — | — | — | — |
| " | Hay | — | — | — | June 21 | June 28 | July 7 | 33 cwt. |
| " | Hay | — | — | — | { June 15 | June 31 | July 5 | — |
| " | Hay | — | — | — | { " 22 | — | — | — |
| " | Hay | — | — | — | { " 24 | — | — | — |

* In the case of roots, the dates given are those on which lifting began and finished. † Estimates of standing crops.

CROP YIELDS ON THE EXPERIMENTAL PLOTS.

NOTES.—In each case the year refers to the harvest, *e.g.*, Wheat harvested in 1926.
In the tables, total straw includes straw, cavings and chaff.

CONVERSION TABLE.

| | | |
|--------------------------|--|---------------------------------|
| 1 acre = | 0.405 Hectare = | 0.963 Feddan. |
| 1 bushel (Imperial) = | 0.364 Hectolitre (36.364 litres) ... = | 0.184 Ardeb. |
| 1 lb.(poundavoirdupois)= | 0.453 Kilogramme = | 1.009 Rotls. |
| 1 cwt. (hundredweight)= | 50.8 Kilogrammes = | { 113.0 Rotls. 1.366 Maunds. |
| 1 metric quintal ... = | { 100.0 Kilogrammes = 220.46 lb. = | |
| 1 bushel per acre = | 0.9 Hectolitre per Hectare ... = | 0.191 Ardeb per Feddan. |
| 1 lb. per acre ... = | 1.12 Kilogramme per Hectare ... = | 1.049 Rotls per Feddan. |
| 1 cwt. per acre ... = | 125.60 Kilogrammes per Hectare or 1.256 metric Quintals per Hectare | 117.4 Rotls per Feddan. |

In America the Winchester bushel is used=35.236 litres. 1 English bushel=1.032 American bushels.

CROPS GROWN IN ROTATION. AGDELL FIELD.

PRODUCE PER ACRE.

| Year. | CROP. | O. Unmanured. | | M. Mineral Manure. | | C. Complete Mineral & Nitrogenous M'nure | |
|--|---|------------------|------------------------------|-----------------------|------------------------------|--|------------------------------|
| | | 5. Fallow. | 6. Clover or Beans. | 3. Fallow. | 4. Clover or Beans. | 1. Fallow. | 2. Clover or Beans. |
| AVERAGE OF THE FIRST NINETEEN COURSES, 1848-1923. | | | | | | | |
| | Roots (Swedes) cwt.* | 32.7 | 11.2 | 175.7 | 195.9 | 355.3 | 302.0 |
| | Barley— | | | | | | |
| | Dressed Grain bush. | 22.7 | 20.9 | 23.8 | 27.9 | 32.2 | 36.8 |
| | Total Straw ... cwt. | 13.9 | 13.7 | 14.0 | 16.0 | 19.5 | 22.6 |
| | Beans— | | | | | | |
| | Dressed Grain bush. | — | 13.1 | — | 18.2 | — | 22.3 |
| | Total Straw ... cwt. | — | 9.2 | — | 13.2 | — | 15.3 |
| | Clover Hay ... cwt. | — | 28.3 | — | 54.1 | — | 55.0 |
| | Wheat— | | | | | | |
| | Dressed Grain bush. | 24.2 | 22.8 | 28.5 | 31.2 | 29.5 | 31.2 |
| | Total Straw ... cwt. | 23.7 | 21.7 | 29.0 | 30.3 | 31.4 | 30.4 |
| PRESENT COURSE (20th), 1924, 1925 and 1926. | | | | | | | |
| 1924 | Roots (Turnips) ... cwt. | 2.9 | 0.7 | 42.8 | 31.5 | 127.4 | 104.7 |
| 1925 | Barley— | | | | | | |
| | Dressed Grain bush. | 10.86 | 7.35 | 10.09 | 16.70 | 10.35 | 8.60 |
| | Offal Grain ... lb. | 42.0 | 49.0 | 94.0 | 38.0 | 53.0 | 59.0 |
| | Straw lb. | 633.0 | 678.0 | 602.0 | 866.0 | 626.0 | 541.0 |
| | Total Straw ... cwt. | 7.2 | 7.5 | 7.4 | 9.3 | 7.0 | 6.5 |
| | Wt. of Dressed Grain per bush. } lb. | 52.7 | 51.6 | 52.5 | 53.6 | 53.3 | 54.3 |
| | Proportion of Total Grain to 100 of Total Straw } | 76.3 | 50.7 | 75.5 | 89.2 | 77.0 | 72.4 |
| 1926 | Clover Hay ... cwt. | — | 14.2 | — | 32.2 | — | 26.3 |

* Plots 1, 3 and 5 based upon 18 years. Plots 2, 4 and 6 based upon 17 years.

METEOROLOGICAL RECORDS, 1925 and 1926.

| | Rain. | | Drainage through soil. | | | Bright Sunshine. | Temperature (Mean). | | | | |
|---------------|--|---|------------------------|---------------|---------------|------------------|---------------------|------|------------------|------------|------------|
| | Total Fall. $\frac{1}{1000}$ Acre Gauge. | No. of Rainy Days. (0.01 inch or more) $\frac{1}{1000}$ Acre Gauge. | 20 ins. deep. | 40 ins. deep. | 60 ins. deep. | | Max. | Min. | 1 ft. in ground. | Solar Max. | Grass Min. |
| 1925 | Inches. | No. | Inches. | Inches. | Inches. | Hours. | °F. | °F. | °F. | °F. | °F. |
| Jan. ... | 2.053 | 18 | 1.804 | 1.870 | 1.845 | 52.7 | 44.6 | 34.6 | 39.6 | 64.2 | 32.2 |
| Feb. ... | 3.940 | 16 | 3.413 | 3.452 | 3.457 | 68.3 | 45.3 | 35.7 | 40.0 | 83.4 | 31.7 |
| Mar. ... | 1.219 | 12 | 0.340 | 0.442 | 0.426 | 89.3 | 45.0 | 34.5 | 39.2 | 91.8 | 30.2 |
| April ... | 1.703 | 16 | 0.149 | 0.183 | 0.169 | 139.6 | 52.1 | 37.1 | 44.3 | 106.8 | 32.7 |
| May ... | 2.480 | 18 | 0.391 | 0.534 | 0.486 | 204.7 | 60.8 | 44.7 | 52.4 | 121.2 | 40.7 |
| June ... | 0.121 | 2 | 0.002 | 0.033 | 0.043 | 259.5 | 68.0 | 48.2 | 59.6 | 119.4 | 43.1 |
| July ... | 4.428 | 15 | 1.573 | 1.343 | 1.284 | 183.6 | 70.9 | 53.4 | 62.4 | 125.5 | 48.4 |
| Aug. ... | 2.972 | 15 | 1.048 | 1.180 | 1.095 | 133.1 | 65.8 | 52.8 | 60.1 | 116.9 | 49.1 |
| Sept. ... | 3.287 | 18 | 1.528 | 1.605 | 1.501 | 124.3 | 58.6 | 46.0 | 53.7 | 112.0 | 40.9 |
| Oct. ... | 3.013 | 14 | 2.078 | 2.203 | 2.037 | 102.9 | 56.5 | 44.2 | 51.0 | 97.7 | 39.9 |
| Nov. ... | 2.241 | 15 | 1.481 | 1.706 | 1.616 | 90.6 | 43.4 | 34.1 | 42.2 | 76.6 | 29.8 |
| Dec. ... | 2.127 | 16 | 1.900 | 2.052 | 1.903 | 57.8 | 41.3 | 31.3 | 36.3 | 60.6 | 27.6 |
| Total or Mean | 29.584 | 175 | 15.707 | 16.603 | 15.862 | 1506.4 | 54.4 | 41.4 | 48.4 | 98.0 | 37.2 |
| 1926 | | | | | | | | | | | |
| Jan. ... | 3.511 | 19 | 3.169 | 3.387 | 3.260 | 45.7 | 43.9 | 32.5 | 38.4 | 66.2 | 29.6 |
| Feb. ... | 2.494 | 17 | 2.112 | 2.431 | 2.298 | 40.6 | 48.4 | 39.5 | 42.1 | 72.5 | 35.4 |
| Mar. ... | 0.215 | 5 | 0.003 | 0.049 | 0.041 | 119.9 | 49.4 | 36.9 | 42.3 | 99.3 | 30.5 |
| April ... | 2.963 | 16 | 0.861 | 0.938 | 0.862 | 108.2 | 55.3 | 40.7 | 46.4 | 105.9 | 35.3 |
| May ... | 1.945 | 18 | 0.369 | 0.653 | 0.581 | 153.6 | 57.4 | 42.9 | 50.5 | 117.1 | 38.3 |
| June ... | 3.014 | 13 | 0.943 | 1.258 | 1.157 | 180.7 | 63.3 | 47.9 | 57.8 | 123.9 | 42.9 |
| July ... | 2.787 | 11 | 0.291 | 0.442 | 0.384 | 151.5 | 68.6 | 54.5 | 61.5 | 123.9 | 50.5 |
| Aug. ... | 1.190 | 9 | — | 0.035 | 0.033 | 195.2 | 69.0 | 52.8 | 60.9 | 122.8 | 47.4 |
| Sept. ... | 1.788 | 11 | 0.576 | 0.659 | 0.600 | 133.2 | 65.8 | 51.3 | 59.3 | 112.8 | 46.3 |
| Oct. ... | 2.672 | 14 | 1.149 | 1.230 | 1.135 | 98.5 | 52.4 | 40.3 | 48.9 | 95.9 | 35.7 |
| Nov. ... | 5.321 | 24 | 4.520 | 4.840 | 4.644 | 45.0 | 47.7 | 37.4 | 43.3 | 75.8 | 33.0 |
| Dec. ... | 0.477 | 6 | 0.329 | 0.525 | 0.467 | 64.5 | 42.3 | 33.8 | 38.8 | 67.8 | 29.9 |
| Total or Mean | 28.377 | 163 | 14.322 | 16.447 | 15.462 | 1336.6 | 55.3 | 42.5 | 49.2 | 98.7 | 37.9 |

RAIN AND DRAINAGE.
MONTHLY MEAN FOR 56 HARVEST YEARS, 1870-1—1925-6.

| | Rainfall. | Drainage. | | | Drainage % of Rainfall. | | | Evaporation. | | |
|-------------|-----------|--------------|--------------|--------------|-------------------------|--------------|--------------|--------------|--------------|--------------|
| | | 20-in. Gauge | 40-in. Gauge | 60-in. Gauge | 20-in. Gauge | 40-in. Gauge | 60-in. Gauge | 20-in. Gauge | 40-in. Gauge | 60-in. Gauge |
| September | Ins. | Ins. | Ins. | Ins. | % | % | % | Ins. | Ins. | Ins. |
| October ... | 3.161 | 1.830 | 1.789 | 1.662 | 57.9 | 56.6 | 52.6 | 1.331 | 1.372 | 1.499 |
| November | 2.725 | 2.055 | 2.091 | 1.971 | 75.4 | 76.7 | 72.3 | 0.670 | 0.634 | 0.754 |
| December | 2.857 | 2.439 | 2.525 | 2.411 | 85.4 | 88.4 | 84.4 | 0.418 | 0.332 | 0.446 |
| January ... | 2.389 | 1.942 | 2.123 | 2.043 | 81.3 | 88.9 | 85.5 | 0.447 | 0.266 | 0.346 |
| February | 2.039 | 1.515 | 1.618 | 1.545 | 74.3 | 79.4 | 75.8 | 0.524 | 0.421 | 0.494 |
| March ... | 2.027 | 1.091 | 1.221 | 1.154 | 53.8 | 60.2 | 56.9 | 0.936 | 0.806 | 0.873 |
| April ... | 2.053 | 0.660 | 0.730 | 0.696 | 32.1 | 35.6 | 33.9 | 1.393 | 1.323 | 1.357 |
| May ... | 2.054 | 0.484 | 0.550 | 0.516 | 23.6 | 26.8 | 25.1 | 1.570 | 1.504 | 1.538 |
| June ... | 2.245 | 0.560 | 0.588 | 0.567 | 24.9 | 26.2 | 25.3 | 1.685 | 1.657 | 1.678 |
| July ... | 2.746 | 0.726 | 0.748 | 0.696 | 26.4 | 27.2 | 25.3 | 2.020 | 1.998 | 2.050 |
| August ... | 2.662 | 0.699 | 0.704 | 0.660 | 26.3 | 26.4 | 24.8 | 1.963 | 1.958 | 2.002 |
| Year ... | 29.342 | 14.786 | 15.440 | 14.610 | 50.4 | 52.6 | 49.8 | 14.556 | 13.902 | 14.732 |

Area of each gauge $\frac{1}{10000}$ acre.

MANGOLDS, BARN FIELD, 1925 and 1926.

Roots since 1856. Mangolds since 1876.

Produce per Acre.

| Strip. | Strip Manures. | Cross Dressings. | | | | |
|--------|--|---|---|------------------------|-----------------------------|------------------------|
| | | O. | N. | A. | A.C. | C. |
| | | None. | Nitrate of Soda. | Ammon. Salts. | Ammon. Salts and Rape Cake. | Rape Cake. |
| | | Tons | Tons | Tons | Tons | Tons |
| 1 | 1925. Dung only | { R. 14.28 L. 2.77 | { 25.55 5.98 | { 19.14 6.35 | { 18.99 6.74 | { 18.20 5.77 |
| 2 | Dung, Super., Potash ... | { R. 16.19 L. 2.98 | { 27.13 6.41 | { 25.21 6.26 | { 23.22 7.28 | { 23.25 6.49 |
| 4 | Complete Minerals ... | { R. 3.25 ^a L. 0.93 ^b | { R.16.84* L. 4.98 R.16.90 L. 5.65 | { 14.27 3.68 | { 22.43 6.05 | { 16.07 3.98 |
| 5 | Superphosphate only ... | { R. 3.64 L. 1.12 | { 14.01 4.32 | { 6.10 3.69 | { 6.30 4.51 | { 6.63 4.26 |
| 6 | Super. and Potash ... | { R. 4.16 L. 1.11 | { 14.31 4.36 | { 13.91 3.59 | { 18.18 5.90 | { 13.46 3.66 |
| 7 | Super., Sulphate of Mag., and Sodium Chloride | { R. 3.49 L. 1.00 | { 14.81 3.23 | { 14.21 3.05 | { 13.37 5.25 | { 12.09 3.38 |
| 8 | None | { R. 2.32 L. 1.01 | { 4.94 3.37 | { 2.81 2.23 | { 5.25 3.39 | { 4.03 2.32 |
| 9 | Sodium Chloride, Nit. Soda, Sulph. Potash, and Sulph. Mag. ... | { R. 17.08 L. 3.83 | { — — | { — — | { — — | { — — |
| | | Tons | Tons | Tons | Tons | Tons |
| 1 | 1926. Dung only | { R. 21.16 L. 3.39 | { 31.39 4.58 | { 21.77 4.24 | { 18.35 3.81 | { 19.39 4.88 |
| 2 | Dung, Super., Potash ... | { R. 23.80 L. 3.25 | { 34.72 4.83 | { 30.84 5.22 | { 30.08 6.07 | { 27.90 5.47 |
| 4 | Complete Minerals ... | { R. 4.75 ^a L. 0.85 ^b | { R.24.07* L. 3.93 R.23.75 L. 4.51 | { 19.52 2.92 | { 25.77 4.12 | { 16.39 2.52 |
| 5 | Superphosphate only ... | { R. 4.81 L. 0.86 | { 18.39 2.67 | { 9.25 2.17 | { 8.29 2.25 | { 10.28 2.39 |
| 6 | Super. and Potash ... | { R. 5.41 L. 0.89 | { 20.80 3.02 | { 17.86 2.58 | { 21.05 4.12 | { 13.29 1.94 |
| 7 | Super., Sulphate of Mag., and Sodium Chloride | { R. 5.28 L. 0.96 | { 21.27 3.24 | { 18.86 3.08 | { 20.00 3.94 | { 11.66 2.36 |
| 8 | None | { R. 3.36 L. 0.81 | { 13.97 3.72 | { 7.83 3.02 | { 7.73 2.41 | { 8.04 2.57 |
| 9 | Sodium Chloride, Nit. Soda, Sulph. Potash and Sulph. Mag. ... | { R. 25.09 L. 3.11 | { — — | { — — | { — — | { — — |

R.=roots. L.=leaves.

* From 1904 onwards plot 4 N has been divided, 4a receiving Sulphate of Potash, Sulphate of Magnesia, Sodium Chloride and Nitrate of Soda; 4b receiving Calcium Chloride, Potassium Nitrate and Calcium Nitrate.

HAY. THE PARK GRASS PLOTS. 1925, 1926.

| Plot | Manuring per acre | 1925 | | | | | | 1926 | | | | | | | |
|------|---|-----------------------|----------|-------|---------------------|----------|-------|-----------------------|----------|-------|---------------------|----------|-------|------|------|
| | | Yield of Hay per acre | | | Dry Matter per acre | | | Yield of Hay per acre | | | Dry Matter per acre | | | | |
| | | 1st Crop | 2nd Crop | Total | 1st Crop | 2nd Crop | Total | 1st Crop | 2nd Crop | Total | 1st Crop | 2nd Crop | Total | | |
| 1 | Single dressing Amm. Salts (=43 lb. N.) ; (with Dung also 8 years, 1856-63) ... | 15.1 | 16.1 | 31.2 | 1418 | 1268 | 2686 | lb. | 16.5 | 9.8 | 26.3 | 1602 | 881 | 2483 | 1 |
| 2 | Unmanured (after Dung 8 years, 1856-63) | 23.1 | 14.9 | 38.0 | 2075 | 1155 | 3230 | lb. | 21.9 | 9.2 | 31.1 | 2111 | 823 | 2934 | 2 |
| 3 | Unmanured ... | 21.7 | 12.8 | 34.5 | 1987 | 884 | 2871 | lb. | 16.3 | 8.5 | 24.8 | 1406 | 764 | 2170 | 3 |
| 4-1 | Superphosphate of Lime ... | 12.5 | 11.7 | 24.2 | 1122 | 871 | 1993 | lb. | 11.8 | 9.0 | 20.8 | 1026 | 807 | 1833 | 4-1 |
| 4-2 | Superphosphate of Lime and double dressing Amm. Salts (=86 lb. N.) ... | 22.5 | 14.6 | 37.1 | 1971 | 971 | 2942 | lb. | 16.8 | 10.6 | 27.4 | 1511 | 952 | 2463 | 4-2 |
| 5-1 | (N. half) Unmanured following double dressing Amm. Salts (=86 lb. N.) 1856-97 ... | 20.2 | 12.3 | 32.5 | 1839 | 839 | 2678 | lb. | 14.5 | 7.6 | 22.1 | 1229 | 682 | 1911 | 5-1 |
| 5-2 | (S. half) Superphosphate, Sulphate of Potash; following double dressing Amm. Salts (=86 lb. N.) 1856-68 ... | 22.6 | 7.3 | 29.9 | 1812 | 570 | 2382 | lb. | 24.6 | 6.6 | 31.2 | 1843 | 595 | 2438 | 5-2 |
| 6 | Complete Mineral Manure as plot 7; following double dressing Amm. Salts (=86 lb. N.) 1856-68 ... | 32.7 | 16.7 | 49.4 | 3126 | 1326 | 4452 | lb. | 37.1 | 10.1 | 47.2 | 3259 | 901 | 4160 | 6 |
| 7 | Complete Mineral Manure ... | 13.9 | 8.0 | 21.9 | 1288 | 625 | 1913 | lb. | 12.6 | 8.8 | 21.4 | 1169 | 792 | 1961 | 7 |
| 8 | Mineral Manure without Potash ... | 24.2 | 14.9 | 39.1 | 2187 | 1152 | 3339 | lb. | 24.2 | 10.6 | 34.8 | 2137 | 951 | 3088 | 8 |
| 9 | Complete Mineral Manure and double dressing Amm. Salts (=86 lb. N.) ... | 26.6 | 23.1 | 49.7 | 2320 | 1329 | 3649 | lb. | 31.5 | 15.7 | 47.2 | 2835 | 1402 | 4237 | 9 |
| 10 | Mineral Manure (without Potash) and double dressing Amm. Salts (=86 lb. N.) ... | 28.2 | 22.8 | 51.0 | 2480 | 1403 | 3883 | lb. | 32.2 | 18.4 | 50.6 | 2949 | 1651 | 4600 | 10 |
| 11-1 | Complete Mineral Manure and treble dressing Amm. Salts (129 lb. N.) ... | 35.0 | 19.1 | 54.1 | 2900 | 1051 | 3951 | lb. | 32.8 | 14.0 | 46.8 | 3450 | 1251 | 4701 | 11-1 |

| | | | | | | | | | | | | | | |
|------|--|------|------|------|-------|------|------|------|------|------|------|------|------|------|
| 11-2 | As plot 11-1 and Silicate of Soda ... | 49.5 | 30.3 | 79.8 | 44.24 | 1950 | 6374 | 59.4 | 24.4 | 83.8 | 4535 | 2188 | 6723 | 11-2 |
| 12 | Unmanured ... | 63.9 | 26.4 | 90.3 | 5847 | 2150 | 7997 | 58.3 | 28.4 | 86.7 | 4860 | 2548 | 7408 | 12 |
| 13 | Dung 1905, and every fourth year since (omitted 1917), Fish Guano in 1907 and every fourth year since ... | 17.6 | 13.1 | 30.7 | 1630 | 1000 | 2630 | 18.4 | 15.4 | 33.8 | 1646 | 1376 | 3022 | 13 |
| 14 | Complete Mineral Manure and double dressing Nitrate of Soda (=86 lb. N.) ... | 45.8 | 26.9 | 72.7 | 4099 | 1461 | 5560 | 45.6 | 24.0 | 69.6 | 3767 | 2147 | 5914 | 14 |
| 15 | Complete Mineral Manure as plot 7; following double dressing Nitrate of Soda (=86 lb. N. 1858-1875) ... | 38.5 | 25.3 | 63.8 | 3536 | 1406 | 4942 | 41.1 | 21.9 | 63.0 | 3465 | 1964 | 5429 | 15 |
| 16 | Complete Mineral Manure and single dressing Nitrate of Soda (=43 lb. N.) ... | 61.1 | 25.2 | 86.3 | 4709 | 1886 | 6595 | 56.2 | 23.0 | 79.2 | 4626 | 2064 | 6690 | 16 |
| 17 | Complete Mineral Manure and single dressing Nitrate of Soda (=43 lb. N.) ... | 58.4 | 20.0 | 78.4 | 4725 | 1413 | 6138 | 55.9 | 16.7 | 72.6 | 4469 | 1499 | 5968 | 17 |
| 18 | Single dressing Nitrate of Soda (=43 lb. N.) ... | 47.4 | 11.3 | 58.7 | 3846 | 781 | 4627 | 49.6 | 5.2 | 54.8 | 4181 | 465 | 4646 | 18 |
| 19 | Mineral Manure (without Super.), and double dressing Sulphate of Amm. (=86 lb. N.) 1905 and since; following Minerals and Amm. Salts supplying the constituents of 1 ton of Hay, 1865-1904 ... | 33.5 | 25.8 | 59.3 | 2854 | 1650 | 4504 | 30.5 | 19.5 | 50.0 | 2674 | 1750 | 4424 | 19 |
| 20 | Farmyard Dung in 1905 and every fourth year since (omitted in 1917) following Nitrate of Soda (=43 lb. N.) and Minerals, 1872-1904 ... | 29.2 | 21.6 | 50.8 | 2676 | 1555 | 4231 | 25.8 | 13.4 | 39.2 | 2442 | 1196 | 3638 | 20 |
| | | 41.6 | 21.6 | 63.2 | 3480 | 1409 | 4889 | 38.6 | 18.8 | 57.4 | 2993 | 1688 | 4681 | |
| | | 44.1 | 18.5 | 62.6 | 3763 | 1273 | 5036 | 38.0 | 17.1 | 55.1 | 3046 | 1529 | 4575 | |
| | | 26.8 | 15.9 | 42.7 | 2137 | 1024 | 3161 | 23.1 | 12.5 | 35.6 | 2010 | 1123 | 3133 | |
| | | 30.8 | 14.9 | 45.7 | 2633 | 999 | 3632 | 28.5 | 12.6 | 41.1 | 2571 | 1127 | 3698 | |
| | | 17.7 | 14.6 | 32.3 | 1690 | 895 | 2585 | 19.4 | 24.0 | 43.4 | 1689 | 2147 | 3836 | |
| | | 36.5 | 21.7 | 58.2 | 3372 | 1253 | 4625 | 49.7 | 24.2 | 73.9 | 4086 | 2165 | 6251 | |
| | | 29.8 | 18.7 | 48.5 | 2757 | 974 | 3731 | 39.1 | 19.0 | 58.1 | 3039 | 1703 | 4742 | |
| | | 29.4 | 18.4 | 47.8 | 2398 | 1123 | 3521 | 34.7 | 17.5 | 52.2 | 3106 | 1569 | 4675 | |
| | | 27.0 | 15.0 | 42.0 | 2421 | 849 | 3270 | 30.0 | 8.7 | 38.7 | 2771 | 777 | 3548 | |
| | | 27.0 | 14.1 | 41.1 | 2282 | 902 | 3184 | 32.6 | 10.8 | 43.4 | 3108 | 970 | 4078 | |
| | | 35.8 | * | * | 3010 | * | * | 47.1 | 18.1 | 65.2 | 4693 | 1625 | 6318 | |
| | | 30.4 | 15.0 | 45.4 | 2654 | 1033 | 3687 | 43.9 | 11.5 | 55.4 | 4246 | 1027 | 5273 | |
| | | 41.6 | 14.4 | 56.0 | 3513 | 977 | 4490 | 43.3 | 16.6 | 59.9 | 4243 | 1488 | 5731 | |

Ground lime was applied to the Southern portion (limed) of the plots at the rate of 2,000 lb. to the acre in the Winter of 1903-4, 1907-8, 1915-16, 1923-24, and at the rate of 2,500 lb. to the acre in the Winter of 1920-21, except where otherwise stated.

Up to 1914 the limed and unlimed plot results were not separately given in the Annual Report, but the mean of the two was given. From 1915 onwards the separate figures are given.

* Figures for this plot not recorded.

§ The second crop was carted green; the figures given are estimated hay yields, calculated from the dry matter.

The Park Grass Plots.
 BOTANICAL COMPOSITION, PER CENT. 1923, 1st CROP.

| Plot | Manuring | Liming | Gramineæ | Leguminosæ | Others | "Other Orders" consist largely of | Plot |
|------|--|--------------|----------|------------|--------|--|------|
| 3 | Unmanured | Limed ... | 63.7 | 4.6 | 31.6 | Plantago lanceolata; Poterium sanguisorba; Luzula campestris | 3 |
| 7 | Complete Mineral Manure | Unlimed ... | 63.6 | 10.6 | 25.8 | Plantago lanceolata; Centaurea nigra; Poterium sanguisorba | 7 |
| 9 | Complete Mineral Manure and double Amm. Salts | Limed ... | 52.7 | 40.1 | 7.1 | Achillea millefolium; Ranunculus sp. | 9 |
| 14 | Complete Mineral Manure and double Nitrate of Soda | Unlimed ... | 69.1 | 15.2 | 15.7 | Plantago lanceolata; Spiræa ulmaria, etc. | 14 |
| 15 | As plot 7 following double Nitrate of Soda, 1858-75 | Limed ... | 99.4 | — | 0.6 | Rumex acetosa | 15 |
| 17 | Single Nitrate of Soda | Unlimed ... | 99.7 | — | 0.2 | Rumex acetosa | 17 |
| 18 | Mineral Manure (without Super.) and double Sulphate Amm. 1905 and since | Limed ... | 96.0 | 0.8 | 3.2 | Taraxacum vulgare | 18 |
| 19 | Farmyard Dung in 1905 and every 4th year since (omitted in 1917) | Unlimed ... | 93.7 | 0.1 | 6.2 | Taraxacum vulgare; Anthriscus sylvestris; Rumex acetosa | 19 |
| 20 | Farmyard Dung in 1905 and every 4th year since (omitted in 1917), each intervening year Sulphate Potash, Super., and Nitrate of Soda | Limed ... | 69.2 | 18.3 | 12.4 | Plantago lanceolata; Conopodium denudatum; Taraxacum vulgare | 20 |
| | | Unlimed ... | 57.8 | 15.4 | 26.8 | Plantago lanceolata; Luzula campestris; Conopodium denudatum | |
| | | Limed ... | 73.9 | 1.2 | 24.9 | Plantago lanceolata; Leontodon hispidus; Centaurea nigra | |
| | | Unlimed ... | 65.6 | 0.1 | 34.3 | Plantago lanceolata; Leontodon hispidus; Centaurea nigra | |
| | | L. 6,788 lb. | 87.4 | — | 12.6 | Rumex acetosa | |
| | | L. 3,951 lb. | 85.6 | — | 14.4 | Rumex acetosa | |
| | | Unlimed ... | 96.8 | 0.1 | 3.0 | Rumex acetosa | |
| | | L. 3,150 lb. | 72.4 | 17.0 | 10.6 | Ranunculus sp.; Plantago lanceolata; Conopodium denudatum | |
| | | L. 570 lb. | 79.2 | 10.0 | 10.7 | Ranunculus sp.; Rumex acetosa; Conopodium denudatum | |
| | | Unlimed ... | 78.5 | 7.4 | 14.1 | Ranunculus sp.; Rumex acetosa; Anthriscus sylvestris | |
| | | L. 2,772 lb. | 82.7 | 5.3 | 11.9 | Anthriscus sylvestris; Ranunculus sp.; Conopodium denudatum; Tragopogon pratensis | |
| | | L. 570 lb. | 82.5 | 10.6 | 6.8 | Ranunculus sp.; Conopodium denudatum | |
| | | Unlimed ... | 88.2 | 2.5 | 9.3 | Anthriscus sylvestris; Rumex acetosa; Ranunculus sp. | |

The Park Grass Plots—contd.
BOTANICAL COMPOSITION, PER CENT. 1924, 1st CROP.

| Plot | Manuring | Liming | Gramineae | Leguminosae | Other Orders | "Other Orders" consist largely of | Plot |
|------|---|--------------|-----------|-------------|--------------|---|------|
| 3 | Unmanured | Limed ... | 51.2 | 14.5 | 34.3 | Centaurea nigra; Scabiosa arvensis; Plantago lanceolata; Poterium sanguisorba | 3 |
| 5-1 | (N. half), Unmanured following double dressing of Amm. Salts (=86 lb. N.), 1856-97 | Unlimed ... | 50.0 | 8.0 | 42.0 | Plantago lanceolata; Centaurea nigra; Leontodon hispidus; Poterium sanguisorba | 5-1 |
| 5-2 | (S. half), Super., Sulphate of Potash; following double dressing of Amm. Salts (=86 lb. N.), 1856-97 | Unlimed ... | 68.0 | 1.7 | 30.3 | Centaurea nigra; Scabiosa arvensis; Rumex acetosa | 5-2 |
| 7 | Complete Mineral Manure | Unlimed ... | 57.3 | 17.6 | 25.1 | Rumex acetosa; Centaurea nigra; Luzula campestris; Achillea millefolium | 7 |
| 9 | Complete Mineral Manure and double Amm. Salts | Limed ... | 36.9 | 51.8 | 11.3 | Heracleum sphondylium; Centaurea nigra | 9 |
| 14 | Complete Mineral Manure and double Nitrate of Soda | Unlimed ... | 47.1 | 33.3 | 19.6 | Plantago lanceolata; Heracleum sphondylium; Conopodium denudatum; Achillea millefolium | 14 |
| 18 | Mineral Manure (without Super.) and double Sulphate Amm., 1905 and since | Limed ... | 98.8 | 0.1 | 1.0 | Conopodium denudatum; Rumex acetosa | 18 |
| 19 | Farmyard Dung in 1905 and every 4th year since (omitted in 1917) | Unlimed ... | 98.7 | 0.2 | 1.0 | Heracleum sphondylium; Potentilla reptans | 19 |
| 20 | Farmyard Dung in 1905 and every 4th year since (omitted in 1917) each intervening year Sulphate of Potash, Super. and Nitrate of Soda | Limed ... | 84.9 | 6.2 | 8.9 | Taraxacum vulgare; Anthriscus sylvestris | 20 |
| | | Unlimed ... | 90.2 | 0.4 | 9.4 | Anthriscus sylvestris; Taraxacum vulgare | |
| | | L. 6,788 lb. | 91.8 | 0.1 | 8.1 | Rumex acetosa; Conopodium denudatum | |
| | | L. 3,951 lb. | 86.6 | 0.2 | 13.2 | Rumex acetosa; Conopodium denudatum; Centaurea nigra | |
| | | Unlimed ... | 86.2 | — | 13.8 | Heracleum sphondylium; Rumex acetosa | |
| | | L. 3,150 lb. | 66.9 | 19.7 | 13.3 | Ranunculus sp.; Taraxacum vulgare; Conopodium denudatum | |
| | | L. 570 lb. | 69.0 | 21.6 | 9.4 | Ranunculus sp.; Conopodium denudatum; Cerastium vulgatum | |
| | | Unlimed ... | 66.7 | 20.0 | 13.3 | Ranunculus sp.; Conopodium denudatum; Centaurea nigra; Rumex acetosa | |
| | | L. 2,772 lb. | 65.5 | 23.4 | 11.1 | Taraxacum vulgare; Anthriscus sylvestris; Ranunculus sp. | |
| | | L. 570 lb. | 57.8 | 30.5 | 11.7 | Ranunculus sp.; Taraxacum sp. | |
| | | Unlimed ... | 71.2 | 16.8 | 12.0 | Ranunculus sp.; Centaurea nigra | |

WHEAT. BROADBALK FIELD, 1925.

| Plot. | Manurial Treatment. | Top Portion. | | | | | | Bottom Portion. | | | | | | 74 year Average 1852-1925. | |
|-------|--|-----------------|--------------------|----------------------|-----------------|-----------------------|-----------------------------------|-----------------|--------------------|----------------------|-----------------|-----------------------|-----------------------------------|----------------------------|-----------------------|
| | | Dressed Grain. | | Ofal Grain per Acre. | Straw per Acre. | Total Straw per Acre. | Proportion of Total Grain to 100. | Dressed Grain. | | Ofal Grain per Acre. | Straw per Acre. | Total Straw per Acre. | Proportion of Total Grain to 100. | Dressed Grain per Acre. | Total Straw per Acre. |
| | | Yield per Acre. | Weight per Bushel. | | | | | Yield per Acre. | Weight per Bushel. | | | | | | |
| | | bush. | lb. | lb. | lb. | lb. | bush. | lb. | lb. | lb. | lb. | lb. | bush. | cwt. | |
| 2A | Farmyard Manure ... | 10.5 | 58.4 | 88 | 1500 | 17.7 | 35.3 | 14.9 | 58.5 | 82 | 1591 | 19.1 | 44.6 | 26.8* | 32.1* |
| 2B | Farmyard Manure ... | 15.1 | 59.1 | 151 | 1807 | 21.3 | 43.9 | 19.1 | 58.6 | 228 | 1907 | 22.8 | 52.9 | 33.5 | 34.2 |
| 3 | Unmanured ... | 6.7 | 58.8 | 49 | 518 | 5.8 | 68.3 | 5.7 | 58.1 | 37 | 569 | 6.5 | 50.8 | 11.7 | 9.8 |
| 5 | Complete Mineral Manure ... | 6.8 | 58.8 | 68 | 502 | 5.6 | 74.4 | 6.8 | 58.5 | 51 | 462 | 5.3 | 76.7 | 13.5 | 11.5 |
| 6 | As 5, and Single Amm. Salts ... | 10.1 | 58.7 | 87 | 707 | 8.1 | 74.7 | 10.1 | 58.7 | 80 | 784 | 9.2 | 65.5 | 21.7 | 20.3 |
| 7 | As 5, and Double Amm. Salts ... | 18.6 | 59.2 | 93 | 1558 | 17.9 | 59.6 | 21.4 | 54.7 | 100 | 1768 | 20.0 | 56.7 | 30.4 | 32.1 |
| 8 | As 5, and Treble Amm. Salts ... | 19.5 | 59.7 | 106 | 2182 | 25.0 | 45.5 | 21.7 | 59.0 | 95 | 1868 | 22.0 | 56.0 | 34.5 | 39.8 |
| 9 | As 5, and Single Nitrate of Soda ... | 16.3 | 58.2 | 45 | 1362 | 15.9 | 55.7 | 16.0 | 57.0 | 55 | 1534 | 17.6 | 49.1 | 18.8† | 24.6† |
| 10 | Double Amm. Salts alone ... | 14.0 | 59.2 | 138 | 1162 | 13.6 | 63.6 | 10.6 | 58.5 | 126 | 797 | 10.1 | 66.3 | 18.7 | 17.8 |
| 11 | As 10, and Superphosphate ... | 20.5 | 58.3 | 143 | 1558 | 17.7 | 62.3 | 16.9 | 57.2 | 142 | 1042 | 13.2 | 75.0 | 21.3 | 21.4 |
| 12 | As 10, and Super. and Sulph. Soda ... | 18.8 | 59.1 | 189 | 1496 | 17.3 | 67.0 | 18.0 | 58.6 | 189 | 1698 | 19.7 | 56.2 | 27.0 | 26.8 |
| 13 | As 10, and Super. and Sulph. Potash ... | 24.3 | 59.4 | 87 | 1832 | 21.4 | 63.8 | 22.2 | 59.0 | 64 | 2192 | 24.4 | 50.2 | 29.2 | 30.6 |
| 14 | As 10, and Super. and Sulph. Magnesia ... | 20.2 | 58.5 | 77 | 1556 | 17.9 | 62.8 | 21.7 | 58.9 | 78 | 2275 | 24.3 | 49.9 | 26.7 | 26.8 |
| 15 | Double Amm. Salts in Autumn and Minerals ... | 20.6 | 59.7 | 64 | 1460 | 16.6 | 69.6 | 16.3 | 59.5 | 66 | 1184 | 13.9 | 66.6 | 27.8 | 28.2 |
| 16 | Double Nitrate and Minerals ... | 21.2 | 59.5 | 104 | 2002 | 22.7 | 53.7 | 22.0 | 59.6 | 118 | 2175 | 24.5 | 52.0 | 29.9† | 35.2† |
| 17) | Minerals alone, or double Amm. Salts alone in | 9.7 | 59.6 | 68 | 624 | 7.1 | 81.1 | 10.7 | 60.0 | 56 | 692 | 8.0 | 77.9 | 27.8 | 27.7 |
| 18) | alternate years ... | 15.7 | 60.0 | 133 | 1272 | 14.6 | 65.8 | 14.2 | 59.8 | 157 | 1510 | 16.9 | 53.2 | 14.1 | 12.5 |
| 19 | Rape Cake alone ... | 12.6 | 59.7 | 55 | 1102 | 12.5 | 57.7 | 6.7 | 58.9 | 43 | 971 | 11.1 | 35.5 | 20.8† | 22.0† |
| 20 | Mineral Manure (without Super.) and Amm. Salts | 7.7 | 60.0 | 47 | 1045 | 11.6 | 39.1 | — | — | — | — | — | — | 16.5§ | 18.6§ |

* 26 years only, 1900-1925. † 41 years only, 1885-1925. ‡ 33 years only, 1893-1925. § 18 years only, 1906-1925 (no crop in 1912 and 1914).

WHEAT. BROADBALK FIELD, 1926.
Top portion fallowed.

| Plot | Manurial Treatment | Dressed Grain | | Offal Grain per Acre | Straw per Acre | Total Straw per Acre | Proportion of Total Grain to 100 of Total Straw |
|------|---|----------------------|-----------------------|----------------------|----------------|----------------------|---|
| | | Yield per Acre bush. | Weight per Bushel lb. | | | | |
| 2A | Farmyard Manure | 6.8 | 54.8 | 113 | 1979 | 24.6 | 17.6 |
| 2B | Farmyard Manure | 6.5 | 55.5 | 133 | 2675 | 33.6 | 13.2 |
| 3 | Unmanured | 0.9 | 57.5* | 9 | 135 | 1.8 | 30.2 |
| 5 | Complete Mineral Manure | 2.2 | 57.5 | 17 | 285 | 3.5 | 38.8 |
| 6 | As 5, and Single Amm. Salts | 5.9 | 56.8 | 50 | 1030 | 13.0 | 26.5 |
| 7 | As 5, and Double Amm. Salts | 5.7 | 55.1 | 91 | 1985 | 23.3 | 15.4 |
| 8 | As 5, and Treble Amm. Salts | 7.5 | 50.4 | 118 | 2973 | 33.5 | 13.2 |
| 9 | As 5, and Single Nitrate of Soda | 5.8 | 54.0 | 72 | 1293 | 16.0 | 21.8 |
| 10 | Double Amm. Salts alone | 4.4 | 51.3 | 84 | 1030 | 12.5 | 22.0 |
| 11 | As 10, and Superphosphate | 4.2 | 53.0 | 113 | 1360 | 17.7 | 16.8 |
| 12 | As 10, and Super. and Sulph. Soda | 7.1 | 54.1 | 149 | 1733 | 21.7 | 21.9 |
| 13 | As 10, and Super. and Sulph. Potash | 9.3 | 56.3 | 123 | 2205 | 26.4 | 21.7 |
| 14 | As 10, and Super. and Sulph. Magnesia | 8.6 | 54.6 | 135 | 1838 | 22.7 | 24.1 |
| 15 | Double Amm. Salts in Autumn and Minerals | 5.5 | 56.4 | 107 | 1408 | 18.9 | 20.4 |
| 16 | Double Nitrate and Minerals | 7.5 | 54.4 | 141 | 2283 | 27.8 | 17.8 |
| 17 | Minerals alone or Double Amm. | 6.4 | 56.0 | 88 | 1508 | 18.0 | 22.9 |
| 18 | Salts alone in alternate years | 3.6 | 56.0 | 60 | 668 | 9.0 | 27.2 |
| 19 | Rape Cake alone | 4.4 | 53.4 | 98 | 1503 | 17.6 | 16.6 |

* Adopted from plot 5.

RED CLOVER grown year after year on rich Garden Soil,
Rothamsted Garden.

Hay, Dry Matter, and Nitrogen per Acre, 1925 and 1926.

| Year | No. of Cuttings | As Hay | Dry Matter | Nitrogen | Seed Sown |
|---------------------|-----------------|--------|------------|----------|--|
| 1925 | 2 | 1525 | 1270 | 33 | April 17th, Re-seeded June 1st, Patched |
| 1926 | 2 | 1248 | 1040 | 32 | |
| Averages: | | | | | |
| 25 years, 1854—1878 | | 7664 | 6387 | 179 | |
| 25 years, 1879—1903 | | 3924 | 3270 | 101 | |
| 20 years, 1904—1923 | | 2640 | 2200 | 65 | |

WHEAT AFTER FALLOW (without Manure 1851,
and since).

Hoos Field, 1925 and 1926.

| | 1925 | 1926 | Average 70 years 1856-1925 |
|---|-------|-------|----------------------------------|
| Dressed Grain { Yield per Acre—bushels | 5.9 | 5.24 | 14.70 |
| { Weight per Bushel—lb. | 58.9 | 58.2 | 58.8 |
| Offal Grain per Acre—lb. | 33.5 | 96.0 | 50.7 |
| Straw per Acre—lb. | 623.0 | 780.0 | — |
| Total Straw per Acre—cwt. | 6.8 | 9.0 | 12.7 |
| Proportion of Total Grain to 100 of Total Straw | 49.8 | 39.7 | — |

AVERAGE WHEAT YIELDS of VARIOUS COUNTRIES.

| Country | Mean Yield per Acre 1901-10 bushels | Country | Mean Yield per Acre 1901-10 bushels |
|----------------------|-------------------------------------|------------------------|-------------------------------------|
| Great Britain | 31.6 | Denmark | 41.3 |
| England | 31.7 | Argentina | 10.6 |
| Hertfordshire | 30.5 | Australia | 10.1 |
| France | 20.2 | Canada | 19.5 |
| Germany | 29.1 | United States | 14.3 |
| Belgium | 35.1 | Russia—European | 10.0 |

NOTE.—Figures for Great Britain, England and Hertfordshire are taken from the Board of Agriculture's "Agricultural Statistics," Vol. 46. Other figures from "Annuaire International de Statistique Agricole," 1910-12, and converted at the rate of 60 lb. per bushel.

PERMANENT BARLEY PLOTS. Hoos Field, 1925, 1926.
PRODUCE PER ACRE.

| Plot. | Manuring. | 1925. | | | | | | 1926. | | | | | | 74 years Average Yield 1852-1926. † | |
|-------|---|-----------------------|---------------------------|--------------------------|--------------------|--------------------------|--|-----------------------|---------------------------|--------------------------|--------------------|--------------------------|--|--|--------------------------|
| | | Yield per Acre. | Dressed Grain. Bushel. | Offal Grain per Acre. | Straw per Acre. | Total Straw per Acre. | Proportion of Total Grain to 100 of Total Straw. | Yield per Acre. | Dressed Grain. Bushel. | Offal Grain per Acre. | Straw per Acre. | Total Straw per Acre. | Proportion of Total Grain to 100 of Total Straw. | Dressed Grain per Acre. | Total Straw per Acre. |
| 1 O | Unmanured ... | 6.7 | 50.8 | 33 | 396 | 5.0 | 66.9 | 6.1 | 51.5 | 30 | 382 | 7.1 | 42.9 | 13.6 | 7.9 |
| 2 O | Superphosphate only ... | 10.9 | 52.3 | 44 | 594 | 7.6 | 72.5 | 12.2 | 53.1 | 32 | 569 | 8.3 | 72.6 | 19.2 | 9.8 |
| 3 O | Alkali Salts only ... | 5.0 | 50.0 | 26 | 355 | 4.5 | 54.0 | 4.8 | 50.8 | 41 | 374 | 5.7 | 39.6 | 14.5 | 8.6 |
| 4 O | Complete Minerals ... | 7.1 | 51.8 | 43 | 470 | 6.3 | 58.0 | 12.9 | 52.5 | 135 ^a | 875 | 13.2 | 55.3 ^a | 19.3 | 11.0 |
| 5 O | Potash and Superphosphate ... | 8.1 | 52.3 | 33 | 451 | 5.9 | 69.4 | 9.9 | 52.5 | 39 | 622 | 9.6 | 51.9 | 15.7 | 9.5 |
| 1 A | Ammonium Salts only ... | 9.4 | 49.5 | 39 | 693 | 8.5 | 53.3 | 12.0 | 51.9 | 52 | 836 | 11.9 | 50.5 | 24.0 | 13.8 |
| 2 A | Superphosphate and Amm. Salts ... | 19.0 | 52.5 | 147 | 1037 | 13.2 | 77.5 | 26.1 | 52.1 | 63 | 1546 | 18.9 | 67.4 | 36.4 | 20.7 |
| 3 A | Alkali Salts and Amm. Salts ... | 11.0 | 51.8 | 55 | 864 | 10.9 | 51.2 | 11.3 | 50.6 | 65 | 1009 | 13.0 | 43.8 | 26.2 | 16.1 |
| 4 A | Complete Minerals and Amm. Salts ... | 19.3 | 51.7 | 116 | 1372 | 16.6 | 59.6 | 30.1 | 51.7 | 74 | 2054 | 23.6 | 61.7 | 39.9 | 23.8 |
| 5 A | Potash, Super. and Amm. Salts ... | 21.8 | 53.3 | 122 | 1436 | 17.5 | 65.4 | 24.2 | 53.0 | 50 | 1645 | 20.7 | 57.4 | 34.4 | 21.9 |
| 1 AA | Nitrate of Soda only ... | 12.6 | 50.3 | 52 | 825 | 10.2 | 60.0 | 15.9 | 52.9 | 62 | 1084 | 16.0 | 50.6 | 24.5* | 15.4* |
| 2 AA | Super. and Nitrate of Soda ... | 29.9 | 53.7 | 165 | 1623 | 20.2 | 78.3 | 31.0 | 52.1 | 78 | 1986 | 23.6 | 64.1 | 39.3* | 23.3* |
| 3 AA | Alkali Salts and Nitrate of Soda ... | 10.0 | 50.0 | 76 | 803 | 10.4 | 49.4 | 10.3 | 50.3 | 70 | 1051 | 16.8 | 31.4 | 24.9* | 16.5* |
| 4 AA | Complete Minerals and Nitrate of Soda ... | 18.7 | 53.0 | 96 | 1342 | 15.5 | 62.5 | 27.9 | 51.0 | 89 | 2167 | 24.5 | 55.1 | 38.2* | 23.7* |
| 1 AAS | As Plot 1 AA and Silicate of Soda ... | 13.8 | 52.0 | 69 | 941 | 12.1 | 58.0 | 21.0 | 53.3 | 70 | 1359 | 17.7 | 59.9 | 30.5* | 18.4* |
| 2 AAS | " " 2 AA " " " " | 26.3 | 53.6 | 124 | 1381 | 17.5 | 78.2 | 37.8 | 52.0 | 94 | 2316 | 27.7 | 66.4 | 40.3* | 24.2* |
| 3 AAS | " " 3 AA " " " " | 12.5 | 52.3 | 96 | 963 | 12.2 | 54.8 | 16.6 | 52.3 | 107 | 1271 | 17.9 | 48.7 | 31.7* | 20.1* |
| 4 AAS | " " 4 AA " " " " | 17.9 | 52.9 | 66 | 1364 | 16.8 | 53.6 | 35.0 | 51.5 | 95 | 2299 | 27.0 | 62.8 | 40.6* | 25.7* |
| 1 C | Rape Cake only ... | 24.5 | 52.9 | 87 | 1955 | 18.1 | 68.3 | 24.5 | 52.4 | 60 | 1559 | 19.1 | 62.9 | 35.9 | 20.7 |
| 2 C | Superphosphate and Rape Cake ... | 21.9 | 54.4 | 128 | 1331 | 15.8 | 74.8 | 33.6 | 51.5 | 63 | 2019 | 23.4 | 68.4 | 38.4 | 22.1 |
| 3 C | Alkali Salts and Rape Cake ... | 12.7 | 52.8 | 99 | 1001 | 12.5 | 54.7 | 20.4 | 52.2 | 39 | 1570 | 18.8 | 52.3 | 34.2 | 20.6 |
| 4 C | Complete Minerals and Rape Cake ... | 21.3 | 53.2 | 85 | 1298 | 15.5 | 69.9 | 34.6 | 52.0 | 64 | 2107 | 25.5 | 65.2 | 38.0 | 22.8 |
| 7-1 | Unmanured (after dung 20 years, 1852-71) | 7.0 | 51.5 | 76 | 475 | 6.3 | 61.7 | 11.0 | 53.3 | 48 | 725 | 10.9 | 51.6 | 22.8† | 13.7† |
| 7-2 | Farmyard Manure ... | 22.0 | 52.3 | 121 | 1158 | 15.9 | 71.4 | 35.8 | 52.1 | 88 | 2331 | 27.6 | 63.3 | 45.1 | 28.1 |
| 6-1 | Unmanured ... | 5.7 | 50.5 | 54 | 354 | 4.9 | 62.6 | 7.1 | 51.5 | 50 | 485 | 7.8 | 47.3 | 14.9 | 8.7 |
| 6-2 | Ashes from Laboratory furnace ... | 7.5 | 51.0 | 36 | 431 | 5.6 | 66.4 | 9.6 | 52.4 | 43 | 620 | 8.7 | 55.6 | 15.9 | 9.4 |
| 1 N | Nitrate of Soda only ... | 11.8 | 51.8 | 80 | 820 | 11.3 | 54.7 | 14.3 | 52.0 | 70 | 1078 | 16.0 | 45.4 | 29.0\$ | 18.0\$ |
| 2 N | " " " " " " | 16.8 | 53.3 | 63 | 1172 | 14.0 | 61.2 | 20.0 | 52.5 | 85 | 1436 | 19.6 | 51.6 | 32.1\$\$ | 20.1\$\$ |

* 58 years, 1868-1926. † 54 years, 1872-1926. §§ 67 years, 1853-1926. \$ 73 years, 1853-1926. \$ 67 years, 1859-1926.
^a A large amount of black medic seed in Offal Grain.

Little Hoos Field. Swedes, 1926.
Produce per acre. Roots and Leaves in Tons.

| Manurial Treatment | | Roots | Leaves | Total | Season of last Dressing |
|--------------------|--|---|--------------|--------------|-------------------------|
| A 1 | Control | <i>12.61</i> | <i>2.31</i> | <i>14.92</i> | — |
| 2 | Ordinary Dung, 16 tons | 21.79 | 3.94 | 25.73 | 1926 |
| 3 | | 11.46 | 2.87 | 14.33 | 1921 |
| 4 | | 8.25 | 2.44 | 10.69 | 1922 |
| 5 | | 9.20 | 2.53 | 11.73 | 1924 |
| B 1 | Cake-fed Dung, 16 tons | 21.11 | 3.75 | 24.86 | 1926 |
| 2 | Control | <i>13.30</i> | <i>2.62</i> | <i>15.92</i> | — |
| 3 | Cake-fed Dung, 16 tons | 14.95 | 2.99 | 17.94 | 1921 |
| 4 | | 13.88 | 3.09 | 16.97 | 1922 |
| 5 | | 12.74 | 2.94 | 15.68 | 1924 |
| C 1 | | Shoddy; Superphosphate; Sulphate of Potash | 16.74 | 3.01 | 19.75 |
| 2 | Control | <i>13.44</i> | <i>2.62</i> | <i>16.06</i> | 1921 |
| 3 | Control | <i>10.28</i> | <i>2.32</i> | <i>10.60</i> | — |
| 4 | Shoddy; Superphosphate | 5.72 | 1.56 | 7.28 | 1922 |
| 5 | Sulphate of Potash | 1.87 | 0.56 | 2.43 | 1924 |
| D 1 | Guano; Sulphate of Ammonia | 17.31 | 3.20 | 20.51 | 1926 |
| 2 | Sulphate of Potash | 13.71 | 2.68 | 16.39 | 1921 |
| 3 | Control | <i>12.96</i> | <i>2.79</i> | <i>15.75</i> | 1922 |
| 4 | Control | <i>11.34</i> | <i>2.36</i> | <i>13.70</i> | — |
| 5 | | Guano; Sulphate of Ammonia, Sulphate of Potash | 13.79 | 3.41 | 17.20 |
| E 1 | Rape Dust; Superphosphate | 16.86 | 2.89 | 19.75 | 1926 |
| 2 | Sulphate of Potash | 11.64 | 2.55 | 14.19 | 1921 |
| 3 | Control | 8.71 | 2.08 | 10.79 | 1922 |
| 4 | Control | 14.36 | 2.62 | 16.98 | 1924 |
| 5 | Control | <i>10.81</i> | <i>2.42</i> | <i>13.22</i> | — |
| F 1 | Control | <i>7.20</i> | <i>1.78</i> | <i>8.98</i> | — |
| 2 | Superphosphate; Sulphate of Ammonia; Sulphate of Potash | 15.54 | 2.85 | 18.39 | 1926 |
| 3 | | 5.95 | 1.40 | 7.35 | 1921 |
| 4 | | 6.59 | 1.41 | 8.00 | 1922 |
| 5 | | 11.60 | 2.00 | 13.60 | 1924 |
| G 1 | Bone Meal; Sulphate of Ammonia; Sulphate of Potash | 14.46 | 2.97 | 17.43 | 1926 |
| 2 | Control | 7.08 | 1.88 | 8.96 | 1921 |
| 3 | Control | <i>3.86</i> | <i>1.09</i> | <i>4.95</i> | — |
| 4 | Bone Meal; Sulphate of Ammonia; Sulphate of Potash | 6.84 | 1.75 | 8.59 | 1922 |
| 5 | Control | 8.89 | 2.02 | 10.91 | 1924 |
| H 1 | Basic Slag; Sulphate of Ammonia; Sulphate of Potash | 13.40 | 2.08 | 15.48 | 1926 |
| 2 | Control | 9.50 | 1.88 | 11.38 | 1921 |
| 3 | Control | 9.47 | 1.94 | 11.41 | 1922 |
| 4 | Control | 9.88 | 1.85 | 11.73 | 1924 |
| 5 | Control | <i>4.76</i> | <i>1.47</i> | <i>6.23</i> | — |

1925, field fallowed.

NOTES.—Since 1919 the manure for each plot (except of series A and B) has been rationed at 40 lb. Nitrogen, 100 lb. Calcium Phosphate and 50 lb. Potash per acre. Each plot has been supplied with as much of its particular manure (shoddy, guano, etc.) as possible without exceeding the receipt in any of the three rationed ingredients. Any deficit in either of these three has been made good by adding the necessary quantity of Sulphate of Ammonia, Superphosphate, or Sulphate of Potash. No manure was applied for 1923 crop.

Figures in italics denote unmanured plots. The yield on the plots to which the manure was applied in a given season are printed in heavy type.

Hay. Great Field, 1925 and 1926.

| Plot. | Manurial Treatment. Quantities per Acre. | Yield per Acre. | | Yield per Acre. | | Dry Matter per Acre. | |
|-------|---|-----------------------|-------------------------|-----------------------|-------------------------|-------------------------|------------------------|
| | | 1925. | | 1926. | | 1926.* | |
| | | No Potash. cwt. | With Potash. cwt. | No Potash. cwt. | With Potash. cwt. | No Potash. lb. | With Potash. lb. |
| 1 A | High Grade Slag, No. 12, 1,170 lb. | 38.2 | 34.8 | 41.6 | 40.4 | 3628 | 3519 |
| 1 B | | 48.4 | 42.9 | 43.2 | 37.5 | 3776 | 3381 |
| 2 A | Open Hearth Slag, No. 13, 1,925 lb. | 36.3 | 37.9 | 36.3 | 42.3 | 3159 | 3741 |
| 2 B | | 45.0 | 35.0 | 37.3 | 39.5 | 3214 | 3688 |
| 3 A | Open Hearth Slag, No. 14, 1,930 lb. | 39.8 | 34.3 | 35.5 | 38.4 | 3198 | 3336 |
| 3 B | | 40.7 | 32.3 | 37.5 | 40.9 | 3384 | 3730 |
| 4 A | Gafsa Phosphate 750 lb. ... | 47.0 | 32.7 | 39.6 | 41.1 | 3358 | 4129 |
| 4 B | | 42.5 | 32.7 | 37.3 | 42.3 | 3252 | 3940 |
| A C | Control. No Manure ... | 37.0 | 34.1 | 31.8 | 43.0 | 2853 | 3648 |
| B C | | 45.2 | 35.7 | 40.2 | 38.2 | 3154 | 3397 |
| 7 C | Nauru Phosphate 263 lb. ... | 37.1 | 35.5 | — | — | — | — |
| 7 D | | 33.6 | 32.9 | — | — | — | — |
| 8 C | Nauru Slag Phosphate, No. 8, 411 lb. | 36.4 | 31.3 | — | — | — | — |
| 8 D | | 30.7 | 31.4 | — | — | — | — |
| 1 C | High Soluble Slag, No. 1, 872 lb. | 33.6 | 38.8 | — | — | — | — |
| 2 C | Low Soluble Slag, No. 2, 1,225 lb. | 30.7 | 33.4 | — | — | — | — |
| 3 C | Gafsa Phosphate, 347 lb. ... | 30.5 | 36.1 | — | — | — | — |
| 4 C | Tunisian Phosphate, 336 lb. ... | 33.4 | 34.8 | — | — | — | — |
| 5 C | Florida Phosphate, 292 lb. ... | 36.4 | 35.5 | — | — | — | — |
| C C | Control. No Manure ... | 27.9 | 32.0 | — | — | — | — |
| D C | | 30.0 | 27.1 | — | — | — | — |

Kainit at 4 cwt. per acre, applied January 28th, 1924.
 * Dry Matter determinations were not made in 1925.
 Series C and D were discarded in 1926.

Great Knott Field, 1926.

Produce per Acre.

| Wheat Varieties | Dressed Yield per Acre. bush. | Grain Weight per bush. lb. | Straw per Acre lb. | Total Straw per Acre. cwt. | Proportion of Total Grain to 100 Total Straw |
|--------------------|--|-------------------------------------|--------------------------|-------------------------------------|---|
| Red Standard ... | 30.7 | 61.4 | 3105 | 31.4 | 54.6 |
| Browick A ... | 36.8 | 58.7 | 4118 | 42.8 | 49.4 |
| Browick B ... | 36.2 | 57.7 | 3406 | 35.5 | 53.5 |
| Little Joss A ... | 45.9 | 62.6 | 4795 | 48.3 | 55.5 |
| Little Joss B ... | 46.5 | 61.8 | 4630 | 47.2 | 57.4 |
| R. Million A ... | 37.1 | 61.4 | 3900 | 43.5 | 48.9 |
| R. Million B ... | 37.4 | 61.2 | 3224 | 38.9 | 54.8 |