


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Report for 1934

[Full Table of Content](#)



The Classical Experiments

Rothamsted Research

Rothamsted Research (1935) *The Classical Experiments* ; Report For 1934, pp 135 - 162 - DOI:
<https://doi.org/10.23637/ERADOC-1-66>

CROPS GROWN IN ROTATION, AGDELL FIELD

PRODUCE PER ACRE.

Year.	Crop.	O. Unmanured since 1848.		M. Mineral Manure.‡ No Nitrogen.		C. Complete Mineral and Nitrogenous Manure	
		5. Fallow.	6. Clover or Beans.	3. Fallow.	4. Clover or Beans.	1. Fallow.	2. Clover or Beans.
Average of first twenty-one Courses, 1848-1931							
	Roots (Swedes) .. cwt.*	32.0	16.1	174.0	206.5	352.0	310.0
	Barley—						
	Dressed Grain .. bush.	21.6	19.8	22.7	26.6	30.3	35.0
	Total Straw .. cwt.†	13.3	13.2	13.6	15.6	18.4	21.7
	Beans—						
	Dressed Grain .. bush.‡‡	—	13.1	—	18.2	—	22.3
	Total Straw .. cwt.‡‡	—	9.2	—	13.2	—	15.3
	Clover Hay .. cwt.§	—	25.6	—	52.1	—	52.0
	Wheat—						
	Dressed Grain .. bush.	23.1	21.6	26.9	29.4	27.5	29.0
	Total Straw .. cwt.†	22.9	21.2	28.2	29.8	29.4	29.3
Present Course (22nd), 1932-34							
1932	Roots (Turnips) .. cwt.	20.2	5.4	86.0	118.0	120.0	98.6
1933	Barley—						
	Dressed Grain .. bush.	6.0	2.2	9.5	13.9	3.7	5.4
	Total Grain .. cwt.	3.3	1.3	5.2	7.4	2.0	2.9
	Weight per bushel lb.	54.8	50.2	55.2	55.0	52.9	53.0
	Total Straw .. cwt.†	6.3	4.8	7.4	11.4	9.1	14.0
1934	Beans—						
	Dressed Grain .. bush	—	9.2	—	24.6	—	13.1
	Total Grain .. cwt.	—	5.6	—	15.2	—	8.1
	Weight per bushel lb.	—	67.8	—	69.4	—	69.6
	Total Straw .. cwt.	—	10.5	—	28.3	—	16.5

* Plots 1, 3 and 5 based upon 19 courses. Plots 2, 4 and 6 based upon 18 courses.

† Includes straw, cavings and chaff.

‡ Mineral Manure : 528 lb. Superphosphate (35%) ; 500 lb. Sulphate of Potash ; 100 lb. Sulphate of Soda ; 200 lb. Sulphate of Magnesia, all per acre. Nitrogenous Manure—206 lb. Sulphate of Ammonia and 2,000 lb. Rape Dust per acre. Manures applied once every four years, prior to sowing of Swedes.

‡‡ Based on 8 courses.

§ Based on 13 courses.

CULTIVATIONS, ETC.—Cropped sections: Clover sown : May 3. Variety: Broad Red. Clover failed. Ploughed : Feb. 5 and 6. Harrowed : Feb. 21. Seed sown : Feb. 21. Variety: Spring beans. Hoed : May 14, 28 and 29. Cut : August 4. Fallow sections : Ploughed : November 29-December 1, May 25, July 22 and August 30-September 7.

WHEAT AFTER FALLOW—HOOS FIELD

Without Manure 1851, and since.

SCHEME FOR COMPARING A THREE YEAR FALLOW WITH A ONE YEAR FALLOW.

Each of the two strips on Hoos Wheat after Fallow is to be divided into four parts. In the year when a strip is in crop, one quarter is to continue to be fallowed, so that this quarter has a three-year fallow. Different quarters are to be selected for fallow in successive years in the rotation given in the following table :

A N B

Cropping of strips A and B.

C=Crop. F=Fallow.

1	1
2	2
3	3
4	4

Year.	A1.	A2.	A3.	A4.	B1.	B2.	B3.	B4.
1932	F	C	C	C	F	F	F	F
1933	F	F	F	F	C	C	F	C
1934	C	F	C	C	F	F	F	F
1935	F	F	F	F	C	C	C	F
1936	C	C	F	C	F	F	F	F
1937	F	F	F	F	F	C	C	C
1938	C	C	C	F	F	F	F	F
1939	F	F	F	F	C	F	C	C
1940	F	C	C	C	F	F	F	F

A comparison of the effect of a three year fallow with the effect of a one year fallow will be possible in ever year.

Half the experiment will continue to be wheat after one year fallow, and continuity with previous results will thus be maintained.

PRODUCE PER ACRE, 1934.

	A1.	A3.	A4.	Mean.	Average, 78 years, 1856-1933.
Dressed Grain—bushels	27.9	19.6	28.1	25.2	14.3
Total grain—cwt.	16.8	12.3	17.2	15.4	8.1
Weight per bushel—lb.	63.9	63.4	63.5	63.6	58.8
Total Straw—cwt.	24.6	15.2	19.0	19.6	12.7

CULTIVATIONS, ETC.—Cropped sections. Ploughed : September 12. Harrowed : April 7. Rolled : April 9. Seed sown : October 13. Variety : Red Standard. Harvested : August 8. Fallowed section. Ploughed : February 7. Tractor cultivated : April 20, May 4. Horse harrow : May 17. Plough : May 17. Tractor cultivated : June 16. Two horse roll : June 22. Plough : June 28. Tractor cultivated : July 24.

MANGOLDS—BARNFIELD, 1934

Roots each year since 1856.

PRODUCE PER ACRE.

Mangolds each year since 1876.

Strip	Manures (Amounts stated are per acre.)	1934						53 Year Average, 1876-1933†					
		Cross Dressings						Cross Dressings					
		O	N	A	AC	C		O	N	A	AC	C	
	None	Nitrate of Soda (550 lb.)	Sulphate of Ammonia (412 lb.)	Sulphate of Ammonia (412 lb.) & Rape Cake (2,000 lb.)	Rape Cake (2,000 lb.)		None	Nitrate of Soda (550 lb.)	Sulphate of Ammonia (412 lb.)	Sulphate of Ammonia (412 lb.) & Rape Cake (2,000 lb.)	Rape Cake (2,000 lb.)		
Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons	Tons
Roots													
1	Dung only (14 tons)	25.31	42.88	34.01	33.67	32.02	17.45	26.14	21.57	23.38	23.43		
2	Dung, Superphosphate (3½ cwt.), Sulphate of Potash (600 lb.)	33.50	49.49	39.19	41.42	37.84	19.00	26.71	24.68	27.46	26.50		
4	Complete Minerals : Super. and Potash as 2, Salt (200 lb.)		(a) 37.70**					(a) 17.41					
	Sulphate of Magnesia (200 lb.)		(b) 37.08**					(b) 18.01*					
5	Superphosphate only (3½ cwt.)		8.74	25.33	39.66	32.03	4.64	14.48	25.97	20.95			
6	Super. (3½ cwt.) Sulphate of Potash (500 lb.)		7.89	28.27	6.55	8.15	8.80	4.47	14.71	6.80	9.34	10.03	
7	Super. (3½ cwt.) Sulphate of Magnesia (200 lb.) and Sodium Chloride (200 lb.)		8.11	30.09	20.77	28.79	27.79	4.04	15.16	13.56	22.30	18.02	
8	No Minerals												
9	Sodium Chloride (200 lb.), Nit. Soda (550 lb.), Sulph. Potash (500 lb.) and Sulph. Mag. (200 lb.)		5.92	30.52	21.14	26.85	30.02	4.83	16.16	14.80	21.97	19.04	
			15.77	4.97	9.59	13.41	3.34	9.76	5.40	8.38	8.87		
			30.66	—	—	—	—	—	—	—	—	—	
Leaves													
1	Dung only (14 tons)		4.68	6.44	6.76	7.81	8.10	3.02	4.60	4.83	5.16	4.50	
2	Dung, Superphosphate (3½ cwt.), Sulphate of Potash (600 lb.)		4.60	6.92	6.18	6.90	6.87	3.13	5.11	5.41	6.18	4.77	
4	Complete Minerals : Super. and Potash as 2, Salt (200 lb.)		(a) 5.44		(b) 6.35				(a) 3.83				
	Sulph. of Magnesia (200 lb.)		2.12	4.40	6.12	4.25	1.04	(b) 4.01*	2.87	5.25	3.35		
5	Superphosphate only (3½ cwt.)		1.73	5.68	3.44	4.18	4.32	1.05	3.15	2.59	3.22	2.80	
6	Super. (3½ cwt.) Sulphate of Potash (500 lb.)		1.57	5.29	4.21	6.35	4.48	0.93	3.01	2.77	5.09	2.84	
7	Super. (3½ cwt.) Sulphate of Magnesia (200 lb.) and Sodium Chloride (200 lb.)		1.63	5.23	4.85	7.23	5.64	1.10	3.31	3.00	5.14	3.30	
8	No Minerals												
9	Sodium Chloride (200 lb.), Nit. Soda (550 lb.), Sulph. Potash (500 lb.) and Sulph. Mag. (200 lb.)		1.49	4.95	2.98	4.57	4.94	0.97	3.16	2.48	3.22	2.81	
			4.50	—	—	—	—	—	—	—	—	—	

** From 1904 onwards plot 4N has been divided, 4(a) receiving Superphosphate, Sulphate of Potash, Sulphate of Magnesia, Sodium Chloride and Nitrate of Soda, amounts as above; (4b) receiving Superphosphate, Calcium Chloride, (190 lb.), Potassium Nitrate (570 lb.), and Calcium Nitrate (100 lb.). Nitrogenous manures are applied as to one-third at time of sowing and two-thirds as top dressing at a later date, except with Rape Cake which all goes on with the seed.

† Excluding 1885 when nitrogenous fertilisers were not applied, owing to poor crop, 1908 and 1927 when the crop was swedes, 1930 when the spacing of the rows was changed and 1931 when the crop was a mixture of mangolds and swedes.

* 26 years only, 1904-1933, excluding 1908, 1927, 1930 and 1931. For this period the average yield of plot 4(a) was 18.16 for roots and 3.95 for leaves.

Cultivations, Etc.—Ploughed; November 3-10 and 25-27, Cultivated April 25 and 30 and May 1. Harrowed: May 1, 2 and 4. Rolled: May 3, 6 and 10. Hoed: May 31-June 4, June 13-16, July 16-18 and August 8 and 9. Singled: June 23-28. Seed sown: May 4. Variety: Yellow Globe. Manures applied: November 22-24, April 23-25 and July 9 and 10. Lifted: October 31; November 16.

HAY—THE PARK GRASS PLOTS, 1934

Plot.	Manuring (amounts stated are per acre).	Yield of Hay per acre.			Dry Matter per acre.		
		1st Crop	2nd* Crop	Total	1st Crop	2nd* Crop	Total
1	Single dressing (206 lb.) Sulphate of Ammonia (=43 lb. N.), (with Dung also 8 years, 1856-63)	not limed 7.2	limed 16.5		cwt. cwt. cwt.	6.4 14.2	
2	Unmanured (after Dung 8 years, 1856-63)	not limed 11.8	limed 12.4			10.3 10.6	
3	Unmanured	not limed 12.3	limed 9.4			10.6 7.8	
4-1	Superphosphate of lime (3½ cwt.)	not limed 12.3	limed 9.6			10.9 7.9	
4-2	Superphosphate of lime (3½ cwt.), and double dressing (412 lb.) Sulphate of Ammonia (=86 lb. N.)	not limed 13.7	3.7	17.4	12.3 0.5	12.8	
5-1	(N. half) Unmanured following double dressing Ammonia salts (=86 lb. N.) 1856-97	limed 29.7	2.6	32.3	26.6 0.7	27.3	
5-2	(S. half) Superphosphate (3½ cwt.) Sulphate of Potash (500 lb.) following double dressing Amm. salts (=86 lb. N.) 1856-97	not limed 8.3			7.5		
6	Complete Mineral Manure as Plot 7; following double dressing Amm. salts (=86 lb. N.) 1856-68	not limed 17.6			15.8		
7	Complete Mineral Manure: Super. (3½ cwt.); Sulphate of Potash (500lb.); Sulphate of Soda (100lb.); Sulphate Magnesia (100lb.)	not limed 25.8			22.2		
8	Mineral Manure without Potash	not limed 28.0			24.1		
9	Complete Mineral Manure and double dressing (412 lb.) Sulphate of Ammonia (=86 lb. N.)	not limed 32.7			28.0		
10	Mineral Manure (without Potash) and double dressing Amm. salts (=86 lb. N.)	not limed 17.8			15.1		
11-1	Complete Mineral Manure and treble dressing (618lb.) Sulphate of Amm. (129 lb. N.)	not limed 13.6			10.8		
11-2	As Plot 11-1 and Silicate of Soda	not limed 33.0	10.5	43.5	28.9 1.3	30.2	
12	Unmanured	not limed 48.1	3.7	51.8	41.1 1.0	42.1	
13	Dung (14 tons) in 1905, and every fourth year since (omitted 1917), Fish Guano (6 cwt.) in 1907 and every fourth year since	not limed 20.1			18.2		
14	Complete Mineral Manure and double dressing (550 lb.) Nitrate of Soda (=86 lb. N.)	not limed 33.1	4.4	37.5	29.0 1.4	30.4	
15	Complete Mineral Manure as Plot 7; following double dressing Nitrate of Soda (=86 lb. N., 1858-75)	not limed 36.1	39.1	75.2	29.7 4.3	34.0	
16	Complete Mineral Manure and single dressing (275 lb.) Nitrate of Soda (=43 lb. N.)	not limed 54.4	14.1	68.5	44.4 3.0	47.4	
17	Single dressing (275 lb.) Nitrate of Soda (43 lb. N.)	not limed 50.3	31.9	82.2	42.9 3.8	46.7	
18	Mineral Manure (without Super.), and double dressing Sulphate of Amm. (= 86 lb. N.), 1905 and since ; following Minerals and Amm. salts supplying the constituents of 1 ton of hay, 1865-1904	not limed 54.7	17.8	72.5	45.5 4.2	49.7	
19	Farmyard Dung (14 tons) in 1905 and every fourth year since (omitted in 1917), following Nitrate of Soda (=43 lb. N.) and Minerals, 1872-1904	not limed 14.9			12.8		
20	Farmyard Dung (14 tons) in 1905 and every fourth year since (omitted in 1917); each intervening year Plot 20 receives Sulphate of Potash (100 lb.); Superphosphate (200 lb.) and 1½ cwt. Nitrate of Soda (=26 lb. N.); following Nitrate of Potash and Superphosphate, 1872-1904	not limed 34.8			30.3		
		not limed 31.4			26.4		
		not limed 52.8	7.4	60.2	46.3 1.6	47.9	
		Limed Sun 50.6			44.1		
		," Shade 42.1			34.0		
		not limed 22.3			19.5		
		limed 17.9			16.0		
		not limed 39.2			34.0		
		limed 31.2			28.7		
		not limed 22.5			18.5		
		limed 23.1			21.4		
		not limed 13.6			12.1		
		limed (6788 lb.) 26.6			22.8		
		limed (3951 lb.) 21.0			18.2		
		not limed 24.2			23.1		
		limed (3150 lb.) 24.5			21.2		
		limed (570 lb.) 26.0			21.8		
		not limed 36.6			31.6		
		limed (2772 lb.) 35.6			30.6		
		limed (570 lb.) 39.3			32.2		

Ground Lime was applied to the southern portion (limed) of the plots at the rate of 2,000 lb. to the acre in the Winters of 1903-4, 1907-8, 1915-16, 1923-24, 1927-28, 1931-32, and at the rate of 2,500 lb. to the acre in the Winter of 1920-1921, except where otherwise stated.

Up to 1914 the limed and unlimed plot results were not separately given in the Annual Report but the mean of the two was given. From 1915 onwards the separate figures are given.

*All plots cut but only those weighed as shown. Other plots were too small to weigh.

CULTIVATIONS, ETC.—Harrowed: Feb. 20-23. Rolled: Feb. 24-26. Mannres applied: Mar. 5-7 Apr. 4-5 and 27. Cut: 1st crop, June 26-28; 2nd crop, Jan. 15-17.

PARK GRASS PLOTS

In 1914 and 1919 complete botanical analyses were made of all the Park Grass Plots, and were compared with the earlier analyses carried out by Sir John Lawes, the results being published in the Rothamsted Monograph "Manuring of Grassland for Hay." Since 1919 special attention has been given to the effect of season, and of changes in manurial treatment, on the botanical composition of the hay. The necessary information has been gained by making complete botanical analyses of certain typical plots over periods of consecutive years or in alternate years, and the data have been dealt with in a paper published in the Annals of Applied Biology, May, 1935, in which skeleton tables only are given. The complete figures for all the plots thus analysed from 1920-1933 are given in the present report. In future it is intended to publish the complete analyses year by year as they become available, in addition to the partial analyses which have hitherto appeared.

BOTANICAL COMPOSITION PER CENT—1931 (1st Crop)

Plot	Manuring	Liming.	Gramineae.	Legumi-nosae.	Other Orders.	"Other Orders" consist largely of
3	Unmanured	Limed Unlimed	59.1 60.9	12.1 10.1	28.8 29.0	(None dominant) <i>Pimpinella saxifraga</i>
7	Complete Mineral Manure.	Limed Unlimed	76.9 60.2	9.4 18.7	13.7 21.1	<i>Heracleum sphondylium</i> <i>Achillea millefolium</i>
9	Complete Mineral Manure and double Amm. Salts.	Limed Unlimed	99.0 99.5	— —	1.0 0.5	— —
14	Complete Mineral Manure and double Nitrate of Soda.	Limed (sun) Limed (shade) Unlimed	94.5 96.1 99.2	3.5 2.8 0.03	2.0 1.1 0.77	<i>Taraxacum vulgare</i> <i>Anthriscus sylvestris</i> <i>Anthriscus sylvestris</i>
15	As plot 7 following double Nitrate of Soda, 1858-75.	Limed Unlimed	81.3 80.2	5.8 7.7	12.9 12.1	<i>Plantago lanceolata</i> <i>Achillea millefolium</i> <i>Achillea millefolium</i>
17	Single Nitrate of Soda.	Limed Unlimed	78.5 80.4	3.3 0.4	18.2 19.2	<i>Plantago lanceolata</i> <i>Plantago lanceolata</i> <i>Centaurea nigra</i>
18	Mineral Manure (without Super) and double Sulphate Amm. 1905 and since.	L. 6,788 lb. L. 3,951 lb. Unlimed	90.0 89.3 96.2	0.3 — —	9.7 10.7 3.8	<i>Heracleum sphondylium</i> <i>Taraxacum vulgare</i> <i>Centaurea nigra</i> <i>Rumex acetosa</i>
19	Farmyard Dung in 1905 and every fourth year since (omitted 1917).	L. 3,150 lb. L. 570 lb. Unlimed	87.7 83.5 88.6	3.1 7.6 3.3	9.2 8.9 8.1	<i>Cerastium vulgatum</i> <i>Stellaria graminea</i> <i>Achillea millefolium</i>
20	Farmyard Dung in 1905 and every fourth year since (omitted in 1917); each intervening year Sulphate of potash, Super., and Nitrate of Soda.	L. 2772 lb. L. 570 lb. Unlimed	80.8 86.9 84.8	8.8 3.9 3.7	10.4 9.2 11.5	<i>Cerastium vulgatum</i> <i>Anthriscus sylvestris</i>

PARK GRASS PLOTS
BOTANICAL COMPOSITION PER CENT.
1932 (1st Crop)

Plot	Manuring.	Liming.	Gramineæ.	Legumi-nosae.	Other Orders.	"Other Orders" consist largely of
3	Unmanured	Limed Unlimed	73.1 65.7	7.0 6.8	19.9 27.5	(None dominant) (None dominant)
7	Complete Mineral Manure	Limed Unlimed	68.7 71.4	14.4 15.6	16.9 13.0	(None dominant) (None dominant)
9	Complete Mineral Manure and double Amm. Salts	Limed Unlimed	97.7 100.0	— —	2.3 —	<i>Rumex acetosa</i> —
14	Complete Mineral Manure and double Nitrate of Soda	Limed (sun) Limed (shade) Unlimed	97.0 95.2 95.1	0.9 4.0 1.1	2.1 0.8 3.8	<i>Taraxacum vulgare</i> (None dominant) <i>Anthriscus sylvestris</i>
15	As plot 7 following double Nitrate of Soda, 1858-75	Limed				
17	Single Nitrate of Soda	Unlimed				
18	Mineral Manure (without Super) and double Sulphate Amm. 1905 and since.	L. 6,788 lb. L. 3,951 lb. Unlimed	92.8 83.5 96.4	— — —	7.2 16.5 3.6	(None dominant) <i>Taraxacum vulgare</i> <i>Rumex acetosa</i>
19	Farmyard Dung in 1905 and every fourth year since (omitted 1917)	L. 3,150 lb. L. 570 lb. Unlimed	91.2 88.2 87.8	1.8 1.6 1.2	7.0 10.2 11.0	<i>Ranunculus spp.</i> (None dominant) (None dominant)
20	Farmyard Dung in 1905 and every fourth year since (omitted in 1917); each intervening year Sulphate of potash, Super. and Nitrate of Soda	L. 2,772 lb. L. 570 lb. Unlimed	81.6 90.8 89.0	6.8 2.1 1.9	11.6 7.1 9.1	(None dominant) <i>Achillea millefolium</i> (None dominant)

PARK GRASS PLOT
BOTANICAL COMPOSITION PER CENT.
1933 (1st Crop)

Plot	Manuring.	Liming.	Gramineæ.	Legumi-nosæ.	Other orders.	“ Other Orders ” consist largely of
3	Unmanured	Limed	56.2	13.5	30.3	<i>Poterium sanguisorba</i> (None dominant)
7	Complete Mineral Manure	Unlimed	57.8	2.7	39.5	<i>Heracleum sphondylium</i>
		Limed	44.0	43.8	12.2	<i>Rumex acetosa</i>
		Unlimed	51.9	29.9	18.2	<i>Plantago lanceolata</i>
9	Complete Mineral Manure and double Amm. Salts	Limed	96.8	0.4	2.8	<i>Rumex acetosa</i>
14	Complete Mineral Manure and double Nitrate of Soda	Unlimed	100.0	—	—	<i>Taraxacum vulgare</i>
		Limed (sun)	95.9	1.3	2.8	<i>Anthriscus sylvestris</i>
		Limed (shade)	94.2	4.4	1.4	<i>Anthriscus sylvestris</i>
		Unlimed	98.6	0.2	1.2	<i>Plantago lanceolata</i>
15	As plot 7 following double Nitrate of Soda, 1858-75	Limed	58.1	26.4	15.5	<i>Plantago lanceolata</i>
		Unlimed	78.8	9.1	12.1	<i>Plantago lanceolata</i>
17	Single Nitrate of Soda	Limed	76.1	1.8	22.1	<i>Rumex acetosa</i>
		Unlimed	71.5	0.1	28.4	<i>Plantago lanceolata</i>
						<i>Centaurea nigra</i>
18	Mineral Manure (without Super) and double Sulphate Amm. 1905 and since.	L. 6,788 lb.	88.7	0.2	11.1	<i>Taraxacum vulgare</i>
		L. 3,951 lb.	84.8	—	15.2	<i>Taraxacum vulgare</i>
		Unlimed	97.9	—	2.1	<i>Rumex acetosa</i>
19	Farmyard Dung in 1905 and every fourth year since (omitted 1917)	L. 3,150 lb.	91.8	2.8	5.4	<i>Rumex acetosa</i>
20	Farmyard Dung in 1905 and every fourth year since (omitted in 1917); each intervening year Sulphate of potash, Super., and Nitrate of Soda.	L. 570 lb.	87.1	1.5	11.4	<i>Rumex acetosa</i>
		Unlimed	87.5	1.8	10.7	<i>Rumex acetosa</i>
		L. 2,772 lb.	74.8	10.0	15.2	<i>Taraxacum vulgare</i>
		L. 570 lb.	86.0	3.5	10.5	<i>Achillea millefolium</i>
		Unlimed	89.2	2.1	8.7	<i>Rumex acetosa</i>

PLOT 3—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

	Unlimed (U) ; limed (L).		1919		1921		1922		1923		1924		1925		1926		
	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	
<i>Gramineae</i>																	
1. Agrostis vulgaris	8.42	1.54	25.09	1.73	24.41	4.06	21.10	2.81	19.37	2.99	18.78	1.87	18.31	2.19	
2. Aira caespitosa	0.29	0.61	0.71	4.92	3.27	2.94	0.22	6.89	4.63	9.14	
3. Alopecurus pratensis	7.04	3.09	4.09	0.62	0.93	0.21	4.34	0.98	3.72	0.37	7.22	0.49	
4. Anthoxanthum odoratum	0.29	0.51	0.28	0.18	0.50	0.77	0.07	0.14	0.02	0.10	0.17	0.40	
5. Arrhenatherum avenaceum	0.89	2.78	1.27	3.14	2.41	4.33	0.67	2.11	0.53	2.42	0.42	3.39	
6. Avena flavaescens	4.16	19.27	2.68	18.07	2.70	11.53	3.89	17.86	3.62	16.98	5.82	32.10	
7. " pubescens	1.99	8.97	1.41	7.99	1.21	5.03	4.71	6.61	0.48	2.26	1.87	10.39	
8. Briza media	
9. Bromus mollis	
10. Cynosurus cristatus	0.20	0.21	0.06	0.14	
11. Dactylis glomerata	8.42	7.42	12.26	8.30	7.67	7.75	4.56	2.95	4.26	4.31	6.76	6.35	
12. Festuca ovina	6.73	5.06	13.38	13.27	13.20	10.55	20.35	18.00	10.86	8.72	10.63	7.99	
14. Holcus lanatus	9.22	7.73	11.56	8.78	3.41	1.75	3.59	2.81	2.29	1.74	7.82	4.97	6.64
1b. Lolium perenne	
16. Poa pratensis	0.15	1.96	0.14	1.29	0.42	2.10	0.22	2.25	0.16	2.16	0.09	3.06	
17. , trivialis	0.10	..	0.06	0.14	..	0.02	..	0.09	
<i>Leguminosae</i>																	
1. Lathyrus pratensis	0.89	1.24	0.78	1.66	1.00	3.15	0.82	2.47	2.45	6.41	1.95	1.91	
2. Lotus corniculatus	1.59	2.47	3.52	7.19	3.27	3.28	3.07	4.78	4.26	1.74	2.09	3.03	
4. Trifolium pratense	2.08	1.76	0.35	1.54	0.35	1.19	0.60	2.88	1.27	3.15	1.23	1.42	
5. " repens	

PLOT 3—(continued)

	Unlimed (U); limed (L)	1919		1921		1922		1923		1924		1925		1926	
		U	L	U	L	U	L	U	L	U	L	U	L	U	L
<i>Other orders</i>															
1. Ranunculus spp.	..	0.39	2.57	—	0.37	0.14	0.36	—	1.20	—	1.63	0.04	2.44	0.13	1.65
4. Cerastium vulgatum	..	0.60	0.83	0.07	0.06	0.21	0.28	0.45	1.13	0.32	0.16	0.17	0.22	0.07	0.49
5. Stellaria graminea	0.10	0.07	—	0.14	—	0.23	0.14	0.32	0.06	0.09	0.05	0.07	0.05
6. Linum catharticum	0.06	—	—	—	0.07	—	0.02	—	0.01	—	—
7. Agrimonia eupatoria	—	—	—	—	—	—	—	—	—	0.14
10. Potentilla reptans	—	—	—	0.07	—	—	—	—	—	—
11. Poterium sanguisorba	5.76	3.09	3.73	1.90	5.11	3.56	3.96	3.51	5.33	4.36	3.78	1.64	5.80
14. Conopodium denudatum	4.65	0.61	0.35	0.06	1.92	0.47	2.02	0.21	1.92	0.21	3.53	0.36	5.48
16. Pimpinella saxifraga	0.10	0.14	0.24	0.42	0.36	0.23	0.77	0.58	0.90	0.51	0.53
17. Galium verum	—	—	0.50	—	0.07	—	0.27	—	0.26	—
18. Scabiosa arvensis	1.65	0.28	1.17	0.35	1.96	0.23	1.34	0.48	6.78	0.47	2.13
19. Achillea millefolium	1.59	1.86	0.92	1.17	1.35	1.46	1.35	0.98	2.40	2.42	1.28	0.89	1.55
20. Centaurea nigra	5.76	6.19	4.44	4.79	6.81	21.31	2.17	3.65	9.80	9.09	3.32	2.75	2.84
22. Hieracium pilosella	—	—	—	—	0.07	—	—	0.04	0.04	—
24. Leontodon hispidus	6.94	1.54	2.32	0.56	4.26	1.53	2.62	1.76	5.96	1.05	2.55	0.44	5.48
26. Taraxacum vulgare	—	—	0.14	—	0.30	0.05	0.10	0.16	0.09	0.19
27. Tragopogon pratensis	0.07	0.06	—	0.28	0.07	0.29	0.37	0.42	0.43	0.49
28. Primula veris	—	—	—	—	—	—	—	—	0.07	—
29. Plantago lanceolata	19.12	12.06	7.68	9.22	11.07	8.32	11.52	7.31	10.70	4.83	14.91	7.06	17.41
30. Veronica chamaedrys	0.05	0.10	0.14	0.06	0.07	0.07	0.22	0.29	0.21	0.11	0.21	0.18	0.14
31. Ajuga reptans	0.10	—	—	—	—	0.07	0.01	0.03	—	0.13	—
33. Thymus serpyllum	—	—	—	—	—	0.01	—	0.04	—	—
34. Rumex acetosa	1.98	4.23	0.85	0.93	1.49	0.84	1.20	0.43	0.85	0.79	0.76	1.02	2.39
35. Luzula campestris	0.15	0.20	0.43	0.12	0.50	0.14	3.14	1.41	0.37	0.58	0.17	0.13	0.07
36. Carex praecox	0.55	0.36	0.71	0.37	0.42	0.35	1.87	0.84	1.70	0.58	0.85	0.31	0.97

PLOTS 5¹ AND 5²—BOTANICAL COMPOSITION PER CENT (INDIVIDUAL SPECIES)

No liming on these plots.	PLOT 5 ¹				PLOT 5 ²			
	1919	1921	1922	1924	1926	1919	1921	1922
<i>Gramineae</i>								
1. Agrostis vulgaris ..	4.49	12.73	22.31	12.10	23.62	8.42	20.30	15.19
2. Aira cespitosa ..	0.26	—	—	1.49	0.49	—	—	0.07
3. Alopecurus pratensis ..	0.74	1.06	4.65	0.48	0.58	11.15	5.67	20.39
4. Anthoxanthum odoratum ..	11.69	8.77	0.52	4.15	12.30	5.12	3.86	0.44
5. Arrhenatherum avenaceum ..	1.67	7.78	2.38	1.53	4.94	2.28	3.41	4.02
6. Avena flavaescens ..	—	—	0.22	0.02	—	—	—	0.07
7. " pubescens ..	0.21	0.21	0.28	0.28	0.27	1.71	2.04	0.67
9. Bromus mollis ..	—	—	0.06	—	—	0.06	—	—
11. Dactylis glomerata ..	8.87	14.57	9.29	6.37	8.17	7.40	2.04	7.15
12. Festuca ovina ..	46.73	40.74	15.51	39.91	27.88	19.57	27.79	16.97
14. Holcus lanatus ..	0.57	0.28	0.28	0.20	1.21	1.13	2.38	0.97
16. Poa pratensis ..	1.35	0.64	1.75	1.49	1.17	6.26	5.00	5.81
17. " trivialis ..	—	—	—	—	—	0.11	0.06	—
<i>Leguminosae</i>								
1. Lathyrus pratensis ..	—	—	0.06	0.56	0.04	1.26	6.12	1.56
2. Lotus corniculatus ..	0.36	1.13	2.10	1.01	0.45	3.07	2.61	4.09
4. Trifolium pratense ..	—	—	—	0.10	0.09	—	0.11	0.15
5. " repens ..	—	—	—	0.01	—	—	—	0.07
6. Vicia sepium ..	—	—	—	—	—	0.23	0.34	0.30
						—	—	1.70
								1.53

PLOTS 5¹ AND 5²—(continued).

No liming on these plots.	PLOT 5 ¹				PLOT 5 ²			
	1919	1921	1922	1924	1926	1919	1921	1922
<i>Other orders</i>								
1. & 2. <i>Ranunculus</i> spp.	0.31	0.07	0.22	0.28	0.18	0.62	0.46	0.22
4. <i>Cerastium vulgatum</i>	0.42	0.07	0.17	0.13	0.09	0.17	0.11	0.07
5. <i>Stellaria media</i>	0.10	0.07	0.11	0.60	0.18	0.34	0.06	0.22
11. <i>Potentilla sangisorba</i>	—	—	0.06	—	—	—	—	0.34
14. <i>Conopodium denudatum</i>	1.31	0.14	0.22	1.98	1.79	0.11	0.11	—
15. <i>Heracleum sphondylium</i>	—	—	0.17	—	—	0.28	—	—
16. <i>Pimpinella saxifraga</i>	0.05	—	0.11	0.13	0.04	0.40	0.06	0.22
17. <i>Galium verum</i>	1.98	0.64	0.39	1.30	0.31	0.23	—	0.22
18. <i>Scabiosa arvensis</i>	0.62	1.13	5.72	4.19	1.44	0.23	1.36	1.12
19. <i>Achillea millefolium</i>	0.73	0.35	0.46	1.34	1.03	2.84	1.36	1.19
20. <i>Centaurea nigra</i>	3.86	6.65	19.37	14.92	6.29	3.41	3.18	7.22
22. <i>Hieracium pilosella</i>	—	—	—	0.01	0.04	—	—	1.04
23. <i>Hypochaeris radicata</i>	—	—	—	—	—	0.11	—	—
24. <i>Leontodon hispidus</i>	0.10	—	0.22	0.16	0.09	—	—	0.11
27. <i>Tragopogon pratensis</i>	—	—	0.17	—	0.09	0.45	—	—
29. <i>Plantago lanceolata</i>	0.42	0.28	0.91	1.01	1.70	0.57	0.23	0.30
30. <i>Veronica chamaedrys</i>	0.06	0.14	—	0.41	0.13	0.06	—	0.22
34. <i>Rumex acetosa</i>	12.31	2.20	11.21	3.15	4.76	18.09	7.82	9.52
35. <i>Luzula campestris</i>	0.79	0.35	1.08	0.69	0.63	4.32	3.29	0.52

PLOT 7—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

Unlimed (U) or Limed (L) :	1919		1921		1922		1923		1924		1925	
	U	L	U	L	U	L	U	L	U	L	U	L
<i>Gramineae</i>												
1. Agrostis vulgaris ..	5.41	2.02	12.07	5.31	19.30	5.25	15.53	4.85	13.87	1.23	11.05	1.79
3. Alopecurus pratensis ..	1.74	15.23	1.41	12.09	1.36	10.98	1.64	6.67	1.97	9.95	1.07	8.90
4. Anthoxanthum odoratum ..	3.57	0.45	4.70	0.67	1.56	0.09	4.40	0.76	1.10	0.16	2.26	0.09
5. Arrhenatherum avenaceum ..	3.07	2.63	1.18	5.42	1.36	4.15	1.00	3.44	1.71	3.70	5.25	27.32
6. Avena flaveolens ..	0.72	0.96	1.64	1.30	2.36	2.23	0.71	2.32	1.10	1.75	0.56	0.45
7. " pubescens ..	2.86	8.83	2.70	12.09	1.66	4.92	2.85	6.37	1.32	4.77	1.49	8.10
8. Briza media ..	—	—	—	—	0.05	—	—	—	—	—	0.09	—
9. Bromus mollis ..	0.05	0.60	0.12	2.60	0.15	5.42	0.08	7.94	—	—	—	0.04
11. Dactylis glomerata ..	21.64	18.61	14.08	11.87	11.64	10.21	9.80	5.41	11.90	9.95	34.13	22.81
12. Festuca ovina ..	7.15	5.40	18.20	9.83	24.19	13.17	28.33	10.41	10.84	2.46	9.26	3.18
14. Holcus lanatus ..	4.28	1.56	11.73	2.26	3.78	1.87	2.30	1.22	1.35	0.48	6.49	0.67
15. Lolium perenne ..	—	—	—	—	—	—	—	—	0.04	—	—	0.04
16. Poa pratensis ..	1.18	1.61	0.93	1.98	2.11	2.78	2.43	2.94	1.85	2.03	1.02	1.12
17. " trivialis ..	0.36	1.21	0.12	0.62	0.15	0.27	0.08	0.45	0.04	0.40	0.04	0.67
<i>Leguminosae</i>												
1. Lathyrus pratensis ..	7.15	19.57	7.75	21.26	11.64	29.21	8.75	36.62	28.62	51.06	6.10	1.07
2. Lotus corniculatus ..	0.46	0.03	1.53	0.11	0.81	—	0.71	0.10	1.10	0.56	0.04	0.04
4. Trifolium pratense ..	1.03	0.03	3.29	0.11	0.25	0.32	5.57	0.70	3.06	0.12	0.21	0.04
5. " repens ..	0.05	—	0.70	0.56	0.05	0.09	0.17	2.73	0.41	0.04	—	0.04
7. Vicia sepium	0.10	—	0.23	—	—	—	0.07	—	—	—

PLOT 7 (continued)

Unlimed (U); limed (L)	1919			1921			1922			1923			1924			1925		
	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L
<i>Other species</i>																		
1. Ranunculus spp.	1.94	3.99	0.47	1.30	0.15	0.27	1.00	1.77	0.59	0.83	0.81	2.15	—	—	—	—	—	
4. Cerastium vulgatum	0.05	0.10	2.59	—	—	—	—	0.05	—	—	—	—	—	—	—	—	—	
5. Stellaria media	0.15	—	—	3.05	—	1.11	—	0.08	—	0.04	—	0.04	—	—	—	—	—	
12. Spiraea ulmaria	1.89	—	—	—	—	—	—	2.26	—	—	1.21	—	—	1.24	—	—	—	
13. Anthriscus sylvestris	—	—	—	—	—	—	—	0.27	—	0.25	—	0.72	—	—	—	—	1.07	
14. Conopodium denudatum	9.49	3.68	1.53	1.30	0.61	0.83	1.68	0.87	3.21	0.60	2.56	0.45	—	—	—	—	—	
15. Heracleum sphondylium	4.18	1.56	1.53	4.86	2.62	2.28	1.68	0.87	3.25	4.93	4.22	11.99	—	—	—	—	—	
16. Pimpinella saxifraga	—	—	0.03	—	—	—	0.09	—	—	—	—	—	—	0.04	—	—	—	
17. Galium verum	0.20	—	—	—	—	—	—	0.04	—	—	—	—	—	—	—	—	—	
18. Scabiosa arvensis	0.10	0.55	0.35	0.73	—	—	0.45	0.50	0.50	0.33	0.28	0.34	—	—	—	—	—	
19. Achillea millefolium	6.34	0.96	3.69	0.45	2.52	0.45	0.67	0.20	3.14	0.83	1.83	0.63	—	—	—	—	—	
20. Centaurea nigra	2.55	1.11	2.59	1.08	6.40	1.28	0.96	0.91	0.55	1.19	2.22	0.81	—	—	—	—	—	
23. Hypochaeris radicata	—	—	—	—	—	—	—	—	—	0.14	—	—	—	—	—	—	—	
26. Taraxacum vulgare	—	—	0.05	—	—	—	0.65	—	0.20	—	0.36	—	—	—	—	—	1.43	
27. Tragopogon pratensis	0.05	1.46	0.23	1.19	0.25	1.64	0.04	0.45	—	0.32	0.04	0.09	—	—	—	—	—	
29. Plantago lanceolata	1.33	0.50	1.18	0.67	1.96	0.74	3.56	0.45	4.20	0.48	3.33	0.18	—	—	—	—	—	
30. Veronica chamaedrys	—	—	0.10	0.12	0.06	—	0.04	—	0.07	0.08	0.04	0.04	—	—	—	—	—	
34. Rumex acetosa	10.61	7.12	—	0.28	1.81	0.09	1.76	0.45	2.63	0.64	4.01	4.83	—	—	—	—	—	
35. Luzula campestris	—	—	0.30	0.05	0.29	—	0.15	—	1.38	0.10	0.29	0.04	0.22	—	—	—	—	

PLOT 9—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

Unlimed (U); limed (L).	1919			1921			1922			1923			1924			1925			1926		
	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	
<i>Gramineae</i>																					
1. Agrostis vulgaris	12.46	2.27	27.37	4.15	16.48	3.58	23.30	2.39	31.04	3.33	17.08	1.77	24.84	3.04		
3. Alopecurus pratensis	0.68	26.02	0.58	22.56	13.81	27.83	0.32	28.34	0.53	45.66	1.55	42.21	0.13	23.62		
4. Anthoxanthum odoratum	5.44	1.12	25.15	2.08	8.40	0.36	42.81	1.45	22.41	0.40	13.16	0.73	15.84	1.34		
5. Arrhenatherum avenaceum	46.88	47.35	4.44	43.06	11.06	30.57	8.16	35.60	21.71	32.48	20.21	44.84	6.08	49.64		
6. Avena flavescens	0.02	0.06	0.26	0.21	0.10	—	—	—	—	—	—	—	—	—	—		
7. " pubescens	—	—	—	0.05	0.05	—	—	—	—	—	—	—	—	—	—		
9. Bromus mollis	3.30	6.84	1.12	4.92	3.62	8.04	0.70	2.15	0.09	1.71	0.53	4.30	0.36	8.39	..		
11. Dactylis glomerata	3.59	5.92	9.94	7.78	11.17	4.61	12.36	13.25	8.40	3.21	3.59	2.19	1.53	3.13	..		
12. Festuca ovina	12.43	0.76	30.05	2.08	32.53	1.29	11.91	0.23	14.21	0.20	40.45	0.31	50.73	3.19	..		
14. Holcus lanatus	0.21	5.56	0.29	9.34	1.60	22.44	0.24	15.92	0.33	11.97	0.08	3.11	0.10	6.32	..		
16. Poa pratensis	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
17. " trivialis	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
<i>Leguminosae</i>																					
1. Lathyrus pratensis	0.10	—	—	0.11	0.26	—	—	0.24	0.08	—	0.04	—	0.09		
<i>Other orders</i>																					
10. Potentilla reptans	0.09	—	—	—	—	—	—	—	—	0.38	—	0.29	—	0.03	—		
14. Conopodium denudatum	0.05	0.11	—	0.26	—	—	0.31	0.04	0.14	—	0.36	—	0.27	—	0.12		
15. Heracleum sphondylium	0.27	0.12	1.30	0.53	0.10	—	0.14	0.52	—	2.00	—	—	0.36	—		
20. Centaurea nigra	0.14	—	—	—	—	—	—	—	—	—	—	—	—	—	—	..		
26. Taraxacum vulgare	—	—	0.26	—	—	—	—	—	—	0.28	—	—	—	—	0.06		
34. Rumex acetosa	14.87	3.52	0.82	1.95	0.32	0.10	0.16	0.34	0.14	0.32	1.06	0.23	—	—	—	—	0.97		
35. Luzula campestris	—	—	0.06	—	—	—	—	—	—	—	—	—	—	—	—	—		

PLOT 9 (continued)

Unlimed (U); limed (L)	1927			1928			1929			1930			1931			1932			1933		
	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	
<i>Gramineae</i>																					
1. <i>Agrostis vulgaris</i> ..	14.14	2.95	7.57	2.91	4.57	—	3.26	12.12	2.43	—	5.33	—	2.95	—	—	—	—	—	—		
3. <i>Alopecurus pratensis</i> ..	0.25	38.39	0.41	54.39	40.35	—	57.42	0.16	46.79	—	35.13	—	25.56	—	—	—	—	—	—		
4. <i>Anthoxanthum odoratum</i> ..	10.82	1.54	10.32	3.33	1.65	—	0.75	0.33	1.33	—	3.67	—	3.98	—	—	—	—	—	—		
5. <i>Arrhenatherum avenaceum</i> ..	2.98	38.90	4.16	17.88	18.20	—	20.91	2.32	33.97	—	39.34	—	50.91	—	—	—	—	—	—		
6. <i>Avena flavescens</i> ..	—	0.04	—	—	0.04	—	0.07	—	—	—	0.05	—	—	—	—	—	—	—	—		
7. " <i>pubescens</i> ..	—	0.03	0.17	—	—	—	0.03	—	—	—	0.05	—	0.31	—	—	—	—	—	—		
9. <i>Bromus mollis</i> ..	—	—	6.02	0.45	1.70	—	—	0.03	—	0.33	—	0.59	—	1.29	—	—	—	—	—		
11. <i>Dactylis glomerata</i> ..	0.55	—	3.15	1.38	8.36	—	2.32	1.10	4.38	—	4.06	—	2.48	—	—	—	—	—	—		
12. <i>Festuca ovina</i> ..	1.14	—	3.15	1.38	8.36	—	7.08	7.40	2.10	—	1.42	—	1.45	—	—	—	—	—	—		
14. <i>Holcus lanatus</i> ..	69.42	1.70	75.57	0.54	2.70	100.00	0.46	76.08	1.43	100.00	4.01	100.00	4.91	—	—	—	—	—	—		
16. <i>Poa pratensis</i> ..	—	6.27	0.07	9.44	10.18	—	7.05	—	—	6.29	—	3.96	—	2.95	—	—	—	—	—		
17. " <i>trivialis</i> ..	—	—	—	—	—	—	—	—	—	—	—	0.05	—	—	—	—	—	—	—		
<i>Leguminosae</i>																					
1. <i>Lathyrus pratensis</i> ..	0.03	0.25	—	—	—	0.04	—	0.07	—	—	—	—	—	—	—	—	—	—	0.36		
<i>Other orders</i>																					
13. <i>Anthriscus sylvestris</i> ..	0.03	0.04	—	—	0.12	—	0.26	—	0.03	—	—	—	—	—	—	—	—	—	—		
14. <i>Conopodium denudatum</i> ..	—	—	0.52	0.04	—	0.37	—	0.39	—	0.39	—	0.52	—	—	—	—	—	—	—		
15. <i>Heracleum sphondylium</i> ..	0.03	—	—	—	—	0.54	—	—	—	0.04	—	—	—	—	—	—	—	—	—		
19. <i>Achillea millefolium</i> ..	—	—	—	—	—	0.17	—	—	—	—	—	—	—	—	—	—	—	—	—		
20. <i>Centaurea nigra</i> ..	—	—	0.03	0.04	—	0.54	—	—	—	—	—	—	—	—	—	—	—	—	—		
26. <i>Taraxacum vulgare</i> ..	—	—	0.03	0.21	0.07	0.54	—	—	—	—	—	—	—	—	—	—	—	—	—		
29. <i>Plantago lanceolata</i> ..	—	—	0.03	—	—	—	0.15	—	—	0.13	—	0.19	—	—	—	—	—	—	—		
34. <i>Rumex acetosa</i> ..	—	—	—	—	—	—	—	—	—	—	0.45	—	—	—	—	—	—	—	—		
38. <i>Epilobium angustifolium</i> ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		

Herbage killed by frost and drought.

No sample taken.

150

PLOT 14—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

Unlimed (U); limed (L).	1919			1920			1921			1922			1923			1924			1925			1926			
	U	U	L (sun)	U	L (shade)	U	U	L	U	U	L	U	U	L	U	U	L (sun)	U	U	L (sun)	U	L (shade)	L (sun)		
<i>Gramineae</i>																									
1. Agrostis vulgaris ..	0.55	0.05	0.24	2.61	—	0.96	0.17	0.80	0.05	0.61	0.27	0.27	0.26	—	—	0.60	0.23	—	—	—	—	—	—	—	
3. Alopecurus pratensis ..	53.65	47.83	52.87	47.47	34.72	33.34	58.33	42.54	42.28	23.90	43.70	28.65	35.82	18.88	22.05	14.16	—	—	—	—	—	—	—	—	
4. Anthoxanthum odoratum ..	0.37	—	1.25	—	0.60	—	—	—	0.09	—	0.08	0.01	0.01	0.04	0.01	0.04	0.60	0.10	—	—	—	—	—	—	
5. Arrhenatherum avenaceum ..	23.38	37.34	29.99	3.36	48.13	40.89	24.67	24.70	33.17	34.81	39.36	40.04	40.44	53.96	29.63	57.71	—	—	—	—	—	—	—	—	
6. Avena fatescens ..	0.73	0.05	0.30	2.48	0.30	1.11	0.50	0.25	—	0.89	0.04	0.27	—	—	—	—	0.06	0.35	0.20	—	—	—	—	—	
7. " pubescens ..	3.67	0.16	1.07	14.42	1.64	3.71	0.73	1.62	0.83	5.04	0.39	2.79	0.26	—	—	0.12	15.97	1.31	—	—	—	—	—	—	—
9. Bromus mollis ..	0.51	0.52	0.24	0.75	1.05	1.55	4.60	6.55	3.70	15.29	—	—	—	—	—	—	0.03	0.05	—	—	—	—	—	—	
11. Dactylis glomerata ..	3.23	4.61	3.02	3.48	5.14	2.52	3.14	2.42	4.96	0.56	3.46	1.42	6.83	3.05	2.66	6.23	—	—	—	—	—	—	—	—	
12. Festuca ovina ..	5.06	0.05	1.78	11.56	0.15	5.33	0.17	5.34	0.22	9.17	0.17	7.15	0.02	0.12	16.88	3.11	—	—	—	—	—	—	—	—	
14. Holcus lanatus ..	—	—	—	0.74	—	—	—	—	—	—	—	0.04	—	—	—	—	—	—	—	—	—	—	—	—	
15. Lolium perenne ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
16. Poa pratensis ..	0.81	0.52	1.90	4.73	0.82	2.30	1.18	5.45	0.84	2.59	0.81	2.16	0.21	0.73	1.41	1.18	—	—	—	—	—	—	—	—	
17. " trivialis ..	1.02	6.71	3.74	0.31	5.44	3.18	2.91	2.82	7.61	2.78	1.98	2.12	2.47	6.58	0.05	10.72	—	—	—	—	—	—	—	—	
<i>Leguminosae</i>																									
1. Lathyrus pratensis ..	2.38	—	0.89	5.84	0.15	2.96	0.17	1.06	0.11	0.89	0.46	6.21	0.17	0.02	5.73	0.62	—	—	—	—	—	—	—	—	
<i>Other orders</i>																									
1 & 2. Ranunculus spp.	—	—	1.00	0.19	—	0.07	0.96	0.40	—	1.82	—	0.04	—	—	—	—	—	—	—	—	—	—	—	—	
13. Anthriscus sylvestris ..	2.41	—	—	—	—	—	—	—	—	—	—	0.05	—	—	—	—	—	—	—	—	—	—	—	—	
14. Conopodium denudatum ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
18. Scabiosa arvensis ..	0.04	—	—	—	—	0.12	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
19. Achillea millefolium ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
20. Centaurea nigra ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
26. Taraxacum vulgare ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
29. Plantago lanceolata ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
34. Rumex acetosa ..	1.02	0.79	1.24	0.25	0.82	0.22	0.28	0.15	1.77	0.80	0.66	0.27	1.69	1.10	0.05	1.15	—	—	—	—	—	—	—	—	

Part of the limed area of Plot 14 is shaded by a large oak tree. Samples of the shaded and unshaded (sun) areas were taken separately in 1920, and every year from 1925 onwards.

	Plot 15		Unlimed (U); limed (L)		1919 U		1921 U		1923 U		1925 U		1927 U		1929 U		1931 U		1933 U	
	Gramineae	Leguminosae	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L
1. Agrostis vulgaris	11.39	15.20	9.44	—	13.12	13.83	17.00	7.71	12.48	6.31	11.56	2.51	20.51	2.22	16.80	3.58	—	
2. Aira caespitosa	8.55	19.95	8.53	15.48	10.15	—	8.80	13.94	9.20	—	—	—	—	—	8.92	
3. Alopecurus pratensis	5.04	2.49	3.78	0.16	2.65	1.11	3.72	0.92	1.10	0.16	4.88	0.64	8.24	0.82	—	
4. Anthoxanthum odoratum	3.52	—	3.51	1.12	2.60	1.12	1.07	9.30	5.46	0.68	0.33	5.82	5.92	10.84	7.97	
5. Arrhenatherum avenaceum	1.46	2.63	3.51	0.76	0.95	0.61	0.76	2.62	0.89	1.74	0.84	3.51	0.78	1.95	1.95	
6. Avena flavescens	0.86	1.60	1.62	0.74	0.95	0.61	0.76	2.62	0.89	1.74	0.84	3.51	0.78	1.95	1.95	
7. " pubescens	1.97	3.29	3.24	2.00	3.30	1.77	6.33	2.77	10.91	2.99	8.23	3.24	12.57	4.20	12.55	
8. Briza media	0.09	—	—	—	—	—	0.02	—	—	—	—	—	—	—	—	
9. Bromus mollis	4.97	5.26	2.16	3.26	1.59	9.64	6.46	16.41	5.17	5.78	2.78	6.71	6.84	12.34	6.59	
11. Dactylis glomerata	7.36	22.27	20.35	23.72	8.53	9.85	7.89	7.90	17.08	7.90	10.93	12.66	9.18	13.37	13.37	
12. Festuca ovina	5.99	11.02	5.39	2.08	1.65	9.78	3.70	12.19	3.46	2.47	0.87	7.18	4.25	3.99	1.57	
14. Holcus lanatus	—	—	—	—	0.89	—	—	—	0.37	—	—	—	—	0.16	—	
15. Lolium perenne	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
16. Poa pratensis	1.37	1.10	1.69	1.41	1.78	0.84	1.02	0.66	1.11	0.68	0.65	1.50	2.50	1.09	0.31	
17. " trivialis	0.21	0.07	0.13	0.15	0.89	0.09	0.22	0.04	0.07	—	—	0.09	1.20	0.05	0.25	
1. Lathyrus pratensis	5.31	8.32	18.18	15.06	11.22	4.01	14.31	4.84	26.18	16.03	27.24	4.74	3.79	7.67	13.81	
2. Lotus corniculatus	—	0.15	—	—	—	—	—	—	0.04	—	0.16	0.09	—	—	—	
4. Trifolium pratense	0.09	0.15	0.07	0.07	2.60	0.56	0.45	0.62	1.07	3.00	1.31	0.05	0.46	0.52	5.96	
5. " repens	—	0.22	—	0.29	4.57	0.09	0.05	0.45	1.44	2.94	21.22	2.82	1.57	0.88	6.59	
1 & 2. Other orders	2.66	0.51	0.88	1.34	1.40	2.65	3.97	0.83	0.81	0.74	0.55	0.05	0.19	0.36	0.50	
1. Ranunculus spp.	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
4. Cerastium vulgatum	—	0.15	—	—	0.15	—	—	—	—	—	0.05	—	0.05	—	—	
5. Stellaria media	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
11. Poterium sanguisorba	—	—	—	—	0.07	—	—	—	—	—	—	—	—	—	—	
13. Anthriscus sylvestris	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
14. Conopodium denudatum	3.60	0.73	2.02	2.52	1.97	—	2.93	1.20	2.19	0.58	0.11	0.38	0.37	1.50	0.75	
15. Heracleum sphondylium	0.30	—	—	—	—	—	0.23	0.40	—	0.04	0.26	0.65	—	1.29	—	
16. Pimpinella saxifraga	—	—	—	—	—	—	—	—	0.08	0.04	0.16	—	0.05	—	—	
17. Galium verum	—	—	—	—	—	—	—	—	0.33	0.16	0.05	0.09	—	—	—	
19. Achillea millefolium	5.31	2.63	1.75	1.71	1.20	7.08	2.36	6.20	3.13	15.66	3.76	7.74	3.33	1.45	0.88	
20. Centaurea nigra	0.60	0.58	0.07	0.28	0.76	0.70	0.18	0.54	0.26	0.89	—	0.14	1.11	0.42	0.82	
24. Leontodon hispidus	1.29	0.66	1.15	0.82	1.59	—	—	—	0.04	—	—	—	—	—	—	
26. Taraxacum vulgare	0.26	0.29	0.07	0.51	0.13	0.14	0.04	0.41	0.33	0.53	—	0.09	—	0.10	—	
27. Tragopogon pratensis	3.76	7.23	4.51	13.49	4.63	15.19	8.38	6.16	5.10	5.57	4.14	1.92	5.45	4.40	9.67	
29. Plantago lanceolata	—	—	—	—	0.07	0.06	0.05	0.04	0.11	—	—	—	0.18	—	0.13	
30. Veronica chamaedrys	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
31. Ajuga reptans	7.36	1.39	0.60	2.00	0.31	3.17	—	—	—	—	—	—	—	—	—	
34. Rumex acetosa	0.21	0.81	0.13	3.63	0.31	0.37	0.04	0.04	0.04	0.04	0.07	0.42	0.11	0.19	0.99	
35. Luzula campestris	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

151

PLOT 17—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

Unlimed (U); Limed (L).	1919		1921		1923		1925		1927		1929		1931		1933	
	U	L	U	L	U	L	U	L	U	L	U	L	U	L	U	L
<i>Gramineae</i>																
1. Agrostis vulgaris ..	6.06	5.63	4.77	9.22	4.02	3.31	1.95	5.52	3.56	3.13	0.97	4.52	*	5.06	0.49	
3. Alopecurus pratensis ..	12.86	12.19	10.22	13.32	10.35	14.39	13.20	14.04	13.70	17.76	11.62	17.92	12.34	14.31	9.34	
4. Anthoxanthum odoratum ..	7.35	8.67	3.07	7.49	2.88	7.11	0.88	10.53	2.96	2.73	0.23	5.53	0.73	10.31	0.55	
5. Arrhenatherum avenaceum ..	0.32	—	—	0.07	0.35	0.53	0.08	1.35	2.25	0.12	0.14	0.39	0.60	0.05	1.95	
6. Avena flavaescens ..	1.30	0.77	2.45	1.52	1.30	0.40	1.16	0.30	2.19	0.35	1.49	0.47	2.46	0.29	2.57	
7. " pubescens ..	4.86	4.10	6.95	3.53	10.42	2.34	15.56	2.56	17.04	2.32	18.13	3.20	16.32	1.67	10.50	
8. Briza media ..	0.54	0.29	0.75	0.21	0.35	0.09	0.52	0.20	0.38	0.12	0.70	0.15	0.80	0.33	0.98	
9. Bromus mollis ..	—	—	0.07	0.07	0.35	—	—	—	—	—	—	0.08	0.13	0.05	0.06	
10. Cynosurus cristatus ..	0.11	0.23	0.07	—	—	0.04	0.04	0.05	0.11	0.06	—	—	0.07	0.14	0.18	
11. Dactylis glomerata ..	8.33	5.16	10.63	7.27	4.32	28.34	15.01	23.87	7.12	18.63	7.62	23.54	10.48	17.65	10.75	
12. Festuca ovina ..	4.33	11.60	21.55	17.57	35.42	6.18	21.67	5.07	20.77	6.44	26.82	8.26	27.87	6.77	29.31	
14. Holcus lanatus ..	10.92	15.94	12.55	2.74	2.66	9.80	6.46	11.84	10.19	8.76	2.04	15.43	4.44	13.65	5.86	
15. Lolium perenne ..	0.75	0.23	0.81	0.65	0.36	0.80	0.36	0.15	0.60	0.29	0.70	0.62	1.06	0.48	1.71	
16. Poa pratensis ..	0.27	0.35	—	0.58	0.07	0.44	0.08	0.40	0.11	0.23	0.14	0.08	0.07	0.14	—	
17. " trivialis ..	0.54	0.17	0.20	1.44	1.07	0.97	1.12	0.15	1.48	—	0.19	0.23	1.13	0.57	1.89	
<i>Leguminosae</i>																
1. Lathyrus pratensis ..	0.05	0.17	—	0.07	0.07	0.09	—	0.05	—	0.12	—	0.23	—	0.05	—	
2. Lotus corniculatus ..	0.32	—	—	0.89	—	1.07	—	0.52	0.05	1.26	0.12	1.67	0.16	3.32	—	
4. Trifolium pratense ..	—	—	—	—	0.07	—	—	—	—	—	—	—	—	—	—	

PLOT 17 (continued)

Unlimed (U) ; Limed (L).	1919		1921		1923		1925		1927		1931		1933	
	U	L	U	L	U	L	U	L	U	L	U	L	U	L
<i>Other orders</i>														
1 & 2. <i>Ranunculus</i> spp.	0.81	0.23	0.27	0.14	0.28	0.36	0.51	0.36	0.62	0.18	0.67	0.20	0.38	0.42
4. <i>Cerastium vulgatum</i>	..	0.32	0.17	0.31	0.24
5. <i>Stellaria media</i>	..	0.05	0.19	0.49
11. <i>Poterium sanguisorba</i>
14. <i>Conopodium denudatum</i>	..	1.19	0.12	..	0.27	0.51	0.79	1.24	1.00	0.45	0.28	0.11	0.19	0.08
15. <i>Heracleum sphondylium</i>	..	0.27	..	0.20	..	0.07	0.20	0.09	..
16. <i>Pimpinella saxifraga</i>	0.05	..
17. <i>Galium verum</i>
18. <i>Scabiosa arvensis</i>
19. <i>Achillea millefolium</i>	..	0.92	0.12	0.41	0.28	1.07	0.22	0.88	0.30	1.04	0.29	0.14	0.78	0.79
20. <i>Centaurea nigra</i>	..	8.65	1.99	3.62	1.22	1.37	1.68	2.43	1.91	2.25	1.96	2.19	0.78	0.81
22. <i>Hieracium pilosella</i>	2.39	0.31
24. <i>Leontodon hispidus</i>	..	3.36	1.12	1.57	1.94	1.51	1.50	1.31	1.96	1.10	1.10	1.10
26. <i>Taraxacum vulgare</i>	..	0.16	0.17	..	0.36	0.36	0.22	1.24	0.15	0.22	0.40	0.70	0.55	0.26
27. <i>Tragopogon pratensis</i>	0.12	0.20	0.06
29. <i>Plantago lanceolata</i>	..	24.12	29.42	17.73	27.09	15.96	17.40	10.74	16.40	8.22	22.98	14.83	7.95	7.56
30. <i>Veronica chamaedrys</i>	0.23	..	0.28	0.07	0.13	0.04	0.15	0.05	0.12	0.09	0.20
31. <i>Ajuga reptans</i>	0.16	0.17	0.22	0.04	0.75	..	0.06	0.05	..
32. <i>Prunella vulgaris</i>	0.14
33. <i>Thymus serpyllum</i>	0.87	0.12	0.20	0.28	0.51	1.24	1.20	0.90	1.86	0.70	0.56	0.93
34. <i>Rumex acetosa</i>	0.05	0.35	0.07	0.79	0.51	0.22	0.16	0.20	0.16	0.17	0.47	0.33
35. <i>Luzula campestris</i>	0.16	0.17	0.34	0.79	0.93	0.13	0.12	0.20	0.06	0.29	0.14	0.18
36. <i>Carex praecox</i>

* The very small amount of *Agrostis* present was accidentally lost at the time of weighing.

PLOT 18—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

Unlimed (U); lightly limed (LL); heavily limed (HL).	1919			1920			1921			1922			1923		
	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL
<i>Gramineae</i>															
1. Agrostis vulgaris ..	17.59	42.94	—	0.05	—	—	50.87	41.16	42.54	35.90	31.50	47.92	28.29	27.16	—
2. Aira caespitosa ..	5.08	5.99	4.28	6.25	3.50	5.33	9.90	25.45	17.61	5.05	10.32	8.34	—	—	—
3. Alopecurus pratensis ..	3.46	5.85	7.01	5.34	8.07	4.06	4.43	2.68	0.74	1.00	19.82	2.20	2.58	—	—
4. Anthoxanthum odoratum ..	1.62	0.51	0.78	0.53	0.75	1.97	1.69	1.03	2.45	3.15	0.13	1.97	2.65	—	—
5. Arrhenatherum avenaceum ..	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
6. Avena flavescens ..	—	—	—	0.11	—	0.09	—	—	—	0.19	0.20	1.14	1.29	—	—
7. " pubescens ..	—	—	—	—	—	—	—	—	—	0.31	0.40	1.00	—	0.12	—
9. Bromus mollis ..	—	—	—	34.00	16.65	8.38	16.16	11.97	—	—	0.10	0.16	—	—	—
11. Dactylis glomerata ..	—	—	—	4.21	13.88	5.47	12.20	11.44	21.80	12.93	11.75	7.01	8.02	3.24	8.00
12. Festuca ovina ..	—	—	—	2.37	1.03	7.87	1.91	4.04	6.03	7.50	9.79	12.02	18.27	9.67	31.12
14. Holcus lanatus ..	—	—	—	0.43	0.81	2.05	0.61	0.61	3.30	5.25	0.93	1.87	1.07	0.19	1.34
16. Poa pratensis ..	—	—	—	—	—	—	—	—	2.32	2.68	0.72	2.61	4.16	2.20	6.95
17. " trivialis ..	—	—	—	—	—	—	—	—	—	—	—	0.21	—	—	—
1. Lathyrus pratensis ..	—	—	—	0.15	—	—	—	—	—	—	—	—	1.43	0.13	—
<i>Leguminosae</i>															
1 & 2. Ranunculus spp. ..	—	—	—	—	—	—	—	—	—	—	—	—	0.35	—	—
5. Stellaria media ..	4.32	0.51	0.78	0.53	0.40	0.35	0.69	1.76	—	2.37	2.01	0.26	0.06	0.07	—
14. Conopodium denudatum ..	—	0.44	—	—	0.27	—	0.50	3.61	—	—	1.79	0.19	—	0.81	0.46
15. Heracleum sphondylium ..	—	—	—	0.83	—	—	0.06	—	—	—	2.08	—	—	—	—
17. Galium verum ..	—	—	—	—	—	—	0.25	—	—	—	0.58	—	—	0.12	—
18. Scabiosa arvensis ..	—	0.16	—	—	0.43	0.15	0.12	—	0.20	0.16	4.16	0.50	—	0.23	0.26
19. Achillea millefolium ..	—	0.19	2.33	1.80	1.91	1.69	2.32	1.81	9.59	—	—	—	—	—	—
20. Centaurea nigra ..	—	—	—	—	—	—	—	—	—	—	—	0.50	—	—	—
26. Taraxacum vulgare ..	—	—	—	0.26	—	—	—	—	—	—	—	0.21	—	—	—
29. Plantago lanceolata ..	—	—	0.22	15.40	18.45	6.26	11.24	14.05	4.02	5.71	8.16	—	—	2.47	13.10
32. Prunella vulgaris ..	—	—	—	0.07	0.09	0.08	0.13	—	0.06	—	0.07	0.13	11.66	0.06	0.13
34. Rumex acetosa ..	24.61	8.62	—	—	—	—	—	—	—	—	—	—	—	—	—
35. Luzula campestris ..	—	0.05	—	—	—	—	—	—	—	—	—	—	—	—	—

PLOT 18 (continued)

	1924						1925						1926						1927					
	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL
Unlimed (U); lightly limed (LL); heavily limed (HL).																								
1. <i>Agrostis vulgaris</i>	72.70	36.81	26.46	63.03	36.52	13.65	47.05	17.54	11.54	74.63	21.56	10.30	58.99	17.51	4.83
2. <i>Aira caespitosa</i>	4.64	22.65	21.64	4.30	13.91	14.06	6.47	25.07	24.60	3.85	22.23	19.39	0.16	22.68
3. <i>Alopecurus pratensis</i>	4.12	0.14	..	3.27	0.08	0.04	4.38	0.14	0.12	2.76	0.11	..	6.88	0.50
4. <i>Anthoxanthum odoratum</i>	0.28	2.13	10.11	1.34	7.50	1.43	2.38	9.19	11.30	1.56	10.35	19.28	0.40	2.86	18.08
5. <i>Arrhenatherum avenaceum</i>
6. <i>Avena flavesens</i>	0.06	0.02	0.04	0.28	1.06	0.05	0.09	0.19
7. " <i>pubescens</i>
8. <i>Briza media</i>
11. <i>Dactylis glomerata</i>	3.66	13.51	19.53	7.57	20.67	53.48	10.81	30.55	37.38	9.55	32.28	40.02	9.34	21.08	37.72
12. <i>Festuca ovina</i>	5.41	5.55	3.94	6.31	5.12	5.11	2.71	1.66	1.95	2.84	3.40	2.91	6.22	2.79	3.13
14. <i>Holcus lanatus</i>	0.25	0.17	0.03	0.83	0.40	0.29	3.13	1.31	2.40	2.21	3.33	1.71	8.18	0.82	1.64
16. <i>Foa pratensis</i>	0.64	5.27	4.92	0.31	4.30	3.75	0.83	3.74	3.49	0.70	3.46	3.72	0.87	5.36	6.30
17. " <i>trivialis</i>	0.06
<i>Leguminosae</i>																								
1. <i>Lathyrus pratensis</i>	0.06	-	0.17	0.04	-	0.24	-	0.04	0.03	-	-	0.14	-	-	0.21
<i>Other orders</i>																								
5. <i>Stellaria media</i>	0.10	..	0.04	0.17	0.35	0.15	0.05	..	0.02	0.03
14. <i>Conopodium denudatum</i>	2.48	3.05	0.27	0.04	0.53	0.74	1.17	0.07	0.40	0.29	0.07	0.93	0.07	0.75	0.39
15. <i>Heracleum sphondylium</i>	3.96	..	0.67	0.37	3.13
17. <i>Galium verum</i>	0.14
18. <i>Scabiosa arvensis</i>
19. <i>Achillea millefolium</i>	0.09	0.28	0.15	0.08	0.56	0.03
20. <i>Centaurea nigra</i>	0.18	1.18	0.83	1.93	0.15	1.31	0.24	0.12	0.47	0.33
26. <i>Taraxacum vulgare</i>	0.08	0.38	0.18
27. <i>Tragopogon pratensis</i>	0.03
29. <i>Plantago lanceolata</i>
34. <i>Rumex acetosa</i>	2.28	9.70	9.18	9.85	10.65	5.15	20.90	8.60	4.98	1.01	1.36	0.04	1.05
35. <i>Luzula campestris</i>	0.74

	Unlimed (U); lightly limed (LL); heavily limed (HL)	1919			1920			1921			1922			1923		
		U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL
<i>Gramineae</i>																
1. Agrostis vulgaris	..	7.52	17.82	15.01	15.04	21.77	17.75	13.87	13.49	16.44	10.25	14.28	14.71	10.28	0.06	
2. Aira caespitosa	22.36	16.43	15.15	22.02	13.21	20.29	16.35	21.85	20.40	18.79	16.14	14.61	1.37
3. Alopecurus pratensis	4.27	5.27	4.15	1.68	10.27	8.32	2.93	3.66	1.91	0.62	11.57	9.37	7.13
4. Anthoxanthum odoratum	7.91	0.93	4.15	5.37	8.24	0.30	13.76	11.39	1.85	5.02	8.58	1.95	8.73
5. Arrhenatherum avenaceum	3.17	4.65	4.29	5.37	4.98	7.72	10.49	7.14	9.89	17.65	2.00	6.64	4.34
6. Avena flavaescens	3.32	3.41	4.43	4.70	3.89	1.94	4.63	1.64	1.42	2.90	1.93	2.67	..
7. " pubescens
8. Briza media	0.20	0.08	0.47	0.13	0.78	0.42	0.79	1.64	..	1.31	1.00	0.92	1.67
9. Bromus mollis
10. Cynosurus cristatus	15.84	11.31	9.25	9.00	5.60	15.98	9.02	4.91	15.09	11.85	5.28	6.38	3.66
11. Dactylis glomerata	6.10	11.92	11.79	15.31	12.45	9.84	15.51	17.68	15.70	15.04	13.28	14.84	16.59
12. Festuca ovina	2.14	1.94	1.88	2.01	5.44	6.56	1.53	1.12	2.11	0.45	3.36	2.08	0.18
14. Holcus lanatus	0.43	0.20	..	0.39	0.13	0.13	0.31	..
15. Lolium perenne	0.55	1.24	0.47	2.01	0.47	0.79	1.08	0.79	2.53	1.88	0.58	2.01	2.17
16. Poa pratensis	1.43	1.39	0.40	2.15	1.71	2.43	1.81	0.33	0.68	0.45	0.50	1.43	1.30
17. " trivialis
<i>Leguminosae</i>																
1. Lathyrus pratensis	6.02	12.39	17.96	8.33	4.82	2.07	3.38	7.46	5.93	6.95	7.29	9.25	15.54
2. Lotus corniculatus	0.27	0.33
5. Trifolium repens	0.12	2.87	0.40	0.73	0.31	0.18	0.56	0.49	..
<i>Other orders</i>																
1 & 2. Ranunculus spp.	4.75	2.86	2.09	1.14	0.39	1.34	0.73	1.25	1.49	1.31	4.57	5.46	4.89
4. Cerastium vulgatum	0.28	0.93	1.01	0.27	..	0.06	..	0.07	0.12	..	0.14	0.19	0.19
5. Stellaria graminea	0.04	0.13	0.20	0.07
11. Poterium sanguisorba
13. Anthriscus sylvestris	1.75	0.31	0.27	0.40	0.39	0.67	0.17	0.13
14. Conopodium denudatum
15. Heracleum sphondylium
17. Galium verum	1.38	1.70	1.21	1.08	0.08	0.18	0.23
19. Achillea millefolium	0.39	..	2.69	0.13	1.71	0.12	0.28	0.39	0.31	0.80	0.50
20. Centaurea nigra	0.12	..	0.53	0.60	0.31	..	0.17	0.73	0.44	0.51	0.43	0.32	0.56
26. Taraxacum vulgare	0.28	..	0.40	0.33	0.23	..	0.23	0.07	..	0.06	0.14
27. Tragopogon pratensis	0.24	0.31	0.47	0.40	0.39	0.18	0.11	0.58	0.24	0.51	0.79	0.39	1.92
29. Plantago lanceolata	0.20	0.38	0.33	..	0.23	0.06	..	0.21	0.26	..
30. Veronica chamaedrys	0.16	0.31	0.19	0.43
31. Ajuga reptans	8.40	1.47	0.60	0.94	1.95	2.37	0.34	2.09	1.98	0.40	4.29	2.14	0.25
34. Rumex acetosa	0.12	0.08	0.20	0.13	0.15	..	0.11	0.26	0.06	0.06	0.43	0.39	0.31

PLOT 19 (continued)	Unlimed (U); lightly limed (LL);		1924		1925		1926		1927		1928	
	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL
<i>Gramineae</i>												
1. Agrostis vulgaris	12.79	18.29	8.20	10.72	9.53	3.38	8.57	9.49	2.52	5.46
3. Alopecurus pratensis	16.93	16.17	16.71	16.38	22.48	24.34	29.41	29.75	27.20	25.83
4. Anthoxanthum odoratum	3.95	7.18	0.61	9.52	5.12	0.80	7.63	7.67	0.73	9.29
5. Arrhenatherum avenaceum	10.62	1.02	8.01	18.50	2.70	15.48	11.83	6.01	21.54	17.68
6. Avena flavescens	3.91	3.68	7.28	1.25	0.99	2.11	0.85	1.17	2.24	1.91
7. " pubescens	1.44	1.76	5.90	1.69	2.62	6.70	1.95	2.89	7.19	2.92
8. Briza media
9. Bromus mollis	0.04
11. Dactylis glomerata	7.06	4.40	4.66	9.30	20.97	18.39	14.33	15.31	15.76	17.01
12. Festuca ovina	8.08	9.79	12.94	7.67	5.04	6.83	4.46	2.67	4.31	6.18
14. Holcus lanatus	1.32	2.27	0.77	6.15	1.75	1.68	3.80	2.21	1.28	3.35
15. Lolium perenne	0.04	..	0.16	0.13	0.13	0.10
16. Poa pratensis	0.53	1.33	2.58	0.33	1.51	1.86	0.37	1.17	1.63	0.29
17. " trivialis	0.27	0.72	1.34	0.93	3.02	3.21	1.66	5.59	5.85	0.29
<i>Leguminosae</i>												
1. Lathyrus pratensis	19.50	17.83	20.56	3.86	3.14	1.31	2.29	1.33	0.86	1.15
2. Lotus corniculatus	0.23	2.14	0.81	0.01	0.04	0.21	0.11	0.16
5. Trifolium repens	0.32	0.16	0.14
<i>Other orders</i>												
1 & 2. Ranunculus spp.	3.91	7.44	3.39	5.39	11.12	5.44	2.29	4.26	2.11	1.53
4. Ceratium vulgatum	0.15	1.12	..	0.11	0.04	0.13	0.33	0.26	..	0.10
5. Stellaria graminea	0.04	0.04	0.04	0.03	0.04	0.03
6. " holostea
13. Anthriscus sylvestris	0.98	0.21	0.54	1.36	0.84	2.49	0.18	0.33	0.86	1.53
14. Conopodium denudatum	2.21	1.89	1.66	1.47	0.83	0.84	1.00	1.50	0.70	1.15
15. Heracleum sphondylium	0.23	0.13	..	0.07	0.48
16. Pimpinella saxifraga	0.19	0.34	0.19	0.11	0.04	0.03	0.03
19. Achillea millefolium	0.05	0.72	0.04	0.25	0.42	0.34	0.04
20. Centaurea nigra	0.12	0.26	0.07	0.15
26. Taraxacum vulgare	0.76	0.30	2.31	0.27	0.08	1.65	0.11	0.03	0.48	0.05
27. Tragopogon pratensis	0.15	0.22	..	0.04
29. Plantago lanceolata	0.53	0.04	0.23	0.11	..	0.63	0.66	0.06	0.42	0.29
30. Veronica chamaedrys	0.23	0.17	0.03	0.16	0.08	0.04	0.33	0.42	..	0.19
31. Ajuga reptans	0.15	0.17	0.12	0.22	0.40	0.13	0.19	0.72	0.10	0.24
34. Rumex acetosa	1.83	0.85	0.38	3.97	7.50	1.77	7.04	6.31	2.78	2.49
35. Luzula campestris	0.15	0.09	0.19	0.11	0.04	0.03	0.15	0.03	0.10	0.04
36. Carex praecox

PLOT 20—BOTANICAL COMPOSITION PER CENT. (INDIVIDUAL SPECIES)

	Unlimed (U); lightly limed (LL);		1919		1920		1921		1922		1923	
	U	HL	U	LL	HL	U	LL	HL	U	LL	HL	
<i>Gramineae</i>												
1. <i>Agrostis vulgaris</i> ..	6.57	10.96	14.70	6.10	13.38	10.50	4.87	9.79	11.51	4.86	10.44	12.70
2. <i>Alopecurus pratensis</i> ..	30.34	26.82	21.72	30.26	19.43	22.05	30.14	23.30	23.79	33.27	29.36	16.88
4. <i>Anthoxanthum odoratum</i> ..	1.16	0.78	2.25	0.75	1.08	6.40	1.76	1.21	0.37	0.34	0.34	25.23
5. <i>Arrhenatherum avenaceum</i> ..	4.60	5.80	2.38	2.49	10.05	9.56	3.93	6.81	5.53	2.96	5.56	5.71
6. <i>Avena flavescens</i> ..	2.94	4.64	2.91	3.74	6.70	2.94	5.32	9.04	7.46	7.07	6.14	1.21
7. " <i>pubescens</i> ..	10.17	11.22	10.07	14.20	12.32	8.03	18.58	7.77	5.15	7.41	7.71	5.28
9. <i>Bromus mollis</i> ..	0.23	0.19	0.26	0.56	0.47	1.21	0.72	1.76	2.19	2.81	2.44	11.65
11. <i>Dactylis glomerata</i> ..	12.52	9.87	7.68	9.09	5.62	8.76	8.50	9.04	10.74	8.16	6.00	2.87
12. <i>Festuca ovina</i> ..	4.44	10.06	9.80	9.96	8.71	8.71	7.07	15.64	13.59	12.64	13.85	4.08
14. <i>Holcus lanatus</i> ..	6.85	3.35	4.77	6.60	10.05	10.29	7.73	3.51	4.17	3.06	1.95	2.70
15. <i>Lolium perenne</i> ..	0.30	0.26	0.07	0.12	0.40	0.42	0.11	0.37	0.33	0.10	0.79	1.90
16. <i>Poa pratensis</i> ..	0.62	1.93	0.46	1.50	0.73	0.26	0.87	3.20	0.61	2.61	2.44	0.23
17. " <i>trivialis</i> ..	1.09	1.61	1.59	0.94	1.21	1.84	0.99	0.32	0.54	0.40	1.13	1.06
<i>Leguminosae</i>												
1. <i>Lathyrus pratensis</i> ..	4.72	3.87	15.10	4.60	3.35	4.68	3.51	0.95	7.89	4.41	1.95	9.88
2. <i>Lotus corniculatus</i> ..	—	—	—	—	0.27	—	—	0.16	—	—	—	5.33
4. <i>Trifolium pratense</i> ..	—	—	0.64	0.20	0.06	0.40	0.16	—	—	0.20	0.29	—
6. " <i>repens</i> ..	—	—	—	—	—	—	—	0.05	0.05	—	0.24	—
1. & 2. <i>Ranunculus</i> spp.												
4. <i>Cerastium vulgatum</i> ..	2.13	0.84	0.66	0.69	0.94	0.84	1.15	1.02	0.99	1.30	1.95	2.59
5. <i>Stellaria graminea</i> ..	0.25	0.26	0.46	0.56	—	—	0.05	—	0.05	—	—	0.06
13. <i>Anthriscus sylvestris</i> ..	2.50	0.78	0.07	1.06	0.20	0.10	0.93	0.69	0.27	2.15	2.68	3.27
14. <i>Conopodium denudatum</i> ..	1.50	0.13	0.26	0.50	0.87	0.47	1.04	1.70	1.32	2.46	0.78	1.55
15. <i>Heracleum sphondylium</i> ..	—	—	—	0.25	0.27	—	—	—	—	—	—	—
17. <i>Galium verum</i> ..	0.47	0.06	—	0.43	—	—	0.06	0.42	—	0.81	—	0.17
19. <i>Achillea millefolium</i> ..	1.19	2.71	0.73	1.25	0.73	0.16	1.16	1.70	0.49	0.30	0.68	0.33
20. <i>Centaurea nigra</i> ..	1.44	1.93	1.40	2.61	0.67	—	0.49	0.74	—	0.50	0.49	1.03
26. <i>Taraxacum vulgare</i> ..	0.05	—	—	0.12	—	0.31	0.60	0.16	0.27	0.72	0.24	1.21
27. <i>Tragopogon pratensis</i> ..	0.16	—	1.20	0.50	—	0.47	0.16	—	0.16	0.35	0.10	1.32
29. <i>Plantago lanceolata</i> ..	0.06	—	0.13	0.06	0.47	0.42	—	0.32	0.33	0.15	0.10	—
30. <i>Veronica chamaedrys</i> ..	0.31	—	0.13	0.06	0.13	—	—	0.05	0.05	—	0.10	0.06
34. <i>Rumex acetosa</i> ..	3.31	1.29	0.73	0.94	1.48	1.32	1.04	1.23	1.26	0.81	2.15	0.82
35. <i>Luzula campestris</i> ..	0.03	—	—	0.07	—	0.07	0.05	0.11	0.05	0.15	0.10	0.17

PLOT 20 (continued)

	Unlimed (U); lightly limed (LL);		Heavily limed (HL)		1924		1925		1926		1927		1928		1929			
	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U	LL	HL	U		
<i>Gramineae</i>																		
1. Agrostis vulgaris	..	15.01	9.22	4.26	13.80	3.55	3.00	9.51	4.71	3.57	3.73	2.59	1.87	4.08	4.21	1.17	5.48	
3. Alopecurus pratensis	..	16.01	27.53	20.95	16.61	15.43	19.24	27.45	21.54	25.94	30.25	18.58	23.39	46.23	30.65	24.12	33.88	
4. Anthoxanthum odoratum	..	0.96	1.34	0.23	1.08	1.82	0.41	1.34	3.72	0.97	0.65	3.22	0.84	1.89	6.37	0.73	0.61	
5. Arrhenatherum avenaceum	..	5.97	10.17	0.39	8.88	32.59	3.64	17.62	23.83	8.22	24.58	31.04	7.43	11.28	8.74	3.83	8.85	
6. Avena flavescens	..	4.66	2.83	3.17	1.48	0.58	1.09	0.88	1.40	1.45	2.59	1.59	2.60	3.42	4.11	2.58	3.81	
7. " pubescens	..	5.33	4.86	10.44	6.32	2.58	15.83	5.61	6.82	15.97	4.83	8.00	23.72	6.63	9.35	28.92	8.10	
9. Bromus mollis	0.04	0.06	..	0.08	..	0.04	0.03	0.26	0.20	0.10
10. Cynosurus cristatus
11. Dactylis glomerata	..	6.09	5.31	4.87	10.43	13.38	9.08	11.10	13.71	12.84	14.63	17.90	14.01	7.00	7.12	5.61	9.33	
12. Festuca ovina	..	8.04	6.10	9.90	4.48	1.47	4.39	3.29	3.02	5.38	3.24	2.63	4.76	5.31	6.47	9.04	5.96	
14. Holcus lanatus	..	1.93	2.11	1.46	4.48	4.45	4.76	6.56	6.54	7.62	7.22	7.21	6.96	4.05	7.89	5.00	2.04	
15. Lolium perenne	..	0.21	0.36	0.19	0.04	0.04	—	0.06	0.07	0.03	0.05	0.04	0.07	0.43	0.32	0.08	0.17	
16. Poa pratensis	..	0.92	0.65	1.59	0.96	0.27	1.20	1.89	0.77	1.93	1.10	0.52	2.31	1.43	0.65	2.90	1.36	
17. " trivialis	..	0.35	0.65	0.39	0.28	0.62	0.64	2.35	4.36	3.66	0.35	0.88	0.77	1.23	3.46	2.99	1.40	
<i>Leguminosae</i>																		
1. Lathyrus pratensis	..	21.59	16.70	30.47	10.32	2.49	4.20	3.54	1.09	3.41	1.64	0.52	3.22	2.59	1.78	5.13	6.95	
2. Lotus corniculatus	
4. Trifolium pratense	..	0.67	
5. " repens.	..	1.10	0.11	
<i>Other orders</i>																		
1. & 2. Ranunculus spp.	..	2.89	4.14	2.13	3.80	3.56	9.00	1.77	1.72	1.84	0.55	0.56	0.92	0.76	1.29	1.21	0.65	
4. Cerastium vulgatum	..	0.04	0.11	0.08	—	0.13	0.08	—	0.35	0.18	—	0.08	0.11	0.03	0.26	0.24	0.44	
5. Stellaria graminea	
13. Anthriscus sylvestris	..	1.39	1.12	2.36	4.56	6.00	7.88	1.22	0.91	0.97	0.95	0.28	0.62	0.40	0.32	0.28	0.37	
14. Conopodium denudatum	..	1.54	1.46	1.67	1.40	0.89	0.86	0.58	0.31	0.39	0.65	0.76	0.84	0.73	0.68	0.49	0.72	
15. Heracleum sphondylium	
17. Galium verum	
18. Scabiosa arvensis	
19. Achillea millefolium	..	1.17	0.80	
20. Centaurea nigra	
26. Taraxacum vulgare	..	0.53	2.29	2.55	0.24	3.02	6.38	0.18	0.95	1.21	—	0.36	1.39	0.10	1.00	2.16	0.17	
27. Tragopogon pratensis	..	0.18	1.01	1.27	0.08	0.62	2.70	0.03	0.56	0.60	—	0.12	0.88	—	0.13	0.32	—	
29. Plantago lanceolata	
30. Veronica chamaedrys	..	0.07	
31. Ajuga reptans	..	0.04	0.04	
34. Rumex acetosa	..	1.24	0.76	0.62	2.24	4.85	2.17	3.17	2.57	1.33	1.64	0.48	0.37	1.43	1.62	0.06	—	
35. Luzula campestris	

WHEAT—BROADBALK FIELD, 1934

Plot.	Manurial Treatment (amounts stated are per acre).	Dressed Grain, bushels per acre (in some cases estimated from half or quarter-bushel).					Total Grain, cwt. per acre					74-year Average 1852-1925 (prior to fallow). Total Grain, cwt.				
		I	II	III	V	Mean	I	II	III	V	Mean	I	II	III	IV.	V.
2A	Farmyard Manure (14 tons)	34.6	44.6	42.5	23.3	36.2	21.4	27.6	25.7	16.5	22.8	16.3**	19.4	19.4		
2B	Farmyard Manure (14 tons)	32.2	36.1	37.6	22.8	32.2	19.9	23.3	24.0	15.9	20.8					
3	Unmanured since 1839	17.2	16.6	21.4	29.7	21.2	9.9	10.5	13.2	18.4	13.0	6.7				
5	Complete Mineral Manure \$	12.5	15.3	13.3	24.6	16.4	8.2	9.7	8.6	14.9	10.4	7.8				
6	As 5, and 206 lb. Sulphate of Ammonia	26.1	30.9	31.3	28.7	29.3	16.0	18.8	19.1	18.0	18.0	12.5				
7	As 5, and 412 lb. Sulphate of Ammonia	45.2	46.1	44.8	27.1	40.8	27.7	28.6	27.1	18.0	25.4	17.6				
8	As 5, and 618 lb. Sulphate of Ammonia	43.1	53.2	52.4	28.8	44.4	26.9	33.8	32.1	18.3	27.8	20.1				
9	As 5, and 275 lb. Nitrate of Soda	32.8	35.3	29.6	32.6	32.6	19.7	21.9	18.4	20.4	20.1	13.9†				
10	412 lb. Sulphate of Ammonia	39.2	45.6	44.2	32.1	40.3	25.2	27.8	27.5	20.7	25.3	10.9				
11	As 10, and Superphosphate (3½ cwt.)	35.8	36.1	37.3	29.2	34.6	21.6	22.1	23.1	20.1	21.7	12.3				
12	As 10, and Super (3½ cwt.) and Sulph. Soda (366 lb.)	40.2	45.1	44.7	33.7	40.9	24.2	27.1	26.7	22.7	25.2	15.7				
13	As 10 and Super (3½ cwt.) and Sulph. Potash (200 lb.)	38.0	47.5	42.6	28.6	39.2	23.1	28.6	26.1	19.5	24.3	17.0				
14	As 10, and Super (3½ cwt.) and Sulph. Magnesia (280 lb.)	42.6	46.3	43.6	34.7	41.8	25.4	27.1	26.2	22.9	25.4	15.5				
15	As 5, and 412 lb. Sulphate Amm. all applied in Autumn	33.6	48.7	41.0	34.3	39.4	21.0	28.7	24.7	22.4	24.2	16.1				
16	As 5, and 550 lb. Nitrate of Soda	41.5	49.8	43.4	29.6	41.1	25.5	30.4	26.2	19.2	25.3	17.8††				
17	{ Minerals alone as 5 or 412 lb. Sulphate of Ammonia alone in alternate years	A36.1	41.2	37.9	34.2	37.4	21.9	24.8	23.6	21.7	23.0	A16.1*				
18	{ Rape Cake (1,889 lb.)	M13.2	18.7	14.5	36.0	20.6	8.3	11.5	9.1	22.6	12.9	M8.1				
19	As 7, without Supper	35.9	40.2	32.4	26.7	33.8	21.9	24.3	20.2	17.7	21.0	12.6†				
20		31.9	40.6	—	—	36.2	19.8	24.8	—	—	22.3	10.3§				

FALLOWING ROTATION. After the fallows of 1925-6 to 1928-9 a regular cycle of fallowing was started in the season 1930-31. This cycle and the preceding fallows are shown in the accompanying diagram (C = crop, F = fallow). The sections (I. to V.) are numbered in order from the upper or western end of the field. Preparatory to the first fallow the field was harvested in five separate sections (1924-5).

For notes, see next page.

BARLEY—HOOS FIELD, 1934

Plot	Manuring (Amounts stated are per acre)	Dressed Grains, cwt., per acre									
		16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928	16 Years, Average, 1852-1928
10	Unmanured
20	Superphosphate only (3½ cwt.)
30	Alkali Salts only (200 lb. Sulphate of Potash; 100 lb. Sulphate of Soda; 100 lb. Sulphate of Magnesia)
40	Complete Minerals, as 30 with Superphosphate (3½ cwt.)
50	Potash (200 lb.) and Superphosphate (3½ cwt.)
IA	Ammonium Salts only (206 lb. Sulphate of Ammonia)
2A	Superphosphate and Ann. Salts
3A	Alkali Salts and Ann. Salts
4A	Complete Minerals and Ann. Salts
6A	Potash Super and Ann. Salts
1AA	Nitrate of Soda only (275 lb.)
2AA	Superphosphate and Nitrate of Soda
3AA	Alkali Salts and Nitrate of Soda
4AA	Complete Minerals and Nitrate of Soda
IAAS	As Plot IAA and Silicate of Soda (400 lb.)
2AAS	As Plot 2AA and Silicate of Soda (400 lb.)
3AAS	As Plot 3AA and Silicate of Soda (400 lb.)
4AAS	As Plot 4AA and Silicate of Soda (400 lb.)
1C	Rape Cake only (1,000 lb.)
2C	Superphosphate and Rape Cake
3C	Alkali Salts and Rape Cake
4C	Complete Minerals and Rape Cake
7-1	Unmanured after dung (14 tons) for 20 years (1852-71)
7-2	Farmyard manure (14 tons)
6-1	Unmanured since 1852
6-2	Ashes from Laboratory furnace
1N	Nitrate of Soda only (275 lb.)
2N	Nitrate of Soda only (275 lb.)

In 1912 and 1933 all plots were fallowed. || 1 cwt. 2.15 bushels. † Total straw includes straw, cavings and chaff. * Sixty years 1868-1928. ‡ 56 years, 1872-1928. § 75 years, 1853-1928.

|| 69 years, 1859-1928.

†† Sulphate of ammonia was applied to four-fifths of this plot by mistake. The yields given refer to the unmanured portion, the yields on the manured portion being :- grain:

31.7 bushels or 17.0 cwt. per acre and straw: 21.3 cwt. per acre. The weight per bushel was 65.2 lb. Harrowed: February 2. Ploughed: Mar. 27 and 28. Rolled: March 27 and 28. Manures applied: January 29, March 26 and 27. Seed sown: March 27 and 28. Variety: Plumage Archer. Harvested: August 9 and 10.