

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Rothamsted Report for 1936

[Full Table of Content](#)



Other Experiments at Rothamsted

Rothamsted Research

Rothamsted Research (1937) *Other Experiments at Rothamsted* ; Rothamsted Report For 1936, pp 205 - 223 - DOI: <https://doi.org/10.23637/ERADOC-1-68>

WHEAT

**Effect of sulphate of ammonia applied at five different times
RW—Great Knott, 1936**

Plan and yields in lb., grain above, straw below

		3	5	1	2	0	4	
		76.9	72.1	81.6	80.3	78.7	75.9	
		129.6	116.2	113.6	133.7	96.0	117.1	
		5	4	0	1	3	2	
		86.3	82.1	84.0	91.1	88.8	89.2	
		120.7	117.6	106.2	126.4	139.4	140.8	
W	↑	2	1	3	0	4	5	
		75.2	84.6	77.9	86.3	77.5	84.3	
		134.6	110.9	141.1	121.4	123.8	126.7	
		4	0	5	3	2	1	
		82.0	85.4	90.1	83.8	84.5	96.4	
		127.0	117.8	129.6	146.0	148.5	139.8	
		1	3	2	4	5	0	
		87.8	76.6	85.9	84.2	86.0	88.3	
		120.2	146.4	156.8	136.8	135.2	119.7	
		0	2	4	5	1	3	
		91.9	77.2	94.5	99.1	95.2	89.3	
31		125.1	148.0	139.0	138.4	148.3	150.2	36

SYSTEM OF REPLICATION : 6 × 6 Latin square.

AREA OF EACH PLOT : 1/40 acre (53.0 lks. × 47.2 lks.).

TREATMENTS : No sulphate of ammonia (0) and sulphate of ammonia at the rate of 0.4 cwt. N per acre applied on Oct. 28 (1), Jan. 20 (2), Mar. 16 (3), Apr. 27 (4) and May 25 (5).

CULTIVATIONS, ETC. : Ploughed on various dates during Sept. Harrowed : Oct. 28 and April 16. Drilled : Oct. 28. Harvested : Aug. 18. Variety : Victor. Previous crop : Beans.

STANDARD ERRORS PER PLOT : Grain : 1.22 cwt. per acre or 4.02% ; Straw : 1.94 cwt. per acre or 4.18%.

Summary of Result : cwt. per acre

	No N	Dates of application of sulphate of ammonia (0.4 cwt. N per acre)					Mean of all N	Standard error
		Oct. 28	Jan. 20	Mar. 16	Apr. 27	May 25		
GRAIN								
	(±0.498)	30.6	31.9	29.3	29.4	29.5	30.8	30.2 ±0.223
Incr. ..	(±0.704)		+1.3	-1.3	-1.2	-1.1	+0.2	-0.4 ±0.546
STRAW								
	(±0.792)	40.8	45.2	51.3	50.8	45.3	45.6	47.6 ±0.354
Incr. ..	(±1.12)		+4.4	+10.5	+10.0	+4.5	+4.8	+6.8 ±0.868

Conclusions

On the average of all times of application, sulphate of ammonia had little effect on the yield of grain, and gave a significant increase in straw. There were, however, significant differences in the yields obtained with the different times of application, the earliest dressing (Oct. 28) producing an increase in the yield of grain which was not quite significant, and the next three times of application (Jan. 20, Mar. 16, and April 27) producing a depression in the yield of grain.

The straw yields at first increased to a maximum and then decreased with later application.

WHEAT

Effect of fallow and of one-year leys of clover, ryegrass and clover and ryegrass, cut once and followed by bastard fallow or green manure crops of mustard or vetches, or cut twice.

RW—Pastures—1936
Plan and yields in lb.

		Grain		Straw				Grain		Straw							
133	OM	N	18.6	48.9	O	11.6	31.2	101	69	CR2	O	19.5	42.9	N	22.1	46.9	37
	C2	O	18.9	43.8	N	15.1	47.9			CR1	N	17.7	45.8	O	17.6	36.9	
	O2	N	17.9	44.6	O	8.7	29.8			O1	O	14.7	34.4	N	23.2	46.8	
	R1	O	8.8	29.2	N	12.1	39.4			CM	N	11.6	46.9	O	15.2	34.3	
	OV	O	15.1	39.2	N	16.8	49.4			CV	O	7.5	33.2	N	13.6	38.4	
	RV	O	11.3	37.2	N	10.7	41.8			CRM	O	12.4	37.1	N	14.8	39.2	
	CRM	O	12.8	38.2	N	12.2	47.0			R1	N	14.2	40.0	O	14.5	27.7	
	CR2	O	13.4	35.4	N	14.8	47.4			OV	O	15.1	37.6	N	18.8	55.0	
	RM	O	13.2	30.3	N	12.0	43.0			O2	O	13.8	38.1	N	21.2	49.3	
	O1	O	23.1	46.7	N	22.8	55.7			OM	O	16.1	39.2	N	21.5	47.8	
	R2	N	16.3	41.4	O	12.4	28.5			C1	O	12.4	42.2	N	16.4	43.8	
	CR1	O	14.0	37.9	N	13.4	46.6			CRV	N	11.6	44.9	O	9.0	31.0	
	CV	N	11.0	63.5	O	9.0	40.2			R2	O	11.8	30.2	N	13.0	31.0	
	CRV	N	10.8	56.4	O	6.7	35.0			C2	O	20.6	47.1	N	21.0	49.2	
	CM	N	10.7	51.8	O	10.6	39.4			RV	N	11.8	42.4	O	8.9	33.4	
	C1	N	13.5	57.0	O	10.5	37.8			RM	O	5.5	27.5	N	14.8	40.4	
	R1	O	9.9	29.5	N	11.8	41.0			O2	O	12.9	43.0	N	20.4	55.6	
	OM	O	14.2	37.3	N	17.4	51.6			CR1	N	8.3	34.0	O	9.0	31.7	
	CM	N	14.4	49.1	O	9.6	35.4			OV	O	14.3	45.7	N	19.0	53.0	
	C2	O	21.1	46.0	N	18.6	45.9			R1	N	7.4	28.8	O	9.1	22.0	
	C1	O	17.1	44.8	N	16.0	49.2			OM	N	21.0	50.8	O	14.5	35.5	
	CRV	O	13.0	40.8	N	12.9	49.1			R2	N	17.3	42.4	O	9.2	23.8	
	RM	O	19.2	41.0	N	15.2	45.0			RV	O	7.8	28.2	N	6.9	30.4	
	CR1	O	18.2	40.3	N	17.6	47.9			C1	O	12.7	36.6	N	13.9	41.8	
	O2	N	24.4	51.4	O	18.5	38.2			CRV	O	8.3	37.7	N	9.1	45.9	
	O1	N	26.1	56.6	O	17.2	36.3			CV	O	10.8	44.7	N	12.3	46.7	
	CV	O	15.5	40.8	N	12.0	47.5			CR2	N	19.4	39.8	O	17.1	31.9	
	CRM	N	17.2	45.8	O	9.0	23.5			CM	N	14.1	48.4	O	13.6	31.9	
	CR2	O	16.7	34.2	N	17.9	41.8			CRM	N	15.1	43.2	O	12.0	28.2	
	RV	N	11.9	42.1	O	10.9	26.1			RM	N	20.4	42.6	O	14.0	26.2	
164	OV	N	19.5	54.8	O	17.5	32.5	132	100	O1	O	24.1	48.3	N	23.9	53.1	68
	R2	O	13.6	30.2	N	16.7	34.8			C2	O	20.4	37.9	N	13.4	33.4	

SYSTEM OF REPLICATION ; 4 randomised blocks of 16 plots each, the plots being split for sulphate of ammonia at the rate of 0.3 cwt. N per acre applied as a top dressing to the wheat.

AREA OF EACH SUB-PLOT ; 0.0099 acre (46.8 lks. × 21.2 lks.)

TREATMENTS : 4 × 4 factorial design.

No ley (O), broad red clover (C), ryegrass (R), ryegrass and clover (CR). Vetches sown after 1st cut. (V), mustard sown after 1st cut (M), 1 cut taken, followed by summer fallow (1), 2 cuts taken (2).

BASAL MANURING ; Nil.

CULTIVATIONS, ETC. ; Ploughed ; Sept. 28-30. Harrowed ; Oct. 19 and 28. Drilled ; Oct. 28.

Sulphate of ammonia applied ; March 28. Harrowed and rolled ; May 5. Harvested ; Aug. 24 and 25. Variety ; Victor. Previous crop ; Ley.

STANDARD ERRORS : Grain : Per whole plot : 2.07 cwt. per acre or 15.7% ; per sub plot : 1.96 cwt. per acre or 14.9%. Straw : per whole plot : 3.62 cwt. per acre or 9.87% ; per sub plot : 3.68 cwt. per acre or 10.0%.

Green manure crops : Nitrogen added lb. per acre.

	Fallow	Clover	Ryegrass	Clover and ryegrass	Mean
Mustard	11.8	13.0	10.9	11.7	11.9
Vetches	9.2	12.8	12.2	13.0	11.8

	No nitrogen				Nitrogen			
	1 cut	2 cuts	Mustard	Vetches	1 cut	2 cuts	Mustard	Vetches
GRAIN ; cwt. per acre ($\pm 1.04^1$, $\pm 1.39^2$)								
Fallow	15.0	12.7	13.9	20.2	17.7	16.7		
Clover	11.8	18.2	11.0	9.6	13.4	15.3	11.4	11.0
Ryegrass	9.5	10.6	11.7	8.8	10.2	14.2	14.0	9.3
Clover and ryegrass	13.2	15.0	10.4	8.3	12.8	16.7	13.3	10.0
STRAW ; cwt. per acre ($\pm 1.81^1$, $\pm 2.60^2$)								
Fallow	35.4	32.2	34.9	46.4	44.8	47.7		
Clover	36.3	39.3	31.7	35.7	43.2	39.7	44.1	44.1
Ryegrass	24.4	25.4	28.1	28.1	33.6	33.6	38.5	35.2
Clover and ryegrass	33.0	32.5	28.6	32.5	39.2	39.6	39.4	44.2

(¹)For comparisons involving the mean of nitrogen and no nitrogen.

(²)For comparisons involving the difference of nitrogen and no nitrogen.

	Fallow	Clover	Ryegrass	Clover and ryegrass	Mean	Standard errors	Fallow	Mustard	Vetches
No nitrogen	14.1	12.6	GRAIN ; cwt. per acre 10.2	11.7	12.2		12.4	11.4	10.2
Nitrogen	18.7	12.8	11.9	13.2	14.2		14.2	14.1	11.8
Mean ..	16.4	12.7	11.0	12.4	13.2	± 0.518	13.3	12.8	11.0
Increase		-3.7	-5.4	-4.0		± 0.732		-0.5	-2.3
Difference	+4.6	+0.2	+1.7	+1.5	+2.0	± 0.693	+1.8	+2.7	+1.6
Increase		-4.4	-2.9	-3.1		± 0.980		+0.9	-0.2
No nitrogen	34.5	35.8	STRAW ; cwt. per acre 26.5	31.6	32.1		32.3	30.2	32.8
Nitrogen	46.4	42.8	35.2	40.6	41.2		40.6	41.7	42.8
Mean ..	40.4	39.3	30.8	36.1	36.6	± 0.905	36.4	36.0	37.8
Increase		-1.1	-9.6	-4.3		± 1.28		-0.4	+1.4
Difference	+11.9	+7.0	+8.7	+9.0	+9.1	± 1.30	+8.3	+11.5	+10.0
Increase		-4.9	-3.2	-2.9		± 1.84		+3.2	+1.7

Conclusions

All three leys produced large reductions in the yield of grain compared with fallow, the reduction being significantly greater for ryegrass than for clover or with the clover and ryegrass mixture. The yields of grain were in all cases substantially better where two cuts of the leys were taken than with only one cut. In fact, where no nitrogen was applied to the wheat, the yields with two cuts of clover or the clover and ryegrass mixture were as good as with fallow.

The yields of grain following mustard were slightly but not significantly below those following the bastard fallow. The yields following vetches were significantly lower than those following mustard.

The response in grain to sulphate of ammonia was significantly greater after fallow than after any of the leys, the response following clover being negligible.

The effects on straw were in general similar to those on grain, except that vetches did not depress the yields of straw.

BARLEY

Soil fumigation experiment. Residual effect of chlorpicrin, chlordinitrobenzene, "seekay" and "cymag," as controls of wireworm infestation.

RB—Pastures, 1936
Plan and yields in lb.

Grain, straw, plant number (four metre rows) (April 23) and number of wireworms (total of four samples) in descending order

31	P	O	N	K	M	35
	29.0	27.8	32.1	36.0	45.6	
	48.0	54.4	82.4	66.0	89.9	
	124	120	118	217	153	
	3	2	5	1	4	
36	M	K	O	N	P	
	35.1	41.3	33.1	43.6	42.2	
	68.4	69.7	53.2	78.4	64.8	
	196	187	148	148	152	
	6	0	6	4	4	
S	O	M	K	P	N	
↑	36.9	36.0	31.6	38.9	41.2	
	59.1	60.0	58.6	54.6	81.8	
	161	208	151	177	151	
	4	9	1	6	5	
	N	P	M	O	K	
	39.3	36.6	34.5	40.3	37.6	
	77.2	60.4	62.0	65.0	55.2	
	150	166	152	177	179	
	17	8	8	9	0	
	K	N	P	M	O	
	35.8	41.1	38.1	32.5	30.7	
	60.7	71.6	64.2	55.0	55.3	
51	212	180	182	166	95	55
	4	4	2	4	8	

Note.—In the field the plots lay in one line, 36 being next to 35, etc.

SYSTEM OF REPLICATION : 5 × 5 Latin square.
 AREA OF EACH PLOT ; 1/60 acre (60.6 lks. × 27.5 lks.).
 TREATMENTS : No fumigant (O), chlordinitrobenzene (N) and chlorpicrin (P) at the rate of 2.0 cwt. per acre, "cymag" (M) at the rate of 7.5 cwt. per acre and "seekay" (K) at the rate of 5.0 cwt. per acre applied in 1935. (See 1935 Report, p. 188).
 BASAL MANURING ; Nil.
 CULTIVATIONS, ETC. ; Ploughed on various dates during February. Springtime harrowed ; March 18. Rolled, Harrowed and Drilled ; March 20. Rolled ; May 2. Harvested ; Aug. 21 and 22. Variety ; Plumage Archer. Previous crop ; Sugar beet.
 Sampling for wireworms ; Date ; June 19-July 2. Two random samples per half plot were taken, each sample consisting of 6 ins. × 6 ins. × 5 ins. (deep) of soil.
 STANDARD ERRORS PER PLOT ; Grain ; 2.45 cwt. per acre or 12.5%. Straw ; 4.74 cwt. per acre or 13.7%. Plant number : 174.8 thousands per acre or 16.2%.

Summary of Results

	No fumigant	Chlordinitrobenzene	Chlorpicrin	"Cymag"	"Seekay"	Mean	Standard error
GRAIN ; cwt. per acre	18.1	21.1	19.8	19.7	19.5	19.6	±1.10
STRAW ; cwt. per acre	30.8	41.9	31.3	35.9	33.2	34.6	±2.12
PLANT NUMBER ; thousands per acre	930.7	991.8	1063.5	1161.7	1256.0	1080.7	±78.2
NO. OF WIRE-WORMS per square yard	52	63	41	56	11	45	

Conclusions

"Seekay" applied in 1935 produced a significant decrease in the number of wireworms in June 1936. This was reflected in a significant increase in plant number, but the corresponding increase in grain was not significant, the grain yields being moderately good even in the absence of any fumigation. The somewhat higher yields of grain and straw given by chlordinitrobenzene may be attributed to an effect of nitrogen.

The other fumigants had apparently no residual effects on wireworms.

SPRING OATS

Soil fumigation experiment. Residual effect of "cymag," carbon disulphide jelly, chlordinitrobenzene and "seekay." Effect of sulphate of ammonia.

RO—Pastures, 1936

Plan and yields, total produce in grams above, weeds in lb. centre, 3rd cyst count below*

1	O	2CK	1N	1CM	2CM	2S	2CK	O	40
	79 88 248	938 Nil 232	106 128 272	275 181 206	281 291 160	4,033 230 114	2,461 57 114	105 215 266	
S	IS	O	1A	2CM	1CK	1N	1CM	1A	
	4,844 57 154	63 206 150	171 198 274	256 192 264	1,238 53 180	105 247 198	1,068 199 106	962 192 188	
↑	2S	1CK	2A	2N	O	2A	2N	IS	
	1,888 197 229	621 61 273	210 253 275	387 284 212	65 203 102	188 274 82	140 250 120	2,692 214 88	
21	1CK	1A	1S	2CK	2CK	2A	1CK	1CM	36
	885 25 242	385 147 358	868 94 252	616 9 222	338 32 160	550 245 262	495 48 196	555 191 230	
	2A	2N	2S	1N	1A	2N	2S	O	
	1,047 197 412	30 184 353	1,518 137 258	205 144 322	105 196 100	735 201 116	3,021 110 56	809 156 68	
	2CM	O	1CM	O	1S	1N	O	2CM	
	1,425 187 204	165 147 276	358 166 224	285 127 174	1,608 121 88	1,679 135 52	315 148 116	1,203 204 102	

The positions of the blocks in the field were slightly different from those shown above.

*Number of cysts per 400 grams of soil.

AREA OF EACH PLOT; 1/80 acre (30 lks. × 41.7 lks.).

TREATMENTS; No fumigant (O), single (1) and double (2) dressings of "cymag" (CM), carbon disulphide jelly (S), chlordinitrobenzene (N) and "seekay" (CK), at the following rates of application per acre for the single dressing; 1CM, 7.5 cwt.; 1S, 24.3 cwt.; 1N, 2.0 cwt. and 1CK, 5.0 cwt., applied in 1935.

Sulphate of ammonia (A) at the rate of 0.4 cwt. and 0.8 cwt. N per acre. Applied in 1936.

BASAL MANURING; Nil.

CULTIVATIONS, ETC.; Ploughed; Aug. 27-29 and Feb. 10-March 13. Spring-tine harrowed; March 19. Rolled, harrowed, sulphate of ammonia applied and drilled; March 20. Harrowed; May 8. Rolled; May 9. Harvested; Aug. 7. Variety; Marvellous. Previous crop; Oats. (See 1935 Report, p.176).

SPECIAL NOTES; The entire produce was weighed, Oats plus weeds. Weight of oats was determined by sampling a strip 30 lks. × 1 yd. Third cyst count; Dec. 1 and 2. Two random samples of about 50 grms. of soil were taken per half plot.

STANDARD ERRORS PER PLOT; Weeds; 22.4 cwt. per acre or 19.8%. 3rd cyst count: sampling error 32.3 or 16.6%; experimental error 59.9 or 30.7%.

Summary of Results

	TOTAL PRODUCE ; cwt. per acre ²					WEEDS ; cwt. per acre (± 11.2 . Means : ± 7.92)				
	Chlor- dinitro- ben- zene	Carbon disul- phide jelly	" Cy- mag "	" See- kay "	Sul- phate of am- monia	Chlor- dinitro- ben- zene	Carbon disul- phide jelly	" Cy- mag "	" See- kay "	Sul- phate of am- monia
None			3.4					115.2 ¹		
Single	7.6	36.2	8.2	11.7	5.5	116.8	87.0	131.5	33.4	130.8
Double	4.7	37.8	11.4	15.7	7.2	164.1	120.4	156.2	17.5	173.1
<i>Mean of single and double</i>	6.2	37.0	9.8	13.7	6.4	140.4	103.7	143.8	25.4	152.0

Standard error ; (1) ± 7.92 .

(2) No single standard error is applicable to this table.

	Third cyst count (± 30.0)				
	Chlordini- troben- zene	Carbon disulphide jelly	" Cymag "	" Seekay "	Sulphate of ammonia
None			175 ¹		
Single	211	146	192	223	230
Double	202	164	182	182	258
<i>Mean of single and double</i> (± 21.2)	206	155	187	202	244

Standard error : (1) ± 21.2 .

Conclusions

The effects of the fumigants applied in 1935 on the numbers of cysts in December 1936 were not significant.

The yields of oats were negligible except on the plots which had been fumigated with carbon disulphide jelly. In particular, the application of sulphate of ammonia in 1936 failed to produce a reasonable crop.

" Seekay " produced a large reduction in weeds.

POTATOES

Effect of Dung, ploughed in in December or applied in the bouts, of Straw and of Sulphate of Ammonia

RP—Gt. Harpenden, 1936

Total produce in lb. above, percentage ware below

1 ↑ W	— ST N DL — — — ST — DL ST —	DE — — DE ST — DE — N DL ST —	8
	206 366 165 344	317 291 352 397	
	58.8 74.8 58.1 75.2	77.0 75.5 73.0 78.1	
	— — N DE ST N DL ST N DE — —	— — — DL — — — ST — DE ST N	
	249 302 543 200	234 373 169 320	
	68.0 71.4 65.1 64.6	72.1 81.2 56.0 66.1	
— — — DL — N DE ST — DE — N	DL — N — ST N DL ST N — — N		
182 417 266 201	426 322 391 291		
60.1 80.4 69.3 62.8	81.6 77.6 75.6 71.8		
— — — DE ST N — ST — — — N	DE — — — ST N — ST — DE — N		
245 370 246 299	342 335 190 335		
72.5 78.9 72.0 75.9	80.0 78.5 60.7 73.0		
— ST N DE — N DL ST N DL ST —	DL ST N DL — — DL ST — DL — N		
238 342 458 408	458 372 277 356		
66.9 77.2 83.0 80.8	81.8 77.0 71.3 77.5		
DE ST — DE — — DL — — DL — N	DE ST — DE ST N — — N — — —		
230 314 389 336	339 369 322 254		
41 67.6 75.8 80.8 75.5	77.5 77.7 77.7 78.1	48	

SYSTEM OF REPLICATION: 4 randomised blocks of 12 plots each.
 AREA OF EACH PLOT (after rejecting edge bouts): 1/48 acre. Plots actually 1/40 acre (45.5 lks. × 54.9 lks.)

TREATMENTS: 3 × 2 × 2 factorial design.
 Dung: None, 15 tons per acre ploughed in in December (DE), or stored and applied in the bouts (DL).

Straw: None, 40 cwt. per acre (chaffed), ploughed in in December, except when applied with DL, for which straw and dung were mixed and stored (St).

Sulphate of ammonia: None, 0.4 cwt. N per acre applied in the bouts (N).

BASAL MANURING: 0.5 cwt. P₂O₅ per acre as superphosphate, and 1 cwt. K₂O per acre as sulphate of potash applied in the bouts.

CULTIVATIONS, ETC.: Applied dung and chaff: Dec. 10. Ploughed: Dec. 19-21. Springtime harrowed: April 7-10. Horse harrowed: April 15. Bouted: April 16-17. Applied stored dung and artificials: April 17-18. Potatoes planted: April 21-22. Harrowed ridges: April 24. Re-ridged: May 5. Grubbed: May 28-June 12. Earthed up: July 1. Sprayed: July 21. Lifted: Sept. 29-Oct. 1. Variety: Ally. Previous crop: Wheat.

SPECIAL NOTE: Potatoes passed through a 1½ inch riddle to determine the percentage ware.
 STANDARD ERRORS PER PLOT: Total produce: 1.00 tons per acre or 15.0%; Percentage ware: 4.87.

Summary of Results: Yields of separate treatments

		No Dung	Dung ploughed in in the bouts	
TOTAL PRODUCE: tons per acre (±0.500)				
No sulph. amm...	No straw	4.83	6.19	7.92
	Straw	4.06	5.94	7.53
Sulph. amm.	No straw	6.13	6.49	8.10
	Straw	5.81	7.19	9.77
PERCENTAGE WARE: (±2.44)				
No sulph. amm...	No straw	70.7	74.4	78.4
	Straw	61.7	72.5	76.4
Sulph. amm.	No straw	73.4	71.5	78.8
	Straw	70.4	73.5	76.4

Effect of time of application of dung

	No sulph. amm.		Sulph. amm.		Mean
	No straw	Straw	No straw	Straw	
TOTAL PRODUCE : tons per acre (± 0.707 . Mean : ± 0.354)					
Dung in the bouts minus dung ploughed in	+1.73	+1.59	+1.61	+2.58	+1.88
PERCENTAGE WARE : (± 3.45 . Mean : ± 1.72)					
Dung in the bouts minus dung ploughed in	+4.0	+3.9	+7.3	+2.9	+4.5

Effect of Straw

Response to Straw	No Dung	Dung ploughed in in the bouts		Mean
		No Dung	Dung ploughed in in the bouts	
TOTAL PRODUCE : tons per acre (± 0.707)				
No sulph. amm.	-0.77	-0.25	-0.39	
Sulph. amm.	-0.32	+0.70	+1.67	
PERCENTAGE WARE : (± 3.45)				
No sulph. amm.	-9.0	-1.9	-2.0	
Sulph. amm.	-3.0	+2.0	-2.4	

Effect of sulphate of ammonia

Response to sulphate of ammonia	No dung	Dung ploughed in in the bouts		Mean
		No dung	Dung ploughed in in the bouts	
TOTAL PRODUCE : tons per acre (± 0.707 . Mean : ± 0.408)				
No straw	+1.30	+0.30	+0.18	+0.59
Straw	+1.75	+1.25	+2.24	+1.75
PERCENTAGE WARE : (± 3.45 . Mean : ± 1.99)				
No straw	+2.7	-2.9	+0.4	+0.1
Straw	+8.7	+1.0	0.0	+3.2

Conclusions

Dung applied in the bouts gave 1.9 tons per acre more than dung ploughed in, the mean responses being 1.2 tons per acre to dung ploughed in and 3.1 tons per acre to dung applied in the bouts.

Straw gave slight but not significant decreases in yield, except in the presence of both dung and sulphate of ammonia, in which case it gave a significant increase of 1.2 tons per acre.

Sulphate of ammonia produced a significant increase of 1.2 tons per acre. The increase was somewhat greater in the presence of straw than in its absence, particularly where dung was present.

Dung produced a significant increase in percentage ware. Sulphate of ammonia gave a significant increase in the absence of dung, but no increase in its presence, the interaction between sulphate of ammonia and dung being almost significant. The depression due to straw was not significant.

SUGAR BEET

Effect of muriate of potash and agricultural salt, ploughed in in Autumn or harrowed in in Spring, and of dung.

RS—Great Harpenden, 1936

Plan and yields in lb.

Roots (dirty), tops, sugar percentage and plant number in descending order

1	— K A —	— S —	NA K S D	NA K A D	NA K S —	— A —	6
	670	674	733	728	646	600	
	584	616	760	629	500	462	
	17.42	17.68	17.86	17.71	18.49	17.57	
	503	484	500	517	505	498	
	— A —	NA — A —	— K A D	— K S D	NA K A —	NA — S —	
	677	730	755	725	701	641	
	584	634	744	704	550	534	
	17.77	18.26	17.60	17.31	17.77	17.83	
	465	478	474	523	523	510	
	— K S —	NA K A —	NA — A D	— A D	NA — A —	— K S —	
	604	697	790	788	747	648	
	480	570	682	658	480	462	
	18.58	17.22	18.20	17.51	18.32	17.80	
	537	536	538	541	531	497	
	NA K S —	NA — S —	— S D	NA — S D	— K A —	— S —	
	583	618	770	774	670	634	
	448	484	645	724	507	507	
	18.81	17.31	18.32	18.18	18.12	18.26	
	522	526	529	521	509	506	
NE							
	— K S D	— S D	NA K S —	— K A —	NA K S D	— K A D	
	703	664	689	662	747	686	
	554	556	502	508	689	666	
	18.26	17.98	18.24	18.40	18.04	17.70	
	519	515	540	528	517	509	
	— A D	NA — A D	NA — S —	— A —	NA K A D	NA — S D	
	721	799	722	649	774	730	
	590	609	550	456	638	735	
	17.76	17.92	18.20	18.09	17.82	17.92	
	518	541	526	532	543	502	
	NA — S D	— K A D	— S —	— K S —	— S D	— A D	
	804	758	666	659	751	752	
	711	649	518	501	563	700	
	18.40	18.03	17.88	18.46	17.76	17.52	
	470	521	520	520	535	483	
	NA K S D	NA K A D	NA — A —	NA K A —	NA — A D	— K S D	
	812	798	743	716	812	738	
	675	565	490	436	580	623	
	18.19	18.14	18.03	18.19	17.99	18.00	
43	563	580	580	580	560	551	48

SYSTEM OF REPLICATION : 6 randomised blocks of 8 plots each.

AREA OF EACH PLOT (after rejecting edge rows) : 0.01851 acre. Plots actually : 1/45 acre (73.3 lks. × 30.3 lks.).

TREATMENTS : 2⁴ factorial design.

Muriate of potash : None, 1.0 cwt. K₂O per acre (K).

Agricultural salt : None, 5 cwt. per acre (NA).

Minerals ploughed in, in Autumn (A). Harrowed in, in Spring (S).

Dung : None, 10 tons per acre ploughed in, in Autumn (D).

BASAL MANURING : Sulphate of ammonia at the rate of 0.6 cwt. N per acre, superphosphate at the rate of 0.5 cwt. P₂O₅ per acre.

CULTIVATIONS, ETC. : Dung applied : Dec. 9. Autumn artificials applied : Dec. 10. Ploughed : Dec. 19. Spring-tine harrowed : April 7. Rolled and harrowed : April 27. Spring artificials applied : May 1. Harrowed, rolled and drilled : May 1. Harrowed and rolled : May 2. Horse hoed : June 15. Singled : June 16 and 17. Hand hoed : Aug. 13. Lifted : Nov. 19-26. Variety : Kleinwanzleben. Previous crop : Wheat.

STANDARD ERRORS PER PLOT: Total sugar: 2.86 cwt. per acre or 5.36%. Tops: 1.09 tons per acre or 7.74%. Mean dirt tare : 0.138.

Responses to fertilisers

MEAN YIELDS: Total sugar: 53.3 cwt.; Roots (washed): 14.84 tons; Tops: 14.08 tons; Sugar percentage: 17.98; Plant number: 28.2 thousands.

	Mean response	Differential responses					
		Dung		Salt		Mur. pot.	
		Absent	Present	Absent	Present	Absent	Present
TOTAL SUGAR: cwt. per acre (± 1.17 . Means ± 0.825)							
Salt ..	+3.7	+3.1	+4.2	—	—	+4.4	+3.0
Mur. pot.	-1.1	-1.0	-1.2	-0.4	-1.8	—	—
ROOTS (washed): tons per acre							
Salt ..	+0.90	+0.81	+0.99	—	—	+1.04	+0.76
Mur. pot.	-0.32	-0.39	-0.25	-0.18	-0.46	—	—
TOPS: tons per acre (± 0.445 . Means: ± 0.314)							
Salt ..	+0.34	-0.01	+0.70	—	—	+0.72	-0.04
Mur. pot.	-0.12	-0.54	+0.29	+0.26	-0.50	—	—
SUGAR PERCENTAGE							
Salt ..	+0.13	+0.05	+0.22	—	—	+0.20	+0.06
Mur. pot.	+0.06	+0.19	-0.07	+0.13	-0.01	—	—
PLANT NUMBER: thousands per acre							
Salt ..	+0.9	+1.2	+0.6	—	—	+0.7	+1.0
Mur. pot.	+0.5	+0.6	+0.3	+0.3	+0.6	—	—

Effects of time of application of minerals

Minerals applied	None	Salt	Mur. pot.	Salt mur. pot.	No dung	Dung	Mean	Increase
TOTAL SUGAR: cwt. per acre								
Autumn	51.7 ¹	57.9 ²	52.7 ²	54.8 ²	53.1 ³	57.1 ³	55.1 ⁴	
Spring		54.2 ²	49.9 ²	53.8 ²	48.7 ³	56.6 ³	52.6 ⁴	-2.5 ³
Standard errors ..	⁽¹⁾ ± 0.825 , ⁽²⁾ ± 1.17 , ⁽³⁾ ± 0.953 , ⁽⁴⁾ ± 0.674 .							
ROOTS (washed): tons per acre								
Autumn	14.48	15.98	14.76	15.40	14.79	15.97	15.38	
Spring		15.06	13.84	14.72	13.37	15.71	14.54	-0.84
TOPS: tons per acre								
Autumn	13.78 ¹	13.97 ²	14.70 ²	13.62 ²	12.76 ³	15.44 ³	14.10 ⁴	
Spring		15.03 ²	13.36 ²	14.37 ²	11.96 ³	16.55 ³	14.25 ⁴	+0.15 ³
Standard errors ..	⁽¹⁾ ± 0.314 , ⁽²⁾ ± 0.445 , ⁽³⁾ ± 0.363 , ⁽⁴⁾ ± 0.257 .							
SUGAR PERCENTAGE								
Autumn	17.84	18.12	17.88	17.81	17.97	17.90	17.94	
Spring		17.98	18.07	18.27	18.19	18.02	18.11	+0.17
PLANT NUMBER: thousands per acre								
Autumn	27.6	29.0	27.4	29.5	28.6	28.7	28.6	
Spring		27.5	28.4	28.3	28.1	28.0	28.1	-0.5

Conclusions

Salt produced a significant increase in sugar per acre, while muriate of potash gave a decrease which was not significant.

The autumn application of salt gave somewhat higher yields than the spring application, but the difference was hardly significant.

Salt and muriate of potash had little effect on tops.

Dung applied to whole blocks gave an increase of 5.7 cwt. sugar per acre and 3.30 tons tops per acre.

SUGAR BEET

Effect of sowing date, of sulphate of ammonia and of superphosphate and muriate of potash broadcast or drilled with the seed

**RS—Great Harpenden, 1936
Plan and yields in lb.**

		Roots			Tops			Sugar		
		Sugar	Man-	Sugar	Man-	Sugar	Man-	Sugar	Man-	per
		beet	golds	beet	golds	beet	golds	beet	golds	cent.
49	2 Db —	73.5		58.0		18.49				
	3 Db N	122.7	9.0	101.0	0.5	17.42				
	3 Bs —	93.3		71.5		17.77				
	1 Bs —	135.7		81.5		18.15				
	3 — N	115.2	4.5	96.0	0.5	17.80				
	2 Bp —	81.2	3.0	54.5	0.5	17.83				
	1 Bp N	139.0		113.0		18.20				
	3 P —	85.7		66.0		18.29				
	2 — —	88.1		60.5		18.03				
	1 Da —	44.1	98.0	23.0	6.0	18.32				
	2 P N	138.1		103.0		17.74				
	3 Da —	79.0	12.0	67.5	1.0	18.06				
	2 Bs N	155.7		144.0		17.31				
	1 P —	132.5	6.0	94.5	0.5	17.14				
	1 Db N	161.8	58.0	126.5	3.5	17.34				
	3 Bp N	127.9	5.5	126.5	0.5	16.79				
	2 Da N	156.4	11.5	147.0	1.0	17.28				
	1 — N	153.5		108.0		17.31				
	3 Da —	101.0		108.5		17.05				
	3 Bp —	88.7		93.0		17.14				
	2 Bs —	97.2		85.0		17.16				
	1 Da —	51.2	118.0	38.5	14.0	16.99				
	3 — N	93.8		103.5		17.16				
	3 Db N	125.6	11.0	126.0	0.5	17.11				
	1 P —	130.0		101.0		17.34				
	1 Bs N	166.3		130.5		17.37				
	3 P —	91.7	8.0	81.5	0.5	17.28				
	1 Db N	128.0	12.0	89.0	0.5	17.60				
	1 Bp —	98.7		68.5		17.34				
	2 — —	99.5		76.5		17.11				
	1 — N	182.4		146.0		17.25				
	2 Db —	123.8	4.5	113.5	0.5	16.76				
	2 Da N	116.9	7.5	112.0	0.5	17.05				
	3 Bs N	149.0		135.5		16.47				
	2 Bp N	140.0		102.0		16.96				
84	2 P N	146.1		67.5		16.30				
	2 — N	101.5		54.5		18.61				
	1 P N	142.0		75.0		18.91				
	1 Bp —	95.4	4.0	49.5	0.5	18.75				
	1 Bs N	159.9	9.5	91.5	0.5	18.52				
	3 P N	149.1		113.0		18.46				
	2 Db N	104.8		69.0		18.61				
	2 Bp N	121.0		83.5		18.64				
	2 P —	108.9		70.0		18.46				
	3 Db —	88.5	6.5	58.0	0.5	18.87				
	1 Da N	112.8	26.0	58.0	1.0	18.61				
	2 Da —	92.2	12.5	66.0	0.5	18.44				
	3 Bs N	124.3	7.0	99.0	0.5	17.94				
	3 — —	109.0		85.5		17.12				
	3 Bp —	94.6		84.0		17.22				
	3 Da N	122.8	15.5	121.0	2.5	17.34				
	2 Bs —	138.0		112.5		17.57				
	1 — —	101.7		63.0		17.05				
	1 Db —	93.0	7.0	61.5	1.5	16.88				
	2 Bp —	119.6		87.5		17.57				
	2 P —	117.9		83.0		18.32				
	2 Bs N	157.2		115.5		17.68				
	3 Db —	92.0	7.0	78.5	1.5	17.74				
	1 — —	119.5		66.0		17.48				
	1 Bp N	168.2		115.5		17.74				
	3 Bs —	90.0		79.0		17.94				
	3 Da N	143.9	4.0	140.0	0.5	17.40				
	1 Bs —	129.7		81.5		17.77				
	2 Da —	91.0	4.0	73.5	0.5	18.03				
	2 — N	141.0	4.5	104.0	0.5	17.58				
	1 Da N	136.0	54.5	84.5	3.0	17.80				
	3 — —	87.9		69.5		17.60				
	1 P N	176.3		110.5		17.97				
	3 P N	133.8	6.0	120.0	1.0	17.63				
	3 Bp N	120.4	8.5	114.5	1.5	17.37				
	2 Db N	134.6	8.0	119.0	1.0	17.40				
	1 Db —	81.1	7.5	52.0	1.0	17.48				
										120

SYSTEM OF REPLICATION ; 4 randomised blocks of 18 plots each. Certain interactions partially confounded with block differences.

AREA OF EACH PLOT (after rejecting edge-rows) ; 1/190 acre. Plots actually 1/95 (94.8 lks. \times 11.1 lks.).

TREATMENTS ; 6 \times 3 \times 2 factorial design.

Sowing dates ; April 23 (1), May 8 (2), May 26 (3).

Minerals ; None (—), ploughed in (P), broadcast immediately after ploughing (Bp), broadcast at sowing (Bs), drilled below seed (Db), drilled above seed (Da), at the rate of 0.5 cwt. P_2O_5 per acre as superphosphate and 1.0 cwt. K_2O per acre as muriate of potash.

Sulphate of ammonia ; None (—), 0.6 cwt. N per acre drilled where minerals are drilled, otherwise broadcast (N).

BASAL MANURING ; Nil.

CULTIVATIONS, ETC. ; Minerals applied (P) ; March 21 . Ploughed ; March 27-April 1. Minerals applied (Bp) ; April 3. Spring-tine harrowed ; April 7 and 8. Rolled ; April 8. Harrowed ; April 20 and 23. Rolled ; April 23. Minerals applied ; (Bs, Db, Da) ; April 23. Harrowed and rolled ; April 24. Rolled ; May 8. Minerals applied (Bs, Db, Da) ; May 8. Harrowed ; May 8. Harrowed and rolled ; May 26. Minerals applied (Bs, Db, Da) ; May 26. Horse hoed 1st sowing ; May 28. Singled 1st sowing ; June 11. Horse hoed 2nd sowing ; June 12. Singled 2nd sowing ; June 17. Horse hoed 3rd sowing ; June 23. Singled 3rd sowing and hand hoed 1st and 2nd sowings ; July 4. Horse and hand hoed all sowings ; Aug. 8 and 12. Lifted ; Nov. 27-Dec. 3. Variety ; Kleinwanzleben E. Previous crop ; Wheat.

SPECIAL NOTE ; The seed drill used had not been properly cleaned before the experiment and unfortunately contained mangolds seeds, which grew along with the sugar beet on certain plots. The results have been combined by assuming that a sugar beet root weighs one half a mangolds root and that sugar beet tops weigh 2.5 times mangolds tops. The individual weights of sugar beet and mangolds are, however, given in all cases. It will be noticed that the mistake severely affected a few treatments only.

STANDARD ERROR PER PLOT : Roots (Sugar beet + $\frac{1}{2}$ Mangolds) : 1.39 tons per acre or 13.4% ;
Tops (Sugar beet + $\frac{5}{2}$ Mangolds) : 1.52 tons per acre or 19.4%.

Conclusions

Later sowing produced a significant decrease in the yield of roots and a significant increase in the yield of tops. The sugar percentage was also slightly lower with the later sowings, so that the yield of sugar per acre dropped by 4.7 cwt. per acre from the first to the second sowing and by a further 2.6 cwt. per acre from the second sowing to the third.

Sulphate of ammonia produced large increases in the yields of roots and tops. Sulphate of ammonia had little effect on the average on sugar percentage and gave an average increase in sugar per acre of 11.6 cwt. or 32 per cent. of the mean yield.

The average response to minerals was not significant in roots, tops or sugar percentage.

Main effects and interactions of sowings with fertilisers

	Sugar beet			Mangolds			Sugar beet + $\frac{1}{2}$ Mangolds			Mean Increase
	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	S ₁	S ₂	S ₃	
	ROOTS; tons per acre									
N ₀	8.57	8.70	7.79	1.70	0.17	0.24	9.42 ¹	8.78	7.91	8.70 ²
N ₁	12.91	11.40	10.80	1.13	0.22	0.50	13.48	11.51	11.05	12.01 + 3.31 ³
O	11.81	9.12	8.61	0.00	0.10	0.10	11.81 ⁴	9.17	8.66	9.88 ¹
P	12.32	10.84	9.76	0.13	0.00	0.30	12.38	10.84	9.91	11.04 + 1.16 ⁵
Bp	10.63	9.79	9.15	0.08	0.06	0.30	10.67	9.82	9.30	9.93 + 0.05
Bs	12.55	11.62	9.68	0.20	0.00	0.15	12.65	11.62	9.76	11.34 + 1.46
Db	9.84	9.26	9.09	1.79	0.27	0.71	10.74	9.40	9.44	9.86 - 0.02
Da	7.30	9.68	9.47	6.29	0.75	0.67	10.44	10.06	9.80	10.10 + 0.22
Mean Incr'se							11.45 ⁶	10.15	9.48	10.36
								-1.30 ¹	-1.97	
	TOPS; tons per acre									
N ₀	5.52	6.65	6.66	0.17	0.01	0.02	5.94 ⁷	6.67	6.71	6.44 ⁸
N ₁	8.82	8.63	9.87	0.06	0.02	0.06	8.97	8.68	10.02	9.22 + 2.78 ⁹
O	8.12	6.27	7.52	0.00	0.01	0.01	8.12 ¹⁰	6.29	7.54	7.32 ⁷
P	8.08	6.86	8.07	0.01	0.00	0.03	8.10	6.86	8.15	7.70 + 0.38 ¹¹
Bp	7.35	6.94	8.86	0.01	0.01	0.04	7.37	6.96	8.96	7.76 + 0.44
Bs	8.16	9.69	8.16	0.01	0.00	0.01	8.18	9.69	8.18	8.68 + 1.36
Db	6.98	7.62	7.71	0.14	0.03	0.06	7.33	7.70	7.86	7.63 + 0.31
Da	4.33	8.45	9.27	0.51	0.05	0.08	5.61	8.57	9.47	7.88 + 0.56
Mean Incr'se							7.45 ¹²	7.68	8.36	7.83
								+0.23 ⁷	+0.91	
	SUGAR PERCENTAGE					TOTAL SUGAR; cwt. per acre				
	S ₁	S ₂	S ₃	Mean	Increase	S ₁	S ₂	S ₃	Mean	Increase
N ₀	17.56	17.81	17.67	17.68		33.1	31.3	28.0	30.8	
N ₁	17.88	17.60	17.41	17.63	-0.05	48.2	40.5	38.5	42.4	+11.6
O	17.27	17.83	17.42	17.51		40.8	32.7	30.2	34.6	
P	17.84	17.70	17.92	17.82	+0.31	44.2	38.4	35.5	39.4	+4.8
Bp	18.01	17.75	17.13	17.63	+0.12	38.4	34.9	31.9	35.1	+0.5
Bs	17.95	17.43	17.53	17.64	+0.13	45.4	40.5	34.2	40.0	+5.4
Db	17.32	17.82	17.78	17.64	+0.13	37.2	33.5	33.6	34.8	+0.2
Da	17.93	17.70	17.46	17.70	+0.19	37.4	35.6	34.2	35.7	+1.1
Mean Increase	17.72	17.70	17.54	17.66		40.6	35.9	33.3	36.6	
		-0.02	-0.18				-4.7	-7.3		

Standard errors: (1) ± 0.401, (2) ± 0.232, (3) ± 0.328, (4) ± 0.695, (5) ± 0.567, (6) ± 0.284, (7) ± 0.439, (8) ± 0.253, (9) ± 0.358, (10) ± 0.760, (11) ± 0.621, (12) ± 0.310.

MANGOLDS

Effect of sulphate of ammonia, superphosphate, muriate of potash, agricultural salt and dung

RM—GREAT HARPENDEN—1936

Plan and yields in lb., roots above, tops centre, plant number below

1	- PK - D	---	NA D	S - K	---	SPKNAD	---	---	D - PKNA	D	---	K	---	S	---	NA D	8				
	844		1104	1156		1508			1248		1100		784			1376					
	126		157	139		196			120		136		95			152					
	444		461	471		446			484		448		479			478					
NE	SP - NA	---	KN _A	S	---	D	---	P	---	SPK - D	S - KN _A	---	SP	---	P - NA						
	1312		1000	1176				888		1356		1376		1008		964					
	168		150	146				123		158		171		122		130					
↑	466		472	492				504		478		461		493		472					
	---	K - D	SP	---	D	P - NA	---	PKNA	---	S - K - D	SPKNA	---	KN _A D	---	SP - NA						
	896		1284			996		860		1328		1292		1008		1324					
	148		174			146		132		180		167		138		176					
	460		490			469		451		487		458		462		471					
25	S	---	NA	---	SPK	---	---	S - KN _A	D	---	P	---	D	---	PK	---	---	NA	---	S	---
	1184				984			740		1468				1008		692		780			1108
	156				148			110		211				144		110		126			124
	439				474			507		448				447		470		492			487

SYSTEM OF REPLICATION : 4 randomised blocks of 8 plots each. Certain high order interactions are partially confounded with block differences.

AREA OF EACH PLOT (after rejecting edge rows) : 0.02322 acre. Plots actually, 1/45 acre (48.8 lks. × 45.5 lks.)

TREATMENTS : 2⁵ factorial design.

- Sulphate of ammonia : None, 0.6 cwt. N per acre (S).
- Superphosphate : None, 0.5 cwt. P₂O₅ per acre (P).
- Muriate of Potash : None, 1.0 cwt. K₂O per acre (K).
- Agricultural Salt : None, 5 cwt. per acre (NA).
- Dung : None, 10 tons per acre (D).

BASAL MANURING : Nil.

CULTIVATIONS, ETC. : Dung applied : Dec. 11. Ploughed : Dec. 17-19. Spring-tine harrowed : April 7. Artificials applied : May 5. Harrowed : May 5. Rolled : May 5. Drilled : May 5. Horse hoed : June 12, Aug. 12 and 13. Singled : July 2 and 3. Lifted : Nov. 18 and 19. Variety : Yellow Globe. Previous crop : Wheat.

STANDARD ERROR PER PLOT : Roots : 1.91 tons per acre or 7.50%

Responses to fertilisers

MEAN YIELD, Roots : 25.50 tons ; Tops, 3.40 tons ; Plant number : 21.2 thousands.

	Mean response	Differential responses									
		Sulphate of ammonia		Dung		Salt		Muriate of potash		Superphosphate	
		Absent	Present	Absent	Present	Absent	Present	Absent	Present	Absent	Present
ROOTS : tons per acre (± 0.955 , Means : ± 0.675)											
Sulphate of ammonia	+7.73	—	—	+7.87	+7.59	+6.68	+8.79	+5.93	+9.53	+7.58	+7.88
Dung	+4.20	+4.34	+4.06	—	—	+5.17	+3.24	+4.45	+3.96	+4.28	+4.12
Salt	+3.12	+2.07	+4.18	+4.09	+2.16	—	—	+1.68	+4.56	+2.50	+3.75
Muriate of potash ..	+0.22	-1.58	+2.02	+0.46	-0.02	-1.22	+1.66	—	—	+0.87	-0.43
Superphosphate ..	-0.45	-0.60	-0.30	-0.37	-0.53	-1.08	+0.17	+0.20	-1.10	—	—
TOPS : tons per acre.											
Sulphate of ammonia	+0.72	—	—	+0.64	+0.80	+0.62	+0.83	+0.47	+0.98	+0.69	+0.76
Dung	+0.50	+0.42	+0.58	—	—	+0.66	+0.33	+0.46	+0.54	+0.53	+0.46
Salt	+0.50	+0.40	+0.60	+0.66	+0.33	—	—	+0.43	+0.57	+0.57	+0.43
Muriate of potash ..	+0.19	-0.06	+0.44	+0.14	+0.23	+0.12	+0.26	—	—	+0.41	-0.04
Superphosphate ..	+0.05	+0.01	+0.08	+0.08	+0.02	+0.12	-0.02	+0.27	-0.18	—	—
PLANT NUMBER : thousands per acre											
Sulphate of ammonia	0.0	—	—	-0.6	+0.6	+0.4	-0.3	-0.1	+0.2	-0.3	+0.4
Dung	-0.4	-1.0	+0.2	—	—	-0.6	-0.1	-0.4	-0.3	-0.2	-0.5
Salt	-0.8	-0.4	-1.1	-1.0	-0.5	—	—	-0.9	-0.7	-0.9	-0.7
Muriate of potash ..	-0.7	-0.8	-0.5	-0.7	-0.5	-0.8	-0.6	—	—	-0.6	-0.8
Superphosphate ..	-0.3	-0.6	+0.1	-0.1	-0.4	-0.4	-0.2	-0.2	-0.4	—	—

Conclusions

Sulphate of ammonia, dung and salt produced significant increases in the yield of roots. The average response in roots to muriate of potash was small and not significant, but the responses to sulphate of ammonia and salt were increased by the presence of muriate of potash, the increase being significant for sulphate of ammonia and almost so for salt. The results for tops were similar to those for roots. Superphosphate had little effect on roots or tops.

BEANS

Effect of dung, nitro-chalk, superphosphate and muriate of potash
RE—Gt. Harpenden, 1936
Plan and yields in lb., grain above, straw below

28	P	K	D	NPK	NPK	D	P	DNK	35
	45.1	55.0	53.1	36.2	43.1	42.2	38.8	34.2	
	78.4	83.0	89.9	87.0	95.2	96.3	81.0	101.6	
N	DNK	DNP	DPK	N	N	DNP	K	DPK	
	41.0	48.4	54.8	41.7	46.9	52.2	50.2	44.5	
↑	87.8	90.8	81.2	79.0	78.1	99.8	91.0	98.2	
52	DP	NK	DK	PK	NK	DP	O	NP	59
	50.1	44.5	42.6	51.4	42.9	52.4	57.0	39.2	
	86.9	101.2	84.4	80.1	89.4	98.8	73.0	96.0	
	DNPK	O	DN	NP	PK	DK	DNPK	DN	
	44.1	57.5	41.3	49.7	55.8	51.9	54.0	42.5	
	94.2	87.5	102.2	87.3	114.0	110.8	116.8	110.0	

SYSTEM OF REPLICATION : 4 randomised blocks of 8 plots each. Certain interactions confounded with block differences.

AREA OF EACH PLOT : 1/40 acre. (54.5 lks. by 45.9 lks.)

TREATMENTS : 2⁴ factorial design.

Dung : none, 10 tons per acre (D).

Nitrochalk : none, 0.4 cwt. N per acre (N).

Superphosphate : none, 0.6 cwt. P₂O₅ per acre (P).

Muriate of potash : none, 1.0 cwt. K₂O per acre (K).

SPECIAL NOTE : The two right hand blocks failed and were redrilled in the spring with spring beans.

CULTIVATIONS, ETC. : Plots, 32-35, 40-43, 48-51, 56-59. Applied dung and artificials : Oct. 11. Ploughed : Oct. 14-16. Harrowed : Oct. 18. Drilled : Oct. 21. Crop failed. Springtine harrowed : March 16. Harrowed, redrilled and harrowed in : March 19. Horse hoed : May 7 and 29.

Plots, 28-31, 36-39, 44-47, 52-55. Applied dung and artificials : Oct. 11. Ploughed : Oct. 14-16. Harrowed : Oct. 18. Drilled : Oct. 21. Harrowed : March 24. Hand hoed : April 24-27 and June 9. Horse hoed : May 29. All plots harvested : Aug. 24. Previous crop : Wheat.

STANDARD ERROR PER PLOT : Grain : 1.72 cwt. per acre or 10.3%.

Responses to fertilisers

Mean yields : GRAIN, 16.8 cwt ; STRAW, 32.9 cwt.

	Mean response	Differential responses							
		Dung		Nitro-chalk		Superphosphate		Mur. pot.	
		Absent	Present	Absent	Present	Absent	Present	Absent	Present
GRAIN : cwt. per acre (± 0.860 . Means : ± 0.608).									
Dung	-0.1	—	—	-0.8	+0.6	-2.1	+1.8	+0.3	-0.5
Nitro-chalk	-2.2	-3.0	-1.5	—	—	-3.3	-1.2	-1.5	-3.0
Superphosphate	+0.3	-1.6	+2.3	-0.7	+1.4	—	—	-0.3	+1.0
Mur. pot.	-0.3	+0.1	-0.7	+0.4	-1.0	-0.9	+0.4	—	—
STRAW : cwt. per acre									
Dung	+3.3	—	—	+2.6	+4.0	+4.5	+2.1	+5.1	+1.5
Nitro-chalk	+1.8	+1.0	+2.6	—	—	+1.5	+2.1	+2.2	+1.4
Superphosphate	+0.5	+1.6	-0.7	+0.2	+0.8	—	—	+0.2	+0.8
Mur. pot.	+1.8	+3.6	0.0	+2.2	+1.4	+1.5	+2.1	—	—

Conclusions

The average effects of dung, superphosphate and muriate of potash on the yield of grain were negligible, while nitrochalk produced a significant depression in grain. The interaction between the effects of dung and superphosphate was statistically significant, but in view of absence of any apparent average effects of either dung or superphosphate no weight can be attached to it.

Dung produced a significant increase in the straw yields.

KALE

Effect of sulphate of ammonia, poultry manure, soot and rape dust

RK—FOSTER'S, 1936 (3rd year)

Plan and yields in lb.

1	R ₁ 800	N ₀ 650	M ₀ 629	M ₁ 792	M ₂ 724	R ₁ 683	R ₂ 767	N ₁ 836	8
	S ₀ 741	S ₁ 822	N ₁ 883	M ₂ 875	S ₂ 796	M ₁ 622	M ₀ 592	N ₀ 715	
SW ↑	R ₂ 969	S ₂ 927	R ₀ 742	N ₂ 975	R ₀ 764	S ₀ 729	N ₂ 782	S ₁ 848	
	R ₂ 958	N ₀ 798	M ₁ 763	S ₁ 814	R ₂ 874	R ₀ 691	R ₁ 728	M ₁ 856	
	N ₂ 868	M ₀ 746	M ₂ 876	S ₂ 918	M ₂ 785	N ₀ 776	N ₁ 828	S ₀ 756*	
41	R ₁ 720	S ₀ 792	R ₀ 737	N ₁ 776	S ₂ 784	N ₂ 930	S ₁ 830*	M ₀ 657*	48

* Estimated.

SYSTEM OF REPLICATION : 4 randomised blocks of 12 plots each.

AREA OF EACH PLOT : 0.028926 acre (10 yds. × 14 yds.)

TREATMENTS, 1936 : No nitrogen (0), sulphate of ammonia (N) half applied in seed bed and remainder as a top dressing, poultry manure (M), soot (S) and rape dust (R), applied in seed-bed at the rate of 0.4 cwt. N per acre (1) or 0.8 cwt. N per acre (2). Plots receiving treatment O in 1936 had treatment 2 in 1935 and *vice versa*. Plots receiving treatment 1 had this in both years. For N₀, S₀, M₀, and R, (see plan) the fertilizer symbols refer to the 1935 treatment.

BASAL MANURING : All plots were made up to 1.0 cwt. P₂O₅ per acre and 1.0 cwt. K₂O per acre, using superphosphate and muriate of potash (an allowance being made for the P₂O₅ and K₂O contained in the organic manures).

CULTIVATIONS, ETC. Ploughed : March 7, 23-26. Springtine harrowed : April 15. Applied manures (sulphate of ammonia at half-rate) : May 2. Harrowed and rolled, before and after seed sown : May 4. Seed resown : May 26. Harrowed and rolled : May 26. Dusted with Derris powder : June 12 and 17. Hand-hoed : July 9. Horse-hoed : July 22. Applied second half of sulphate of ammonia : July 27. Harvested : Dec. 22-Feb. 2. Variety : Thousand head. Previous crop : Brussels Sprouts. (See 1935 Report, p.191).

SPECIAL NOTE : Kale harvested at weekly intervals as food for stock.

STANDARD ERROR PER PLOT : 0.925 tons per acre or 7.59%.

Summary of Results : tons per acre (±0.462)

Nitrogen, cwt. per acre			Sulph. amm.	Poultry manure	Soot	Rape dust	Mean (±0.231)
1934	1935	1936					
0.0	0.8	0.0	11.34	10.13	11.64	11.32	11.11
0.4	0.4	0.4	12.83	11.71	12.79	11.31	12.16
0.8	0.0	0.8	13.71	12.58	13.21	13.76	13.32
Mean (±0.267)			12.63	11.47	12.55	12.13	12.20

Conclusions

There was a significant response to the 1936 application of nitrogen. Poultry manure gave lower yields than soot or sulphate of ammonia with all three types of dressing, the average differences being significant. Rape dust occupied an intermediate position.