


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Report for 1937

[Full Table of Content](#)



Abbreviated List of the Field Experiments

Rothamsted Research

Rothamsted Research (1938) *Abbreviated List of the Field Experiments* ; Report For 1937, pp 223 - 224 - DOI: <https://doi.org/10.23637/ERADOC-1-69>

ABBREVIATED LIST OF THE FIELD EXPERIMENTS

	<i>Page</i>
Chemical analysis of manures used in replicated experiments	121-122
Average wheat yields of various countries	122
Conversion table	122
Meteorological records	123
 Classical Experiments	
Rotation—Agdell	124
Wheat after fallow—Hoosfield	125
Mangolds—Barnfield	126
Hay—Park grass	127
Botanical Composition—1937 (1st crop)	128
Botanical Composition—1934 (1st crop)	129
Detailed botanical analysis 1937 (1st crop)	130-131
Wheat—Broadbalk	132-133
Barley—Hoosfield	134
 Modern Long Term Experiments	
Four Course Rotation—Residual value of humic and phosphatic fertilisers ..	135-137
Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P ₂ O ₅ and K ₂ O	138-142
Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops	143-145
Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer	146-150
New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw ..	151-152
 Short Term Experiments—Rothamsted	
Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia	153-154
Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash	155-156
Sugar Beet—Four methods of applying salt and muriate of potash; dung	157-159
Mangolds—Sulphate of ammonia, superphosphate, muriate of potash, salt and dung	160-161
Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows	162-163
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects	164
 Short Term Experiments—Woburn	
Sugar Beet—Four methods of applying salt and muriate of potash; dung	165-166
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects	167
Pyrethrum—Lime, fish manure and artificials, applied every year	168-169
 Summaries of Groups of Experiments	
Experiments on poultry manure and sulphate of ammonia	170-174
Sugar beet fertiliser experiments, factory series	175-196

Experiments at Outside Centres		<i>Page</i>
Barley.	Tunstall, Suffolk—Super. or slag, limestone or dolomite	197
	Wye, Kent—S/A or nitro-chalk, super.	197-198
Clover.	Tunstall, Suffolk—Residuals of chalk (O, 1, 2, 3, 4)	198
Potatoes.	Thorney, Cambs.—N (0, 1, 2), P (0, 1, 2), K (0, 1, 2)	199
	March, Cambs.—N (0, 1, 2), P (0, 1, 2), K (0, 1, 2)	200
	Littleport, Cambs.—N (0, 1, 2), P (0, 1, 2), K (0, 1, 2)	201
	Tunstall, Suffolk—P, K, magnesium sulphate, dung	202
Sugar Beet.	Kidderminster, Worcester—S/A or N/S, super. or slag, K (0, 1, 2) ..	202-204

Experiments carried out by Local Workers

Hay.	Harlow, Essex—(0, 1, 2) levels of slag, super, mineral phosphate ; K (0, 1)	205
	Burford, Oxford—Slag (0, 1, 3)	205
	Bakewell, Derby—N, P, K	206
Potatoes.	Bakewell, Derby—Artificial and compost, direct and residual effects	206
	Barnet, Herts—Residuals of K, chalk, phosphates	207
	Loughborough, Leicester—Mixed artificials (0, 1, 2, 3)	207
	Ingham, Lincs.—Mixed artificials (0, 1, 2, 3, 4)	208
	Welton, Lincs.—Mixed artificials (0, 1, 2, 3, 4)	208
Sugar Beet.	Market Rasen, Lincs.—Methods of applying minerals ; S/A, dung ..	209-210
	Wragby, Lincs.—Methods of applying minerals ; S/A, dung ..	210-211
	East Kirkby, Lincs.—Methods of applying minerals ; S/A, depth of ploughing	212-213
	Blyborough, Lincs.—Methods of applying minerals ; S/A, depth of ploughing	213-215
	Blyton, Lincs.—Borax (0, 1, 2), artificials	215
	Digby Fen, Lincs.—Manganese sulphate (0, 1, 3). N/S	216
	Brough, Notts—Mixed artificials (0, 1, 2, 3)	217
	Newton-on-Trent, Lincs.—Mixed artificials (0, 1, 2, 3)	217
	North Muskham, Notts—Mixed artificials (0, 1, 2, 3)	218
	Crowle, Lincs.—Mixed artificials (0, 1, 2, 3), N, time of lifting ..	218-219
	Holton-le-Moor, Lincs.—Methods of singling	220
Thornton, Lincs.—Methods of singling	220	
Kale.	Kneeton, Notts—Cyanamide, nitro-chalk, S/A ; super or slag ..	221
	Loughborough, Leicester—N, thinning	222

Note.—N denotes sulphate of ammonia or nitrate of soda, P denotes superphosphate, and K denotes any potash fertiliser.

ERRATA

1932 Report

p. 203. Salt was applied at the rate of 0.85 cwt. Cl. per acre.

1936 Report

p. 204. The third sentence from the foot of the page should read "The response to each was somewhat greater at the lower level of the other than at the higher level, but the difference was not significant."

p. 223. In harvesting the kale, two guard rows were omitted from each plot. All yields and their standard errors should be multiplied by 18/16 or 1.125.

p. 233, line 4 in the table. For "Blakewell" read "Bakewell."

p. 272, line 3. For "3 x 2 factorial design" read "3² factorial design."