Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Report for 1937

Redmand Experiment Station
Hayendan
State Generalization was
REPORT
for
1937
For Annual on a factor
state Generalization of State Control of S

Full Table of Content

## **Abbreviated List of the Field Experiments**

## **Rothamsted Research**

Rothamsted Research (1938) *Abbreviated List of the Field Experiments*; Report For 1937, pp 223 - 224 - **DOI:** https://doi.org/10.23637/ERADOC-1-69

223

## ABBREVIATED LIST OF THE FIELD EXPERIMENTS

Pyrethrum—Lime, fish manure and artificials, applied every year								Page
Conversion table Meterological records  Classical Experiments  Rotation—Agdell Rotation—Agdell Rotation—Agdell Rotation—Agdell Rotation—Boosfield Rotation—Boosfield Rotation—Composition—1937 (1st crop) Rotanical Composition—1937 (1st crop) Rotanical Composition—1937 (1st crop) Rotanical Composition—1937 (1st crop) Rotatiled botanical analysis 1937 (1st crop) Rotatiled Rotat	Chemical analysis of manures used in replic	cated exper	iments					121-122
Classical Experiments  Rotation—Agdell								122
Classical Experiments  Rotation—Agdell	Conversion table							
Rotation—Agdell	Meterological records	•	••		••	••	••	123
Wheat after fallow—Hoosfield Mangolds—Barnfield Mangolds—Barnfield May—Park grass Botanical Composition—1937 (1st crop) Botanical Composition—1934 (1st crop) Detailed botanical analysis 1937 (1st cr	Classical Experiments							
Mangolds—Barnfield Hay—Park grass Botanical Composition—1937 (1st crop)	Rotation—Agdell							124
Mangolds—Barnfield 129 Botanical Composition—1937 (1st crop) 128 Botanical Composition—1934 (1st crop) 129 Detailed botanical analysis 1937 (1st crop) 130-131 Barley—Hoosfield 132-133 Barley—Hoosf	Wheat after fallow—Hoosfield							
Botanical Composition—1937 (1st crop). 128 Botanical Composition—1937 (1st crop). 129 Detailed Dotanical analysis 1937 (1st crop) 130-131 Wheat—Broadbalk 132-133 Barley—Hoosfield . 132-133 Barley—Hoosfield . 132-133  Modern Long Term Experiments  Four Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops . 143-145 Three Course Rotation—Utilisetion of straw and green manuring. Effect of ploughing in straw and of winter green manure crops . 143-145  New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw . 151-152  Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia . 153-154 Sugar Beet—Four methods of applying salt and muriate of potash; dung . 155-156 Sugar Beet—Four methods of applying salt and muriate of potash and spacing of the rows	Mangolds—Barnfield						• •	
Botanical Composition—1934 (1st crop). 129 Detailed botanical analysis 1937 (1st crop) 130-131 Wheat—Broadbalk 132-133 Barley—Hoosfield	Hay—Park grass							
Detailed botanical analysis 1937 (1st crop) 130–131 Wheat—Broadbalk 132–133 Barley—Hoosfield								
Modern Long Term Experiments  Four Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manure. Effect of ploughing in straw and of winter green manure crops Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw.  Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia. Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash. Sigar Beet—Four methods of applying salt and muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  165–166  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  168–169  Summaries of Groups of Experiments  Experiments on poultry manure and sulphate of ammonia.								and the second s
Modern Long Term Experiments  Four Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw.  Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash Sangar Beet—Four methods of applying salt and muriate of potash; dung Mangolds—Sulphate of ammonia, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and dung Beans—Dung have prephosphate, muriate of potash, salt and Beans—Beans—Beans—Beans—Beans—Beans—Beans—Beans								
Four Course Rotation—Residual value of humic and phosphatic fertilisers Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw.  Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash dung.  Mangolds—Sulphate of ammonia, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows.  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  162-163  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  164-164  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung.  165-166  Summaries of Groups of Experiments  Experiments on poultry manure and sulphate of ammonia.  170-174								
Four Course Rotation—Residual value of humic and phosphatic fertilisers  Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manuring. Effect of plough ing in straw and of winter green manure crops  Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer  New Green Manuring Experiment, Woburn—Effects on kale of clover and rye- grass as leys, of mustard and tares as green manures and of dung, N and straw  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia  Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash Sugar Beet—Four methods of applying salt and muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Sympthylogen and straw at two times of potash; dung  165–166  167  168–169  Summaries of Groups of Experiments  Experiments on poultry manure and sulphate of ammonia	During Troubles of the Control of th							
Four Course Rotation—Residual value of humic and phosphatic fertilisers  Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops  Three Course Rotation—Effects of various types and depths of cultivation. The use of cyanamide as a weed-killer  New Green Manuring Experiment, Woburn—Effects on kale of clover and ryegrass as leys, of mustard and tares as green manures and of dung, N and straw  Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia  Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash  Mangolds—Sulphate of ammonia, superphosphate, muriate of potash; alt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung  Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects  Spyrethrum—Lime, fish manure and artificials, applied every year  165–166  167–168  Summaries of Groups of Experiments  Experiments on poultry manure and sulphate of ammonia								
Six Course Rotation, Rothamsted and Woburn—Seasonal effects of N, P <sub>2</sub> O <sub>5</sub> and K <sub>2</sub> O Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops	Modern Long Term Experiments							
Three Course Rotation—Utilisation of straw and green manuring. Effect of ploughing in straw and of winter green manure crops	Six Course Rotation, Rothamsted and Wo	oburn—Sea	sonal ef	fects of	N, P,	O <sub>5</sub> and	K <sub>2</sub> O	
use of cyanamide as a weed-killer	Three Course Rotation—Utilisation of straw and green manuring. Effect of plough-							
Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia	Three Course Rotation—Effects of vario	ous types a	and dep	ths of	cultiva	tion.	The	140 150
Short Term Experiments—Rothamsted  Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia	use of cyanamide as a weed-killer							
Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia	grass as leys, of mustard and tares as	green man	ures and	d of du	ng, Na	nd stra	w	151-152
Spring Oats—Residual effects of dung and straw at two times of application and of sulphate of ammonia								
Sulphate of ammonia Potatoes—Dung and straw at two times of application; sulphate of ammonia, superphosphate and sulphate of potash Sugar Beet—Four methods of applying salt and muriate of potash; dung								
Sugar Beet—Four methods of applying salt and muriate of potash; dung	Spring Oats—Residual effects of dung ar	nd straw at	two tin	nes of a	pplica	tion an	d of	
Sugar Beet—Four methods of applying salt and muriate of potash; dung	sulphate of ammonia							
Sugar Beet—Four methods of applying salt and muriate of potash; dung	Potatoes—Dung and straw at two tim	ies of appli	ication;	sulpr	nate of	ammo	nia,	155 150
Mangolds—Sulphate of ammonia, superphosphate, muriate of potash, salt and dung Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows	superphosphate and sulphate of potash							
Beans—Dung, nitro-chalk, superphosphate, muriate of potash and spacing of the rows	Sugar Beet—Four methods of applying salt and muriate of potash; dung							
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects	Page Dung nitro-chalk superphosphate muriate of notash and spacing of the							
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects	beaus—Dung, mitto-chain, superphosphato, marketo of potential and spacing of							
Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung	Kale, Sulphate of ammonia poultry manure, soot and rape dust. Direct, cumulative							
Short Term Experiments—Woburn  Sugar Beet—Four methods of applying salt and muriate of potash; dung								164
Sugar Beet—Four methods of applying salt and muriate of potash; dung	and rosadas strong							
Sugar Beet—Four methods of applying salt and muriate of potash; dung								
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects								
Kale—Sulphate of ammonia, poultry manure, soot and rape dust. Direct, cumulative and residual effects	Sugar Beet-Four methods of applying	salt and m	uriate o	f potas	h; dun	g		165-166
Pyrethrum—Lime, fish manure and artificials, applied every year	Kale—Sulphate of ammonia, poultry m	anure, soot	and ra	pe dus	t. Di	ect, cu	mu-	
Summaries of Groups of Experiments  Experiments on poultry manure and sulphate of ammonia	lative and residual effects							
Experiments on poultry manure and sulphate of ammonia 170-174	Pyrethrum—Lime, fish manure and arti	iticials, app	lied eve	ry yea	г			108-109
Experiments on poultry manure and sulphate of ammonia 170-174								
Experiments on poultry manure and sulphate of ammonia 170-174	Summaries of Groups of Experimen	ts						
Experiments on pourtry manare and surplus								170 174
	Experiments on poultry manure and sul Sugar beet fertiliser experiments, factor	y series						

#### 224

Experiments	at Outside Centres	Page
Barley.	Tunstall, Suffolk—Super. or slag, limestone or dolomite	197
	Wye, Kent—S/A or nitro-chalk, super.	197-198
Clover.	Tunstall, Suffolk—Residuals of chalk (O. 1, 2, 3, 4)	198
Potatoes.	Thorney, Cambs.—N (0, 1, 2), P (0, 1, 2), K (0, 1, 2)	199
	March, Cambs.—N (0, 1, 2), P (0, 1, 2), K (0, 1, 2)	200
	Littleport, Cambs.—N $(0, 1, 2)$ , P $(0, 1, 2)$ K $(0, 1, 2)$	201
	Tunstall, Suttolk—P. K. magnesium sulphate dung	202
Sugar Beet.	Kidderminster, Worcester—S/A or N/S, super. or slag, K (0, 1, 2)	202-204
	-,, -, -, -, -, -, -, -, -, -, -, -, -	202-204
Б		
Experiments	carried out by Local Workers	
Hay.	Harlow, Essex—(0, 1, 2) levels of slag, super, mineral phosphate; K (0, 1	205
	Buriord, Oxford—Slag (0, 1, 3)	205
	Bakewell, Derby—N, P, K	206
	Bakewell, Derby—Artificials and compost, direct and residual effects	206
MILTER	Barnet, Herts—Residuals of K, chalk, phosphates	207
Potatoes.	Loughborough, Leicester—Mixed artificials (0, 1, 2, 3)	207
	Ingham, Lines.—Mixed artificials (0, 1, 2, 3, 4)	208
	Welton, Lines.—Mixed artificials (0, 1, 2, 3, 4)	208
Sugar Beet.	Market Rasen, Lines.—Methods of applying minerals: S/A dung	209-210
	Wragby, Lines.—Methods of applying minerals: S/A dung	210-211
	East Kirkby, Lines.—Methods of applying minerals: S/A depth of	
	Plane	212 - 213
	ploughing	
	ploughing	213 - 215
	Divion, Lines.—Dorax (U. 1. 2) artificials	215
	Digby Fen, Lincs.—Manganese sulphate (0, 1, 3). N/S	216
	Brough, Notts—Mixed artificials (0, 1, 2, 3)	217
	Newton-on-Trent, Lincs.—Mixed artificials (0, 1, 2, 3)	217
	North Muskham, Notts—Mixed artificials (0, 1, 2, 3)	218
	Crowle, Lincs.—Mixed artificials (0, 1, 2, 3), N, time of lifting	218-219
	Holton-le-Moor, Lincs.—Methods of singling	220
	Thornton, Lincs.—Methods of singling	220
Kale.	Kneeton, Notts—Cyanamide, nitro-chalk, S/A; super or slag	221
Itale.	Loughborough, Leicester—N, thinning	222

Note.—N denotes sulphate of ammonia or nitrate of soda, P denotes superphosphate, and K denotes any potash fertiliser.

## ERRATA 1932 Report

p. 203. Salt was applied at the rate of 0.85 cwt. Cl. per acre.

## 1936 Report

- p. 204. The third sentence from the foot of the page should read "The response to each was somewhat greater at the lower level of the other than at the higher level, but the difference was not significant."
- p. 223. In harvesting the kale, two guard rows were omitted from each plot. All yields and their standard errors should be multiplied by 18/16 or 1.125.
- p. 233, line 4 in the table. For "Blakewell" read "Bakewell."
- p. 272, line 3. For "3 x 2 factorial design" read "32 factorial design."