

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Report for 1937

[Full Table of Content](#)



Soya Beans

Rothamsted Research

Rothamsted Research (1938) *Soya Beans* ; Report For 1937, pp 51 - 51 - DOI:
<https://doi.org/10.23637/ERADOC-1-69>

place. The resulting seed was, therefore, no longer the pure Dorinni but a back cross, Golden Bantam having been one of the parents of Dorinni. This new strain, which we call Rothamsted Sweet Corn, is now being grown under a variety of conditions.

SOYA BEANS

In the spring of 1934 two varieties were planted, Manitoba Brown and Mandarin; the former being a semi-dwarf, early maturing, brown-seeded variety, developed at the Manitoba Agricultural College, while the latter is a medium sized variety with yellow seeds, much later in maturing.

Manitoba Brown ripened satisfactorily but Mandarin did not. It was, therefore, discarded.

In 1935 three other varieties were sown in addition to Manitoba Brown, namely, The Jap, an early maturing dwarf plant with pale green seeds; J. Yellow, a late maturing plant with yellow seeds; and Black, a medium sized plant with black seed coming later than Manitoba Brown. Frosts in the middle of May severely checked all four varieties but the plants recovered later and gave a fair yield. Manitoba Brown and Jap came out best, followed by Black but J. Yellow was too late to ripen properly. At Woburn also Manitoba Brown did well.

In 1936 at Rothamsted a May frost again checked the plants and a severe hail storm on June 21st did much damage. The yield of seed was small, nevertheless the maturity was good. At Woburn the plants suffered from rabbits, hares and birds.

In 1937 some more varieties were received from the Manitoba Agricultural College, one of which, Tokio, is promising both in yield and early maturity; the seed is dark but it may be possible to remedy this by suitable hybridisation and selection.

Prof. Southworth has now more assistance than before and has been able to commence more intensive study of the morphological and physiological characters of soya bean and he is trying to obtain new varieties better suited to our conditions than the existing sorts. We have been fortunate in securing the help of a collaborator in South Africa who plants the seeds during our winter and returns them to us in time for planting during our summer; we thus secure two crops in one year which saves a good deal of time in making selections.

Vernalisation did not prove helpful either for soya beans or maize.

PYRETHRUM

In view of the importance of pyrethrum as an insecticide and of the fact that it grows well on light sandy soil, a number of experiments have been made to see if by manuring the yields can be raised to levels at which they would become remunerative without at the same time lowering the insecticidal efficiency of the crop.

The experiments were made at Woburn, and were continued over four years: both lime and fertilizers increased the yield of flowers and of pyrethrum, the substance which measures the insecticidal value, but in some seasons the effects were only slight.