

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Rothamsted Report for 1938

[Full Table of Content](#)



Chemical Analyses Etc

Rothamsted Research

Rothamsted Research (1939) *Chemical Analyses Etc* ; Rothamsted Report For 1938, pp 105 - 106 -
DOI: <https://doi.org/10.23637/ERADOC-1-86>

CHEMICAL ANALYSES OF MANURES USED IN REPLICATED EXPERIMENTS, 1938

Manures	%N	%P ₂ O ₅	%K ₂ O
Poultry Manure	3.52	3.09	1.62
" "	3.77	3.17	1.68
" " (1)	4.10	3.57	1.90
" "	3.69	3.52	2.12
Rape Dust	5.41	1.77	1.01
Soot	4.03, 4.83		
Town refuse (2)	1.00		
" (3)	0.77		
" (4)	0.62		
" (5)	0.68		
" (6)	1.01		
Dung (7)	0.66		
" (8)	0.69	0.83	0.75
" (9)	0.70	0.73	0.94
" (10)	0.75	1.00	1.14
" (11)	0.92	0.46	1.07
Rotted normal (12)	0.75	0.37	0.94
Rotted strawy (12)	0.74	0.39	1.06
Fresh normal (12)	0.66	0.28	0.82
Fresh strawy (12)	0.64	0.24	0.74
Sulphate of ammonia	21.1, 21.2		
Nitrate of soda	16.1		
Nitrochalk	16.0		
Cyanamide	20.9, 21.1		
Superphosphate	17.0 Total		
	15.8 Water Sol.		
Basic-slag, high-sol.	15.8 Total		
	15.0 Citric Sol.		
Sulphate of potash	49.1		
Muriate of potash	50.2, 53.2		
Muriate of potash (high grade)	63.1		

- (1) Rothamsted and Woburn Experiments.
- (2) Woburn Experiment.
- (3) Rothamsted Experiment.
- (4) Tunstall Experiment.
- (5) Rochester Experiment.
- (6) Chichester Experiment.
- (7) Rothamsted Sugar Beet Experiment.
- (8) Rothamsted Kale, Organic Manures Experiment.
- (9) Rothamsted Kale, Town refuse Experiment.
- (10) Woburn Kale, Organic Manures Experiment.
- (11) Woburn Kale, Town refuse Experiment.
- (12) Rothamsted Potato Experiment.

Three Course Rotation

Manures	% Organic Matter	% N	% P ₂ O ₅	% K ₂ O
Chaffed straw	77.4	0.54	0.30	1.66
Adco	12.7	0.42	0.39	0.30
Superphosphate	—		16.8 ⁽¹⁾ 17.0 ⁽²⁾	—
Sulphate of ammonia	—	21.1 (1) (2)	—	—
Muriate of potash	—	—	—	51.4 ⁽¹⁾ 53.2 ⁽²⁾
Sulphate of potash	—	—	—	49.1
Nitrate of soda	—	16.1	—	—

(1) Applied in Autumn. (2) Applied in Spring.

Four Course Rotation				
Manures	% Organic Matter	% N	% P ₂ O ₅	% K ₂ O
Chaffed straw	77.4	0.54	0.30	1.66
Dung	14.7	0.52	0.40	1.32
Adco	12.7	0.42	0.39	0.30
Superphosphate	—	—	16.8 (1) (2)	—
Mineral phosphate (90% through 120 mesh)	—	—	25.9 (1) (2)	—
Muriate of potash	—	—	—	51.4 (1) (2)
Sulphate of ammonia	—	21.1	—	—

(1) Applied in Autumn. (2) Applied in Spring.

Six Course Rotation

Sulphate of ammonia	21.2 % N
Superphosphate	16.8(1), 17.0(2), % P ₂ O ₅ (Total)
Muriate of potash	51.4(1), 53.2(2), % K ₂ O

(1) Applied in Autumn. (2) Applied in Spring.

Long Period Cultivation Experiment

Cyanamide	21.1 % N
Nitrochalk	16.0 % N
Superphosphate	17.0 % P ₂ O ₅ (Total)
Muriate of potash	53.2 % K ₂ O

AVERAGE WHEAT YIELDS OF VARIOUS COUNTRIES

Country	Mean yield per acre, 1928-37 cwt.	Country	Mean yield per acre, 1928-37 cwt.
Great Britain	17.8	Denmark	23.4
England and Wales	17.6	Argentina	7.2
Hertfordshire	16.6	Australia	6.2
France	12.0	Canada	7.0
Germany	17.3	United States	7.1
Belgium	20.6	U.S.S.R. (Europe and Asia)	6.4*

Note.—Figures for Great Britain, England and Hertfordshire are taken from the Ministry of Agriculture's "Agricultural Statistics," Vol. 72. Other figures from "International Year Book of Agricultural Statistics," 1930-38.

* Excluding 1931.

CONVERSION TABLE

1 acre (10 sq. chains or 4,840 sq. yards)	0.405 Hectare
1 bushel (Imperial) (8 gallons)	0.364 Hectolitre
1 lb. (pound avoirdupois)	0.454 Kilogramme
1 cwt. (hundredweight, 112 lb.)	50.8 Kilogrammes
1 ton (20 cwt. or 2,240 lb.)	1016 Kilogrammes
1 metric quintal or Doppel Zentner (Dz)	{ 100.0 Kilogrammes
1 metric ton (tonne)	220.46 lb.
1 bushel per acre	1000 Kilogrammes
1 lb. per acre	0.899 Hectolitre per Hectare
1 cwt. per acre	1.121 Kilogrammes per Hectare
1 ton per acre	1.255 dz. per Hectare
1 dz. per Hectare	25.11 dz. per Hectare
1 kg. per Hectare	0.797 cwt. per acre
	0.892 lb. per acre

In America the Winchester bushel is used = 35.238 litres. 1 English bushel = 1.032 American bushels. In America 1 cwt. = 100 lb.

The yields of grain in the replicated experiments are given in cwt. per acre. One bushel of wheat weighs 60 lb., of barley weighs 52 lb., of oats weighs 42 lb. approximately.