


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Rothamsted Experimental Station Report for 1968 - Part 2



[Full Table of Content](#)

9. Microbiology of Broadbalk Soils - Whole Section

Rothamsted Research

Rothamsted Research (1969) *9. Microbiology of Broadbalk Soils - Whole Section* ; Rothamsted Experimental Station Report For 1968 - Part 2, pp 175 - 185 - **DOI:**
<https://doi.org/10.23637/ERADOC-1-2>

9. MICROBIOLOGY OF BROADBALK SOILS

JANE MEIKLEJOHN

Microbiological work began on Broadbalk after interest had been aroused in the practical benefits of partial soil sterilisation in increasing the growth of plants, and reasons for its effects were being sought. The number of bacteria in soil was known to increase greatly after partial sterilisation and this was suspected to be the reason for the increased fertility. Russell and Hutchinson (1913) suggested that partial sterilisation kills some organisms, and they favoured protozoa, that usually restricted bacterial populations and so fertility. Hence, other evidence for the beneficial effects of bacteria and possible harmful ones of protozoa was sought. The extremes of fertility represented by the Broadbalk plots 2, FYM, and 3, unmanured, provided good potential sites and were the first to be examined.

The comparisons of bacterial counts gave some support to the idea that bacteria might be beneficial, because plot 2 contained two or three times as many per gram of soil as plot 3, and the amounts of nitrate in these and several other soils also fluctuated with the numbers of bacteria. That soils did have a specific protozoal fauna was also first shown by work on these plots by Martin and Lewin (1915), but produced little evidence that they were harmful because plot 2 contained more individuals than plot 3, though fewer species. In both plots, numbers were fewer when the soil contained only 10% water than when it contained 20%.

Crump (1920) found that flagellates were the most abundant group of protozoa in plot 2, and averaged 28400 cells per gram in the top inch of soil. Amoeba were also numerous, averaging 7700, whereas ciliates were few, averaging only 56 per gram. However, counts made by Cutler and Crump (1920) at short intervals showed that the populations of three different species of flagellates fluctuated greatly between counts and that the active forms exhibit a daily periodicity, a day with many was followed by a day with few. Their finding that the number of bacteria and active amoeba in the soil were correlated, varying inversely over a period of 16 days, again seemed to suggest that protozoa might be limiting soil fertility, but much work since has shown that although bacterial predators, including protozoa, may limit numbers, they do not restrict bacterial activity (Thornton & Crump, 1953).

The algae have been rather neglected, and studied only by Bristol Roach (1927). Plots 2 and 3 had similar numbers of species, but plot 2 more than twice as many individuals (38000/g of top soil) as plot 3 (15000). These numbers omit blue-green algae, which could not be counted by the method Roach used, but these occurred and were sometimes plentiful enough to form a crust on the soil surface. It was of interest that she found algae living 12 in. below the soil surface, for then it was not known that many algae can live heterotrophically.

ROTHAMSTED REPORT FOR 1968, PART 2

Work since 1930

Since 1930 methods of estimating the numbers of organisms have greatly improved, and although Skinner, Jones and Mollison (1952) confirmed that plot 2 contained more than plot 3, the populations they reported enormously exceeded those reported by Russell and Appleyard (1917). The direct-count method they used gives the best measure of total populations, and showed an average in plot 2 of 2500 and a maximum of 4000 million/g of soil, and in plot 3 an average of 1500 and a maximum of 2500 million. They also examined plot 7 (N₂PKNMg), and this had similar populations to plot 3, although its wheat yields resemble those of plot 2, so bacterial populations did not reflect the fertility as indicated by crop yields.

Singh (1937) was the first to examine fungi and actinomycetes, not only on plots 2B and 3, but also on plot 10 (N₂) plot 11 (N₂P) and plot 13 (N₂PK). He found the same species in all plots, but differences between the numbers in different plots, with plot 2B having most and plot 3 fewest (Table 9-1).

TABLE 9-1
Plate counts of fungi and actinomycetes

Plot	Average numbers in top 6 in.: thousands per gram soil	
	Fungi	Actinomycetes
2B	490	2350
13	320	1500
11	250	1100
10	225	1000
3	175	700

Skinner (1951) also found many actinomycetes in plots 2, 3 and 7, though most were present as spores and not as growing mycelium. A recent series of counts of actinomycetes and fungi (see Etheridge, p. 181) was made to seek a biological explanation for take-all being less prevalent on Broadbalk than on nearby fields and to study the differences between the populations in the soil surrounding the roots (rhizosphere) and those on the roots (rhizoplane).

Protozoa. Singh (1946, 1949) having improved the method for counting soil micro-predators, estimated the population of amoebae in the soils of plots 2, 3 and 7. Plot 3 had fewest, averaging 17000/g, of which 10000/g were active (not encysted) cells and plot 2 most, averaging 72000/g (52000 active), and plot 7, which gives comparable yields to plot 2 average 48000/g (40000 active). Singh (1947a) also isolated some peculiar and little-known soil micro-organisms, including several of the higher Myxobacteria, from plot 3. Myxobacteria, which differ from true bacteria in lacking a rigid cell wall, include the widely-distributed cellulose-decomposer *Sporocytophaga myxococcoides*, discovered at Rothamsted by Hutchinson & Clayton (1919), and several species that break down chitin. However, the species Singh found were all predatory on other bacteria, which they killed by toxins and then digested by extra-cellular enzymes, and they formed characteristic fruiting bodies.

Singh (1947b) also found several members of the Acrasiae in Broadbalk

MICROBIOLOGY OF BROADBALK SOILS

soils. In their active form these unusual organisms are small amoebae, which feed and reproduce as do other soil amoebae, but sometimes they collect into a pseudo-plasmodium which produces a sporangium and the spores hatch into new small amoebae. Species of *Dictyostelium*, a typical member of the Acrasiae, were obtained from the soil of Broadbalk plots 3 and 7.

Another peculiar soil organism isolated was *Leptomyxa reticulata* (Singh, 1948), which is a very large amoeboid creature consisting of a thin multi-nucleate sheet of protoplasm that can attain a width of 3 mm, and feeds on bacteria. As these three organisms occurred in the unmanured plot, Singh concluded that they were indigenous to soil and not introduced in dung, but this conclusion may not be fully justified because Broadbalk was for long cultivated by horse-drawn implements.

Nitrifiers and nitrogen fixers

Nitrifiers. As part of a study on Winogradsky's Azotobacter test for soil fertility, Ziemiecka (1932) counted ammonia-oxidising bacteria on several of the Rothamsted Classical field experiments, using the silica-gel plate method. She found most in plots with bulky organic manures, and fewest in plots with no added nitrogen. The counts for Broadbalk were: plot 2, (FYM) 2243 cells/g; plot 19 (rape cake) 1336/g; plot 8 (N₃PKNaMg) 1949/g; plot 5 (PKNaMg) 406/g; and plot 3 (unmanured) 252/g.

TABLE 9.2
Numbers of nitrifying bacteria per gram dry soil, 1962

Date	Plot	Ammonia oxidisers	Nitrite oxidisers
1958	2A	360	45
1961 July	3 Fallow	6480	100
	3 Cropped	6300	390
1961 Oct.	5 Fallow	26300	412
1961 Aug.	7 Fallow	>11500	>11500
	7 Cropped	>11500	>11500
1961 Oct.	7 Fallow	26500	830
1958	8	720	720
1961 Aug.	10 Fallow	3350	210
	10 Cropped	6670	210
1961 Jan.	Wilderness Wood	4070	128
June	Wilderness Wood	3450	108
1961 Aug.	Wilderness Grass (grazed by sheep)	2870	45
Oct.	Wilderness Grass	2010	15
1962 Jan.	Wilderness Grass	8580	33
May	Wilderness Grass	>16000	65
June	Wilderness Grass	>13000	1700

The ammonia-oxidisers operate the first stage in the process of nitrification that makes combined nitrogen available to crop plants. The nitrite they make by the oxidation of ammonia is further oxidised to nitrate by other nitrifying bacteria, the nitrite-oxidisers. Meiklejohn (1962) found adequate populations of both kinds of nitrifier in six of the Broadbalk plots, 2, 3, 5, 7, 8 and 10, with no evident differences between plots. Soil from the Wilderness at the west end of Broadbalk also contained nitrifiers of both kinds, both in the wooded end and in the end under grass. This is of interest

ROTHAMSTED REPORT FOR 1968, PART 2

because many woodland soils in Britain do not contain nitrifiers (Table 9-2)

An ammonia-oxidising bacterium, *Nitrosomonas europaea*, was isolated from the soil of plot 2 (Meiklejohn, 1949); it, and the nitrite-oxidiser *Nitrobacter winogradskyi*, seem to be the only two abundant nitrifiers in Rothamsted soils.

Nitrogen fixers. The free-living aerobic bacterium *Azotobacter chroococcum* adds nitrogen to the soil by taking it from the air and building it into its own cell substances. Ashby (1907) discovered it in soil from Broadbalk Wilderness and in the Broadbalk drains. Ziemiecka (1932) used Winogradsky's plate method to estimate populations in various fields at Rothamsted in September, October and November 1931. Her counts on Broadbalk were largest on plots 5 (PKNaMg but no nitrogen) 8400/g in September and 2055/g in October, and on plot 9 (N₁*PKMaMg) 1593/g in October. In November she found 338 *Azotobacter* per g. in the Wilderness grass, and 120/g in the Wilderness wood soil. In 1960 Burlingham counted *Azotobacter* in rhizospheres and in soil between the rows of plants in plots 2, 5 and 8 (Table 9-3). Counts in January and March 1960 showed no 'rhizosphere effect', i.e. populations were no larger in the soil immediately around the roots than in the inter-row soil. Plot 2 has many more *Azotobacter* than plots 5 or 8, and contrary to the results of other workers she found no *Azotobacter* in soil under the Wilderness grass.

TABLE 9-3
Numbers of Azotobacter in rhizosphere and in other soil, 1960

		Number of <i>Azotobacter</i> (cells/g dry soil)			
		Plot 2		Plot 5	Plot 8
		Rhizosphere	Soil	Soil	Soil
Jan. 1960		4500	2060	115	14
		4600	3080	74	64
		13800	11000	108	38
		1800	3500	35	69
		Plot 2		Plot 5	
		Rhizosphere	Soil	Rhizosphere	Soil
Mar. 1960		1200	238	0	124
		700	6000	234	126
		12000	56800	160	256
		2300	3500	110	260
		3500	1400	0	87
	1100	5300	0	125	

Wilderness grass: none found in any sample.

After the system of a fallow every fifth year was found to act like a dressing of quick-acting nitrogen, it was thought some of the effect of fallow might reflect nitrogen fixation in the fallow soil. Therefore the populations of nitrogen fixers were estimated before, during and after the fallow year in several of the plots. Most counts were of *Azotobacter chroococcum*, an easily recognisable species, but a few counts were also made of the anaerobe, *Clostridium pasteurianum*. Counts made during the 1955-58 seasons seemed

MICROBIOLOGY OF BROADBALK SOILS

to support the idea that fallowing stimulated nitrogen fixation for the first crop after fallow. *Azotobacter* were few in sections under the 4th successive wheat crop and in the soil being fallowed, but increased greatly in the first wheat crop after the fallow. The increase in *Azotobacter* was greatest in the plots where the increased yield from fallow was greatest (Table 9.4). Plots given ammonium sulphate every year had fewer *Azotobacter* than the other plots (Meiklejohn, 1965).

However, *Azotobacter* in the Broadbalk soils seemed too few to fix enough nitrogen to affect the crop growth. Although an improved technique (Brown, Burlingham & Jackson, 1962) increased the count, the largest population found was only 9400 cells/g dry soil, which again is too few for *Azotobacter* to affect crop growth through nitrogen fixation. Nevertheless the *Azotobacter* numbers indicate that conditions are suitable for nitrogen fixation by other species, such as the anaerobe *Clostridium pasteurianum*, which occurs all over Broadbalk in much larger numbers than *Azotobacter*, ranging up to 100 000 cells per gram dry soil (Meiklejohn, 1956). The possibility that an anaerobe is the active fixing agent in Broadbalk soil is supported by laboratory observations of nitrogen fixation by soil from Broadbalk plot 3 incubated anaerobically with straw (Barrow & Jenkinson, 1962).

TABLE 9.4
Effect of fallow on Azotobacter numbers in 1962

		Largest number (cells/g soil)	Mean
Plots without added nitrogen			
3	Fallow	137	53 ± 16
	1st crop after fallow	1194	430 ± 135
5	Fallow	282	166 ± 30
	1st crop after fallow	894	400 ± 121
Plots with annual ammonium sulphate			
7	Fallow	410	134 ± 49
	1st crop after fallow	1429	631 ± 190
10	Fallow	555	129 ± 76
	1st crop after fallow	260	146 ± 27
Other nitrogen treated plots			
16	Fallow	4524	1003 ± 594
	1st crop after fallow	3782	1803 ± 435
17	Fallow	2620	580 ± 343
	1st crop after fallow	9406	2442 ± 1276

Symbiotic Nitrogen Fixation: Legume Nodule Bacteria

By P. S. NUTMAN

Four species of the genus *Rhizobium*, which form nodules on the roots of leguminous plants occur in Broadbalk soils, but in populations that are much smaller than in neighbouring fields.

Nodule bacteria were counted in samples of soil taken in March 1968

ROTHAMSTED REPORT FOR 1968, PART 2

from plots 3, 5, 8 and 16 in sections I, II, III, IV and VA, using dilutions of soil to inoculate sterile-grown seedlings of the test legumes, viz. *Vicia hirsuta* to test for the presence of *Rhizobium leguminosarum*, *Trifolium pratense* for *R. trifolii*, *Lotus corniculatus* for *R. lupini* and *Medicago sativa* for *R. meliloti*. The test seedlings form nodules and grow only if the inoculum contains the appropriate species. Samples were also taken from plots 2, 11 and 18 to count *R. leguminosarum* only.

Table 9.5 shows the logarithms of estimated numbers per gram dry soil. (Nutman, 1962.)

TABLE 9.5
Rhizobium counts on Broadbalk, March 1968

Section	Herbi- cides since	Log estimated no./g dry soil											
		<i>R. leguminosarum</i>						<i>R. trifolii</i>					
		Plot 3	5	16	8	2	11	18	3	5	16	8	
I	1957	5.12	5.14	4.83	5.16	—	—	—	2.64	2.10	3.12	>4.08	
II	1964	4.42	3.48	4.29	4.55	4.15	3.81	4.20	2.95	1.26	2.29	>3.97	
III	1964	4.89	4.50	3.87	4.02	—	—	—	2.94	1.96	1.41	3.55	
IV	1964	3.68	4.59	2.97	4.89	—	—	—	3.32	2.53	1.85	>4.00	
VA	none	4.52	5.07	4.20	5.70	5.48	3.99	>5.52	3.32	1.38	1.75	>3.20	
		<i>R. lupini</i>				<i>R. meliloti</i>							
		Plot 3	5	16	8	3	5	16	8				
I	1957	1.02	2.38	1.33	<0.51	<0.32	2.32	<0.59	<0.87				
II	1964	<0.33	0.61	<0.96	1.06	<1.18	0.96	<0.45	<0.27				
III	1964	<0.47	1.58	0.75	2.53	0.89	1.71	1.30	<0.42				
IV	1964	0.75	<0.49	1.20	<0.15	0.97	<0.56	0.68	<0.15				
VA	none	2.13	1.16	1.19	1.98	<0.67	0.47	<1.04	<0.15				

Rhizobium leguminosarum is the most abundant species in Broadbalk. The average of all counts was 28000/g dry soil, and ranged from 1800/g dry soil in one sample of plot 11 to >87000 in plot 18. In most plots numbers varied by less than a hundred-fold, and no significant differences were noted between sections or plots. All *R. leguminosarum* cells nodulating the test plant were effective in fixing nitrogen.

The next most abundant species of nodule bacteria is *Rhizobium trifolii* of which most were found in plot 3 (unmanured) and plot 8 (N₃PKNaMg). Maximum, minimum and average counts were <14500, 7 and 320/g soil and populations varied more than *R. leguminosarum*. No consistent differences were found between sections or plots. About 10% of the bacteria that nodulated *Trifolium pratense* when inoculated with the most dilute suspensions did not fix nitrogen or fixed very little on this host.

Rhizobium lupini occurs sparsely and sporadically; it was not found in seven plots (total of 10.4 g sampled) and in the others the average number was 310/g and the maximum 1620/g. Bacteria from all except one sample (plot 3, section II) were effective in fixing nitrogen in nodules of *Lotus corniculatus*.

Rhizobium meliloti occurs infrequently in low numbers. It was absent from nearly half the samples (total weight 19 g) and was few in the samples where it occurred; an average of 13/g and a maximum of 229/g. More than half of the strains were fully effective with *Medicago sativa*.

In fields such as Broadbalk where legume crops are not grown the nodule bacteria may be brought in from neighbouring fields or may persist on leguminous weeds. Whichever way they are introduced their distribution does not correspond with that of leguminous weeds.

MICROBIOLOGY OF BROADBALK SOILS

Medicago lupulina (black medick) is the commonest leguminous weed on the plots sampled, but its nodule bacteria is the rarest of the four species. Black medick is most frequent in plots 3 and 5, much less frequent on plot 16 and rarely occurs on plot 8, which also contains fewer *R. meliloti* than other plots. Herbicides have decreased numbers of *M. lupulina* in most years (see p. 197) but this has had no effect on the numbers of its nodule bacteria in the different sections.

The natural hosts of *R. leguminosarum* found in Broadbalk are common vetch (*Vicia sativa*) and meadow vetchling (*Lathyrus pratensis*), *Vicia sativa* being most frequent, particularly in certain years on plots 3, 5 and 18. *Lathyrus pratensis* occurs rarely and sporadically and very few of either legume occur on plot 16 and none on plot 8. There is no significant difference in numbers of *R. leguminosarum* between plots or between sections treated with herbicides and those not treated, in spite of the effect of herbicides in diminishing populations of *V. sativa* on all plots where it occurs.

Although *Rhizobium trifolii* is more widely distributed on Broadbalk and more abundant than *R. lupini* or *R. meliloti*, its natural host plants, the clovers, occur very rarely and not at all in most years.

No known host plant of *R. lupini* has been recorded on Broadbalk so that the occurrence of this species of *Rhizobium* cannot be related to weed history.

Numbers of *Rhizobium* in Rothamsted soil under leguminous crops are larger by several orders of magnitude than those found in Broadbalk. For example in Sawyers field (limed plots) under *Medicago lupulina*, tick beans and red clover, the average number of the specific rhizobia for these legumes were respectively 500 000, >2 000 000 and 700 000 per g/dry soil (Nutman, 1963). In the long term fallow experiment in Highfield the total *Rhizobium* population is small, but, as on Broadbalk, the clover and bean nodule bacteria are nevertheless more abundant than the lotus and medick bacteria. This suggests that the nodule bacteria found in Broadbalk are unlikely to have multiplied in the rhizospheres of the relatively few leguminous weeds but more likely to have been brought in from neighbouring fields.

Rhizosphere and Rhizoplane Fungi

By JUDITH ETHERIDGE

Take-all, caused by *Ophiobolus graminis*, is usually less severe on wheat on Broadbalk than on wheat grown successively in other fields at Rothamsted. In 1960 the fungi of the rhizosphere and rhizoplane of wheat from Broadbalk were compared with those from other fields in an attempt to identify any organism that might inhibit the development of take-all on Broadbalk. The rhizosphere fungi were studied by the dilution plate method using a glucose peptone medium (Johnson *et al.*, 1959) with 30 mg aureomycin per litre. After the rhizosphere soil had been washed away, the roots were washed in only two changes of sterile water, then cut into 0.5 cm

ROTHAMSTED REPORT FOR 1968, PART 2

TABLE 9-6
Frequency of fungi in rhizosphere and rhizoplane of Broadbalk wheat, 1960

Species	Plot 3 (4th crop after fallow)		Plot 7 (4th crop after fallow)			
	Rhizoplane	Rhizosphere	Rhizoplane	Rhizosphere		
	% root pieces colonised (all dates)	No. of dates when isolated*	% total (all dates)	No. of dates when isolated*	% total (all dates)	No. of dates when isolated*
Aspergillus spp.	0.1	2	0	0	0	0
Aureobasidium spp.	3.2	7	0	0	0	0
Beauveria bassiana	0.05	1	0	0	0	0
Botrytis cinerea	0.8	5	0.04	1	0.5	6
Cephalosporium spp.	2.5	10	3.6	9	2.9	9
Geomyces vulgaris	0.6	7	6.7	11	5.4	11
Gliocladium roseum	1.8	11	0.5	5	1.3	3
Paecilomyces carneus	0.2	2	0.6	4	0.2	2
Penicillium spp.	40.6	11	30.3	11	20.4	11
Sporotrichum carnis	0.05	1	0.2	1	0.9	3
Tilachlidium sp.	0.1	3	0.4	4	0.6	3
Verticillium spp.	0.2	4	0.1	2	0.6	4
Trichoderma spp.	4.0	11	0.8	8	2.2	9
Graphium sp.	0.05	1	0.1	1	0	0
Truncatella truncata	0.05	1	0.2	2	0.2	2
Alternaria spp.	0.9	7	0	0	0.04	1
Cladosporium herbarum	26.2	11	26.7	11	35.9	11
Periconia spp.	2.0	5	0.8	2	0.1	1
Stachybotrys sp.	0	0	0.1	2	0	0
Trichosporium cerealis	0	0	0	0	0.2	2
Cylindrocarpon spp.	0.5	7	0.4	4	0.4	4
Epicoccum nigrum	0	0	0.05	1	0	0
Fusarium avenaceum	3.3	10	0.6	3	0.3	3
F. chlamydosporum	0.4	4	0.2	1	0.2	1
F. culmorum	0.5	6	0.2	2	0.4	3
F. dimerum	0.1	1	0	0	0	0
F. merismoides	0.1	2	2.1	4	0.2	2
F. poae	0.3	4	0	0	0	0
F. sambucinum	0.3	2	0	0	0	0
F. solani	0.05	1	0	0	0	0
Myrothecium sp.	2.3	9	2.3	7	0.1	1
Coniothyrium spp.	0.05	1	0.1	2	0	0
Diplodia sp.	0.8	8	1.0	8	0.4	3
Phoma spp.	1.3	5	0.6	5	0.4	3
Absidia glauca	0.7	7	0	0	0.2	3
Mortierella and other similar						
Phycomycetous fungi	42.5	11	8.0	11	9.0	11
Mucor spp.	7.0	11	1.9	11	2.1	8
Rhizopus sp.	0.05	1	0	0	0	0
Yeasts	1.1	5	1.0	3	0.6	3
Chaetomium indicum	0	0	0.1	1	0	0
Ophiobolus graminis	0.1	1	0	0	0	0
Mycelia sterila, dark	10.0	11	6.6	11	11.0	11
Mycelia sterila, light	5.4	11	2.6	8	3.6	7

* Of 11 sampling dates from January to July.

lengths and 20 pieces were plated at each sampling date. The 'rhizoplane' population of fungi counted by this method are not the same as the 'root surface' fungi obtained by Harley and Waid's method (1955) which employs a more thorough washing technique. Within the limits imposed by these methods (which greatly favour the isolation of fungi present as spores)

MICROBIOLOGY OF BROADBALK SOILS

no consistent differences were found between Broadbalk and other fields. Non-rhizosphere soils were not studied, and this account describes results from Broadbalk only.

The 4th wheat crop after a one-year fallow on plot 3 (no manure) and plot 7 (N₂PKNaMg) was sampled at intervals from January to July. Table 9.6 lists the fungi identified. In addition to these spore-producing species, several sterile fungi were isolated, of which the most frequent has a fast-growing aerial mycelium, pinkish-fawn at first, but becoming dark-grey to black within four days. Twenty-four of the 32 rhizosphere fungi identified occurred on both plots. *Penicillium* spp., *Cladosporium* spp., *Geomyces vulgaris* and the *Mortierella*-type group were the most abundant species. On each of the 11 sampling dates these four groups constituted more than 33% of the total number of fungi and from May to July at least 65%. *Penicillium* spp. were most abundant at the earlier samplings, and *Cladosporium* became dominant later in the growing season.

Actinomycetes in Broadbalk soils. Actinomycetes in rhizosphere and non-rhizosphere soils from plot 7 were counted in May and June 1962, 1963 and 1964 using the dilution plate method (Table 9.7). The rhizosphere/non-rhizosphere ratios (R/S) were small in all 3 years, and the counts showed no evidence that actinomycetes were stimulated in the rhizosphere.

TABLE 9.7
Count of actinomycetes in rhizosphere and non-rhizosphere soils from Broadbalk plot 7 at a dilution of approx 1/40000, 1962-64

Year	Crop	Medium used	Date of sampling	Number actinomycetes millions per g dry soil		R/S
				Rhizo-sphere (R)	Non rhizo-sphere (S)	
1962	4th after fallow	Chitin ¹	16 May	0.67	0.59	1.14
			6 June	0.67	0.69	0.98
			28 June	0.96	0.90	1.07
1963	2nd after fallow	Chitin	8 May	1.25	2.13	0.59
			29 May	0.32	2.30	0.14
			26 June	0.85	0.85	1.00
1964	3rd after fallow	Soil extract agar ²	26 May	1.71	1.73	0.99
			16 June	1.85	2.12	0.87

¹ Lingappa & Lockwood, 1962.

² Flentje, 1956.

The Vesicular-arbuscular Mycorrhizal Fungi

By BARBARA MOSSE

Three plots on Broadbalk were surveyed for spores of *Endogone*, a mycorrhizal fungus, in 1961/2 and 1966 (Mosse & Bowen, 1968a, b), and four plots in 1968 (Hayman, unpublished). Each survey showed spores of four types, designated respectively as yellow vacuolate, laminate, bulbous reticulate and white reticulate, and that they occurred in different proportions in different plots. The unmanured plot contained more spores than the others

ROTHAMSTED REPORT FOR 1968, PART 2

and was especially rich in reticulate spores. Yellow vacuolate spores were most abundant in the FYM plot. Of the four types, laminate spores were fewest, whereas in the nearby field, Little Knott, they were the only ones found and were numerous in some plots, especially those not given nitrogen.

Acknowledgement

We thank G. J. S. Ross for the statistical analyses.

REFERENCES

- ASHBY, S. F. (1907) Some observations on the assimilation of atmospheric nitrogen by a free living organism—*Azotobacter chroococcum* of Beijerinck. *J. agric. Sci., Camb.* **2**, 35–51.
- BARROW, N. J. & JENKINSON, D. S. (1962) The effect of water-logging on fixation of nitrogen by soil incubated with straw. *Pl. Soil* **16**, 258–262.
- BRISTOL ROACH, B. M. (1927) On the algae of some normal English soils. *J. agric. Sci., Camb.* **17**, 563–588.
- BROWN, M. E., BURLINGHAM, S. K. & JACKSON, R. M. (1962) Studies on *Azotobacter* species on soil. I. Comparison of media and techniques for counting *Azotobacter* in soil. *Pl. Soil* **17**, 308–319.
- BURLINGHAM, S. H. (1960) Unpublished results.
- CRUMP, L. M. (1920) Numbers of Protozoa in certain Rothamsted soils. *J. agric. Sci., Camb.* **10**, 182–198.
- CUTLER, D. W. & CRUMP, L. M. (1920) Daily periodicity in the numbers of active soil flagellates; with a brief note on the relation of trophic amoebae and bacterial numbers. *Ann. appl. Biol.* **7**, 11–24.
- ETHERIDGE, J. (1960) Unpublished results.
- ETHERIDGE, J. (1964) Unpublished results.
- FLENTJE, N. J. (1956) Studies on *Pellicularia filamentosa* (Pat) Rogers. I Formation of the perfect stage. *Trans. Br. mycol. Soc.* **39**, 343–356.
- HARLEY, J. L. & WAID, J. S. (1955) A method of studying active mycelia on living roots and other surfaces in the soil. *Trans. Br. mycol. Soc.* **38**, 104–118.
- HAYMAN, D. (1968) Unpublished results.
- HUTCHINSON, H. B. & CLAYTON, J. (1919) On the decomposition of cellulose by an aerobic organism (*Spirochatea cytophaga* n.sp.). *J. agric. Sci., Camb.* **9**, 143–173.
- JOHNSON, L. F. *et al.* (1959) *Methods for studying soil microflora—plant disease relationships*. Minneapolis; Burgess Publishing Co.
- LINGAPPA, Y. & LOCKWOOD, J. L. (1962) Chitin media for selective isolation and culture of actinomycetes. *Phytopathology* **52**, 317–323.
- MARTIN, C. H. & LEWIN, K. R. (1915) Notes on some methods for the estimation of soil protozoa. *J. agric. Sci.* **7**, 106–119.
- MEIKLEJOHN, J. (1949) Isolation of *Nitrosomonas* from Rothamsted soil. *Nature, Lond.* **164**, 667.
- MEIKLEJOHN, J. (1956) Preliminary notes on numbers of nitrogen fixers on Broadbalk field. *Proc. 6th int. Cong. Soil Sci., Paris, C*, 243–248.
- MEIKLEJOHN, J. (1962) Unpublished results.
- MEIKLEJOHN, J. (1965) *Azotobacter* numbers on Broadbalk field, Rothamsted. *Pl. Soil* **23**, 227–235.
- MOSSE, B. & BOWEN, G. D. (1968a) A key to the recognition of some *Endogone* spore types. *Trans. Br. mycol. Soc.* **51**, 469.
- MOSSE, B. & BOWEN, G. D. (1968b) The distribution of *Endogone* spores in some Australian and New Zealand soils, and in an experimental field soil at Rothamsted. *Trans. Br. mycol. Soc.* **51**, 485.
- NUTMAN, P. S. (1963) *Rep. Rothamsted exp. Stn for 1962*, 80, 81.
- NUTMAN, P. S. (1968) Unpublished results.
- RUSSELL, E. J. & APPLEYARD, A. (1917) The influence of soil conditions on the decomposition of organic matter in the soil. *J. agric. Sci.* **8**, 385–417.
- RUSSELL, E. J. & HUTCHINSON, H. B. (1913) The effect of partial sterilisation of soil on the production of plant food. II. The limitation of bacterial numbers in normal soils and its consequences. *J. Agric. Sci., Camb.*, **5**, 152–221.

MICROBIOLOGY OF BROADBALK SOILS

- SINGH, J. (1937) Soil fungi and actinomycetes in relation to manurial treatment, season and crop. *Ann. appl. Biol.* **24**, 154-168.
- SINGH, B. N. (1946) A method of estimating the numbers of soil protozoa, especially amoebae, based on their differential feeding on bacteria. *Ann. appl. Biol.* **33**, 112-119.
- SINGH, B. N. (1947a) Myxobacteria in soils and composts, their distribution, number and lytic action on bacteria. *J. gen. Microbiol.* **1**, 1-10.
- SINGH, B. N. (1947b) Studies on soil Acrasidae. 1. Distribution of species of *Dictyostelium* in soils of Great Britain, and the effect of bacteria on their development. *J. gen. Microbiol.* **1**, 11-21.
- SINGH, B. N. (1948) Studies on giant amoeboid organisms. 1. The distribution of *Leptomyxa reticulata* Goodey in soils of Great Britain, and the effect of bacterial food on growth and cyst formation. *J. gen. Microbiol.* **2**, 8-14.
- SINGH, B. N. (1949) The effect of artificial fertilizers and dung on the numbers of amoebae in Rothamsted soils. *J. gen. Microbiol.* **3**, 204-210.
- SKINNER, F. A. (1951) A method for distinguishing between viable spores and mycelial fragments of actinomycetes in soils. *J. gen. Microbiol.* **5**, 159-166.
- SKINNER, F. A., JONES, P. C. T. & MOLLISON, J. E. (1952) A comparison of a direct and a plate-counting technique for the quantitative estimation of soil micro-organisms. *J. gen. Microbiol.* **6**, 261-271.
- THORNTON, H. G. & CRUMP, L. M. (1953) Micro-predators in soil. *Rep. Rothamsted exp. Stn for 1952*, 164-172.
- ZIEMIECKA, J. (1932) The *Azotobacter* test of soil fertility applied to the classical fields at Rothamsted. *J. agric. Sci., Camb.*, **22**, 797-810.