


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Rothamsted Experimental Station Report for 1968 - Part 2



[Full Table of Content](#)

Symbiotic Nitrogen Fixation: Legume Nodule Bacteria

P. S. Nutman

P. S. Nutman (1969) *Symbiotic Nitrogen Fixation: Legume Nodule Bacteria* ; Rothamsted Experimental Station Report For 1968 - Part 2, pp 179 - 181 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-2>

MICROBIOLOGY OF BROADBALK SOILS

to support the idea that fallowing stimulated nitrogen fixation for the first crop after fallow. *Azotobacter* were few in sections under the 4th successive wheat crop and in the soil being fallowed, but increased greatly in the first wheat crop after the fallow. The increase in *Azotobacter* was greatest in the plots where the increased yield from fallow was greatest (Table 9.4). Plots given ammonium sulphate every year had fewer *Azotobacter* than the other plots (Meiklejohn, 1965).

However, *Azotobacter* in the Broadbalk soils seemed too few to fix enough nitrogen to affect the crop growth. Although an improved technique (Brown, Burlingham & Jackson, 1962) increased the count, the largest population found was only 9400 cells/g dry soil, which again is too few for *Azotobacter* to affect crop growth through nitrogen fixation. Nevertheless the *Azotobacter* numbers indicate that conditions are suitable for nitrogen fixation by other species, such as the anaerobe *Clostridium pasteurianum*, which occurs all over Broadbalk in much larger numbers than *Azotobacter*, ranging up to 100 000 cells per gram dry soil (Meiklejohn, 1956). The possibility that an anaerobe is the active fixing agent in Broadbalk soil is supported by laboratory observations of nitrogen fixation by soil from Broadbalk plot 3 incubated anaerobically with straw (Barrow & Jenkinson, 1962).

TABLE 9.4
Effect of fallow on *Azotobacter* numbers in 1962

		Largest number (cells/g soil)	Mean
Plots without added nitrogen			
3	Fallow	137	53 ± 16
	1st crop after fallow	1194	430 ± 135
5	Fallow	282	166 ± 30
	1st crop after fallow	894	400 ± 121
Plots with annual ammonium sulphate			
7	Fallow	410	134 ± 49
	1st crop after fallow	1429	631 ± 190
10	Fallow	555	129 ± 76
	1st crop after fallow	260	146 ± 27
Other nitrogen treated plots			
16	Fallow	4524	1003 ± 594
	1st crop after fallow	3782	1803 ± 435
17	Fallow	2620	580 ± 343
	1st crop after fallow	9406	2442 ± 1276

Symbiotic Nitrogen Fixation: Legume Nodule Bacteria

By P. S. NUTMAN

Four species of the genus *Rhizobium*, which form nodules on the roots of leguminous plants occur in Broadbalk soils, but in populations that are much smaller than in neighbouring fields.

Nodule bacteria were counted in samples of soil taken in March 1968

ROTHAMSTED REPORT FOR 1968, PART 2

from plots 3, 5, 8 and 16 in sections I, II, III, IV and VA, using dilutions of soil to inoculate sterile-grown seedlings of the test legumes, viz. *Vicia hirsuta* to test for the presence of *Rhizobium leguminosarum*, *Trifolium pratense* for *R. trifolii*, *Lotus corniculatus* for *R. lupini* and *Medicago sativa* for *R. meliloti*. The test seedlings form nodules and grow only if the inoculum contains the appropriate species. Samples were also taken from plots 2, 11 and 18 to count *R. leguminosarum* only.

Table 9.5 shows the logarithms of estimated numbers per gram dry soil. (Nutman, 1962.)

TABLE 9.5
Rhizobium counts on Broadbalk, March 1968

Section	Herbi- cides since	Log estimated no./g dry soil											
		<i>R. leguminosarum</i>						<i>R. trifolii</i>					
		Plot 3	5	16	8	2	11	18	3	5	16	8	
I	1957	5.12	5.14	4.83	5.16	—	—	—	2.64	2.10	3.12	>4.08	
II	1964	4.42	3.48	4.29	4.55	4.15	3.81	4.20	2.95	1.26	2.29	>3.97	
III	1964	4.89	4.50	3.87	4.02	—	—	—	2.94	1.96	1.41	3.55	
IV	1964	3.68	4.59	2.97	4.89	—	—	—	3.32	2.53	1.85	>4.00	
VA	none	4.52	5.07	4.20	5.70	5.48	3.99	>5.52	3.32	1.38	1.75	>3.20	
		<i>R. lupini</i>				<i>R. meliloti</i>							
		Plot 3	5	16	8	3	5	16	8				
I	1957	1.02	2.38	1.33	<0.51	<0.32	2.32	<0.59	<0.87				
II	1964	<0.33	0.61	<0.96	1.06	<1.18	0.96	<0.45	<0.27				
III	1964	<0.47	1.58	0.75	2.53	0.89	1.71	1.30	<0.42				
IV	1964	0.75	<0.49	1.20	<0.15	0.97	<0.56	0.68	<0.15				
VA	none	2.13	1.16	1.19	1.98	<0.67	0.47	<1.04	<0.15				

Rhizobium leguminosarum is the most abundant species in Broadbalk. The average of all counts was 28000/g dry soil, and ranged from 1800/g dry soil in one sample of plot 11 to >87000 in plot 18. In most plots numbers varied by less than a hundred-fold, and no significant differences were noted between sections or plots. All *R. leguminosarum* cells nodulating the test plant were effective in fixing nitrogen.

The next most abundant species of nodule bacteria is *Rhizobium trifolii* of which most were found in plot 3 (unmanured) and plot 8 (N₃PKNaMg). Maximum, minimum and average counts were <14500, 7 and 320/g soil and populations varied more than *R. leguminosarum*. No consistent differences were found between sections or plots. About 10% of the bacteria that nodulated *Trifolium pratense* when inoculated with the most dilute suspensions did not fix nitrogen or fixed very little on this host.

Rhizobium lupini occurs sparsely and sporadically; it was not found in seven plots (total of 10.4 g sampled) and in the others the average number was 310/g and the maximum 1620/g. Bacteria from all except one sample (plot 3, section II) were effective in fixing nitrogen in nodules of *Lotus corniculatus*.

Rhizobium meliloti occurs infrequently in low numbers. It was absent from nearly half the samples (total weight 19 g) and was few in the samples where it occurred; an average of 13/g and a maximum of 229/g. More than half of the strains were fully effective with *Medicago sativa*.

In fields such as Broadbalk where legume crops are not grown the nodule bacteria may be brought in from neighbouring fields or may persist on leguminous weeds. Whichever way they are introduced their distribution does not correspond with that of leguminous weeds.

MICROBIOLOGY OF BROADBALK SOILS

Medicago lupulina (black medick) is the commonest leguminous weed on the plots sampled, but its nodule bacteria is the rarest of the four species. Black medick is most frequent in plots 3 and 5, much less frequent on plot 16 and rarely occurs on plot 8, which also contains fewer *R. meliloti* than other plots. Herbicides have decreased numbers of *M. lupulina* in most years (see p. 197) but this has had no effect on the numbers of its nodule bacteria in the different sections.

The natural hosts of *R. leguminosarum* found in Broadbalk are common vetch (*Vicia sativa*) and meadow vetchling (*Lathyrus pratensis*), *Vicia sativa* being most frequent, particularly in certain years on plots 3, 5 and 18. *Lathyrus pratensis* occurs rarely and sporadically and very few of either legume occur on plot 16 and none on plot 8. There is no significant difference in numbers of *R. leguminosarum* between plots or between sections treated with herbicides and those not treated, in spite of the effect of herbicides in diminishing populations of *V. sativa* on all plots where it occurs.

Although *Rhizobium trifolii* is more widely distributed on Broadbalk and more abundant than *R. lupini* or *R. meliloti*, its natural host plants, the clovers, occur very rarely and not at all in most years.

No known host plant of *R. lupini* has been recorded on Broadbalk so that the occurrence of this species of *Rhizobium* cannot be related to weed history.

Numbers of *Rhizobium* in Rothamsted soil under leguminous crops are larger by several orders of magnitude than those found in Broadbalk. For example in Sawyers field (limed plots) under *Medicago lupulina*, tick beans and red clover, the average number of the specific rhizobia for these legumes were respectively 500 000, >2 000 000 and 700 000 per g/dry soil (Nutman, 1963). In the long term fallow experiment in Highfield the total *Rhizobium* population is small, but, as on Broadbalk, the clover and bean nodule bacteria are nevertheless more abundant than the lotus and medick bacteria. This suggests that the nodule bacteria found in Broadbalk are unlikely to have multiplied in the rhizospheres of the relatively few leguminous weeds but more likely to have been brought in from neighbouring fields.

Rhizosphere and Rhizoplane Fungi

By JUDITH ETHERIDGE

Take-all, caused by *Ophiobolus graminis*, is usually less severe on wheat on Broadbalk than on wheat grown successively in other fields at Rothamsted. In 1960 the fungi of the rhizosphere and rhizoplane of wheat from Broadbalk were compared with those from other fields in an attempt to identify any organism that might inhibit the development of take-all on Broadbalk. The rhizosphere fungi were studied by the dilution plate method using a glucose peptone medium (Johnson *et al.*, 1959) with 30 mg aureomycin per litre. After the rhizosphere soil had been washed away, the roots were washed in only two changes of sterile water, then cut into 0.5 cm

ROTHAMSTED REPORT FOR 1968, PART 2

and was especially rich in reticulate spores. Yellow vacuolate spores were most abundant in the FYM plot. Of the four types, laminate spores were fewest, whereas in the nearby field, Little Knott, they were the only ones found and were numerous in some plots, especially those not given nitrogen.

Acknowledgement

We thank G. J. S. Ross for the statistical analyses.

REFERENCES

- ASHBY, S. F. (1907) Some observations on the assimilation of atmospheric nitrogen by a free living organism—*Azotobacter chroococcum* of Beijerinck. *J. agric. Sci., Camb.* **2**, 35–51.
- BARROW, N. J. & JENKINSON, D. S. (1962) The effect of water-logging on fixation of nitrogen by soil incubated with straw. *Pl. Soil* **16**, 258–262.
- BRISTOL ROACH, B. M. (1927) On the algae of some normal English soils. *J. agric. Sci., Camb.* **17**, 563–588.
- BROWN, M. E., BURLINGHAM, S. K. & JACKSON, R. M. (1962) Studies on *Azotobacter* species on soil. I. Comparison of media and techniques for counting *Azotobacter* in soil. *Pl. Soil* **17**, 308–319.
- BURLINGHAM, S. H. (1960) Unpublished results.
- CRUMP, L. M. (1920) Numbers of Protozoa in certain Rothamsted soils. *J. agric. Sci., Camb.* **10**, 182–198.
- CUTLER, D. W. & CRUMP, L. M. (1920) Daily periodicity in the numbers of active soil flagellates; with a brief note on the relation of trophic amoebae and bacterial numbers. *Ann. appl. Biol.* **7**, 11–24.
- ETHERIDGE, J. (1960) Unpublished results.
- ETHERIDGE, J. (1964) Unpublished results.
- FLENTJE, N. J. (1956) Studies on *Pellicularia filamentosa* (Pat) Rogers. I Formation of the perfect stage. *Trans. Br. mycol. Soc.* **39**, 343–356.
- HARLEY, J. L. & WAID, J. S. (1955) A method of studying active mycelia on living roots and other surfaces in the soil. *Trans. Br. mycol. Soc.* **38**, 104–118.
- HAYMAN, D. (1968) Unpublished results.
- HUTCHINSON, H. B. & CLAYTON, J. (1919) On the decomposition of cellulose by an aerobic organism (*Spirochatea cytophaga* n.sp.). *J. agric. Sci., Camb.* **9**, 143–173.
- JOHNSON, L. F. *et al.* (1959) *Methods for studying soil microflora—plant disease relationships*. Minneapolis; Burgess Publishing Co.
- LINGAPPA, Y. & LOCKWOOD, J. L. (1962) Chitin media for selective isolation and culture of actinomycetes. *Phytopathology* **52**, 317–323.
- MARTIN, C. H. & LEWIN, K. R. (1915) Notes on some methods for the estimation of soil protozoa. *J. agric. Sci.* **7**, 106–119.
- MEIKLEJOHN, J. (1949) Isolation of *Nitrosomonas* from Rothamsted soil. *Nature, Lond.* **164**, 667.
- MEIKLEJOHN, J. (1956) Preliminary notes on numbers of nitrogen fixers on Broadbalk field. *Proc. 6th int. Cong. Soil Sci., Paris, C*, 243–248.
- MEIKLEJOHN, J. (1962) Unpublished results.
- MEIKLEJOHN, J. (1965) *Azotobacter* numbers on Broadbalk field, Rothamsted. *Pl. Soil* **23**, 227–235.
- MOSSE, B. & BOWEN, G. D. (1968a) A key to the recognition of some *Endogone* spore types. *Trans. Br. mycol. Soc.* **51**, 469.
- MOSSE, B. & BOWEN, G. D. (1968b) The distribution of *Endogone* spores in some Australian and New Zealand soils, and in an experimental field soil at Rothamsted. *Trans. Br. mycol. Soc.* **51**, 485.
- NUTMAN, P. S. (1963) *Rep. Rothamsted exp. Stn for 1962*, 80, 81.
- NUTMAN, P. S. (1968) Unpublished results.
- RUSSELL, E. J. & APPLEYARD, A. (1917) The influence of soil conditions on the decomposition of organic matter in the soil. *J. agric. Sci.* **8**, 385–417.
- RUSSELL, E. J. & HUTCHINSON, H. B. (1913) The effect of partial sterilisation of soil on the production of plant food. II. The limitation of bacterial numbers in normal soils and its consequences. *J. Agric. Sci., Camb.*, **5**, 152–221.

MICROBIOLOGY OF BROADBALK SOILS

- SINGH, J. (1937) Soil fungi and actinomycetes in relation to manurial treatment, season and crop. *Ann. appl. Biol.* **24**, 154-168.
- SINGH, B. N. (1946) A method of estimating the numbers of soil protozoa, especially amoebae, based on their differential feeding on bacteria. *Ann. appl. Biol.* **33**, 112-119.
- SINGH, B. N. (1947a) Myxobacteria in soils and composts, their distribution, number and lytic action on bacteria. *J. gen. Microbiol.* **1**, 1-10.
- SINGH, B. N. (1947b) Studies on soil Acrasidae. 1. Distribution of species of *Dictyostelium* in soils of Great Britain, and the effect of bacteria on their development. *J. gen. Microbiol.* **1**, 11-21.
- SINGH, B. N. (1948) Studies on giant amoeboid organisms. 1. The distribution of *Leptomyxa reticulata* Goodey in soils of Great Britain, and the effect of bacterial food on growth and cyst formation. *J. gen. Microbiol.* **2**, 8-14.
- SINGH, B. N. (1949) The effect of artificial fertilizers and dung on the numbers of amoebae in Rothamsted soils. *J. gen. Microbiol.* **3**, 204-210.
- SKINNER, F. A. (1951) A method for distinguishing between viable spores and mycelial fragments of actinomycetes in soils. *J. gen. Microbiol.* **5**, 159-166.
- SKINNER, F. A., JONES, P. C. T. & MOLLISON, J. E. (1952) A comparison of a direct and a plate-counting technique for the quantitative estimation of soil micro-organisms. *J. gen. Microbiol.* **6**, 261-271.
- THORNTON, H. G. & CRUMP, L. M. (1953) Micro-predators in soil. *Rep. Rothamsted exp. Stn for 1952*, 164-172.
- ZIEMIECKA, J. (1932) The *Azotobacter* test of soil fertility applied to the classical fields at Rothamsted. *J. agric. Sci., Camb.*, **22**, 797-810.