

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Rothamsted Experimental Station Report for 1977 Part 2

ROTHAMSTED
EXPERIMENTAL STATION
REPORT FOR 1977
PART 2
PUBLISHED JUNE 1978
ROTHAMSTED RESEARCH
LAWES AGRICULTURAL TRUST

[Full Table of Content](#)

Use of Fertilisers in England and Wales, 1977

B. M. Church

B. M. Church (1978) *Use of Fertilisers in England and Wales, 1977* ; Rothamsted Experimental Station Report For 1977 Part 2, pp 73 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-36543>

Use of Fertilisers in England and Wales, 1977

B. M. CHURCH

In this continuing series of annual surveys done by ADAS Soil Scientists and representatives of the Fertiliser Manufacturers' Association in collaboration with the Statistics Department, Rothamsted (Church & Webber, 1971, Church & Lewis, 1978) a representative sample of 1335 farms was surveyed in England and Wales in 1977.

The main feature of the 1977 results is a marked increase in the use of N on both leys and permanent grass. Use of P and K has changed little.

For nearly 10 years, N use on grassland has been increasing at an average rate of about 7½% per year. This increase was checked by the drought in the summer of 1976, but the upward trend was fully resumed in 1977 when N use on grassland was about 15% more than in 1975 (Table 1).

After little change for several years, there was evidence in both 1976 and 1977 of an overall increase in N use per hectare on tillage crops. Although there may have been changes on other individual crops, the most notable increase is on winter wheat where, possibly due to the introduction of new varieties, N use is estimated to have increased by nearly a quarter in the last 2 years to an average of 115 kg ha⁻¹ (Table 2). As a general average for winter wheat, this would seem to be adequate compared with current recommendations. However, about a tenth of the total N used on this crop is still applied in autumn when it may be of little benefit.

TABLE 1
Fertiliser use on tillage, leys and permanent grass, 1974–77 (kg ha⁻¹)

	Tillage			Ley			P.G. [†]			All crops & grass		
	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
1974	85	51	56	136	36	30	71	23	16	92	40	38
1975	87	46	51	142	34	28	77	20	16	96	35	34
1976	96	50	56	144	36	33	74	22	14	99	39	37
1977*	100	49	56	166	33	32	88	21	16	111	37	39

† Includes grass seeded more than 7 years ago.

* Estimates for previous years adjusted to be comparable with those for 1977.

TABLE 2
Fertiliser use on winter wheat (kg ha⁻¹) 1975–77

N	P ₂ O ₅			K ₂ O		
	1975	1976	1977	1975	1976	1977
93	102	115	39	42	40	33

Estimates of average fertiliser use in 1977 and of the proportions of crop area getting different amounts of nutrients are given for major crops in Tables 3–6. Table 7 gives similar information for grassland classified according to the number of years since seeding, so that use can be related to the revised categories in the Agricultural Census. More detailed information for grassland and for horticultural crops will be reported elsewhere.

REFERENCES

- CHURCH, B. M. & WEBBER, J. (1971) Fertiliser practice in England and Wales: a new series of surveys. *Journal of the Science of Food and Agriculture* 22, 1–7.
CHURCH, B. M. & LEWIS, D. A. (1977) Fertiliser use on farm crops in England and Wales: Information from the Survey of Fertiliser Practice 1942–1976. *Outlook on Agriculture* 9, 186–193.

ROTHAMSTED REPORT FOR 1977, PART 2

TABLE 3
Fertiliser use in England and Wales, 1977

Fields	Hectares (000)	Overall* (kg ha ⁻¹)			% Area receiving			Actual* (kg ha ⁻¹)		
		N	P ₂ O ₅	K ₂ O	N	P	K	FYM	N	P ₂ O ₅
Spring wheat	131	53	86	30	28	92	77	76	94	39
Winter wheat	1947	935	115	40	33	98	77	11	116	52
Spring barley	3071	1458	82	36	37	98	94	17	83	39
Winter barley	610	258	104	44	41	100	91	85	104	49
Spring oats	212	58	67	34	32	96	92	23	70	37
Winter oats	217	64	86	47	42	98	88	11	88	53
Mixed corn	37	13	50	37	33	93	90	86	54	41
Rye	41	10	82	27	31	96	66	20	85	41
Maize	40	10	114	59	57	99	93	87	50	116
Early potatoes	116	29	170	178	216	97	97	50	176	184
Maincrop potatoes	408	128	181	186	250	100	100	51	181	186
Sugar beet	484	222	141	71	157	97	96	33	145	73
Swedes (stock)	121	22	60	127	85	87	96	48	69	132
Turnips (stock)	81	17	62	66	53	90	83	84	42	69
Mangolds	51	5	119	96	134	98	98	62	121	97
Kale and cow cabbage	270	51	110	49	53	95	82	81	116	65
Rape (stock)	62	16	83	112	41	95	94	87	119	46
Beans (stock)	97	33	33	8	33	19	17	42	34	45
Mixed roots and green crops	57	16	82	69	78	92	91	91	68	89
Peas—vining	87	43	7	21	26	20	47	49	4	35
Peas—harvested dry	88	36	6	24	25	24	52	49	1	24
French beans	24	4	151	84	73	95	90	90	0	160
Brussels sprouts	27	6	259	111	164	100	100	100	2	259
Cabbage	68	13	144	68	125	99	96	7	145	71
Cauliflower	60	9	148	84	162	100	89	89	148	95
Carrots	34	9	61	62	77	77	74	4	78	84
Onions	39	8	150	129	188	100	100	15	150	129
Oilseed rape	108	55	238	53	39	100	86	58	3	238
Cereals for grazing	27	8	83	19	19	96	49	27	86	39
Forage maize	59	17	104	58	62	95	95	92	105	61
Arable silage	64	19	78	38	48	97	90	39	81	42
Lucerne	25	7	13	84	163	20	77	85	22	66
All tillage	9298	3776	100	49	56	95	86	82	106	56
One year leys	74	22	134	33	37	94	61	62	143	53
Two to seven year leys	3557	1642	165	33	32	95	67	66	175	49
Permanent grass	3502	2253	88	21	16	74	50	47	119	42
All crops and grass	16431	7692	111	37	39	89	72	69	125	57

* The average application of any fertiliser component over all fields, including those receiving none, is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.

USE OF FERTILISERS IN ENGLAND AND WALES, 1977

TABLE 4

	Percentages of crop area getting different amounts of N (kg ha^{-1})											
Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	131	8	1	2	22	27	24	12	3	0	1	0
Winter wheat	1947	2	0	3	8	19	24	28	15	1	0	0
Spring barley	3071	2	1	7	24	43	17	5	0	0	0	0
Winter barley	610	0	1	2	8	30	36	17	5	0	0	0
Spring oats	212	4	1	11	41	32	7	1	2	0	0	0
Winter oats	217	2	2	8	24	27	24	11	2	0	0	0
Mixed corn	37	11	24	37	15	1	3	3	0	0	0	0
Rye	41	4	11	6	8	49	11	6	5	0	0	0
Maize	40	1	3	0	2	15	33	33	13	0	0	0
Early potatoes	116	3	3	0	1	1	1	3	15	49	17	6
Maincrop potatoes	408	0	1	1	1	1	2	5	8	54	19	3
Sugar beet	484	3	0	1	1	8	12	38	29	5	2	1
Swedes (stock)	121	13	6	22	21	20	10	2	6	0	0	0
Turnips (stock)	81	10	6	21	24	21	4	10	2	1	0	0
Mangolds	51	2	0	8	13	16	15	16	23	4	2	1
Kale and cow cabbage	270	5	1	6	10	19	18	22	14	5	1	0
Rape (stock)	62	5	0	7	13	39	25	4	7	0	0	0
Beans (stock)	97	83	14	0	0	0	0	0	1	1	0	0
Mixed roots and green crops	57	8	3	12	25	18	8	12	8	5	0	0
Peas—vining	87	80	4	10	5	1	0	0	0	0	0	0
Peas—harvested dry	88	76	18	2	4	0	0	0	0	0	0	0
French beans	24	5	0	0	0	0	17	9	40	24	5	0
Brussels sprouts	27	0	0	3	0	8	0	0	4	10	15	2
Cabbage	68	1	9	5	5	17	8	10	13	15	42	33
Cauliflower	60	0	1	2	28	11	6	5	20	15	1	0
Carrots	34	23	5	4	19	30	13	5	2	0	0	0
Onions	39	0	0	0	0	19	26	10	30	2	7	6
Oilseed rape	108	0	0	0	1	4	0	1	11	40	34	7
Cereals for grazing	27	4	1	0	18	39	2	18	4	0	0	0
Forage maize	59	1	0	0	10	25	31	28	4	1	0	0
Arable silage	64	3	2	9	38	23	11	6	5	3	0	2
Lucerne	25	80	3	6	9	0	0	0	0	0	0	2
All tillage	9298	5	1	5	15	27	18	14	10	3	1	0
One year leys	74	6	0	2	20	15	8	12	18	4	9	6
Two to seven year leys	3557	5	1	5	14	13	5	10	13	10	9	5
Permanent grass	3502	26	2	9	17	15	5	8	7	4	3	3
All crops and grass	16431	11	1	6	15	20	11	12	10	5	3	1

ROTHAMSTED REPORT FOR 1977, PART 2

	Percentages of crop area getting different amounts of P_2O_5 ($kg\ ha^{-1}$)											
Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	131	23	6	57	12	1	0	1	0	0	0	0
Winter wheat	1947	23	3	39	29	4	1	0	1	0	0	0
Spring barley	3071	6	8	72	12	1	0	0	0	0	0	0
Winter barley	610	9	3	50	32	4	1	1	0	0	0	0
Spring oats	212	8	10	73	1	1	0	1	1	0	0	0
Winter oats	217	12	4	41	31	6	3	2	1	0	0	0
Mixed corn	37	10	27	39	9	10	0	4	1	0	0	0
Rye	41	34	20	22	23	0	1	0	0	0	0	0
Maize	40	7	0	20	46	14	10	3	0	0	0	0
Early potatoes	116	3	3	0	1	4	6	12	38	17	5	0
Maincrop potatoes	408	0	2	0	1	3	3	7	51	22	5	0
Sugar beet	484	4	2	16	39	24	8	2	4	1	0	0
Swedes (stock)	121	4	1	5	13	19	17	15	9	12	3	0
Turnips (stock)	81	17	3	30	19	6	22	16	14	12	1	0
Mangolds	51	2	6	14	12	22	12	3	3	0	0	0
Kale and cow cabbage	270	18	3	31	28	12	12	3	0	0	0	0
Rape (stock)	62	6	0	27	29	1	1	7	6	13	4	0
Beans (stock)	97	58	0	14	18	2	0	0	0	5	2	0
Mixed roots and green crops	57	9	9	19	27	16	6	5	4	3	2	0
Peas—vining	87	53	3	22	17	5	0	0	0	0	0	0
Peas—harvested dry	88	48	5	25	20	0	0	2	2	18	1	0
French beans	24	10	0	18	9	17	26	3	15	21	0	4
Brussels sprouts	27	0	1	14	5	37	3	0	4	0	0	0
Cabbage	68	4	17	7	20	36	12	0	7	0	1	0
Cauliflower	60	11	1	9	9	26	27	7	7	0	0	0
Carrots	34	26	0	12	13	30	3	12	3	46	11	0
Onions	39	0	6	1	11	8	16	2	5	0	0	0
Oilseed rape	108	14	0	16	55	7	2	5	0	0	0	0
Cereals for grazing	27	51	18	20	8	0	4	0	0	0	0	0
Forage maize	59	5	5	20	58	6	3	0	3	0	0	0
Arable silage	64	10	27	29	21	12	0	0	0	0	0	0
Lucerne	25	23	6	3	8	9	35	6	10	10	1	1
All tillage	9298	14	6	47	21	5	2	1	3	1	0	0
One year leys	74	39	5	29	13	10	1	1	2	1	0	0
Two to seven year leys	3557	33	13	34	10	4	2	2	2	1	0	0
Permanent grass	3502	50	15	25	4	4	1	1	3	1	0	0
All crops and grass	16431	28	10	38	13	4	3	1	1	1	0	0

USE OF FERTILISERS IN ENGLAND AND WALES, 1977

TABLE 6
Percentages of crop area getting different amounts of K_2O ($kg\ ha^{-1}$)

Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	131	24	6	59	10	2	0	0	0	0	0	0
Winter wheat	1947	31	3	40	20	4	1	0	0	0	0	0
Spring barley	3071	9	6	66	17	1	0	0	0	0	0	0
Winter barley	610	15	1	54	25	3	1	1	0	0	0	0
Spring oats	2122	12	8	68	10	1	0	0	0	0	0	0
Winter oats	217	17	3	38	32	8	2	0	0	0	0	0
Mixed corn	37	14	23	38	12	13	0	0	0	0	0	0
Rye	41	34	8	38	16	3	1	0	0	0	0	0
Maize	40	13	0	16	46	10	12	3	0	0	0	0
Early potatoes	116	3	3	0	0	1	4	9	21	17	26	13
Maincrop potatoes	408	0	1	1	0	1	4	4	8	25	29	23
Sugar beet	484	1	0	0	10	12	14	7	25	20	8	2
Swedes (stock)	121	8	2	10	18	24	24	6	7	0	0	0
Turnips (stock)	81	16	7	28	25	9	7	0	1	0	0	0
Mangolds	51	2	1	4	12	14	10	18	20	16	5	0
Kale and cow cabbage	270	19	2	28	28	12	4	4	3	0	0	0
Rape (stock)	62	13	2	43	37	5	0	0	0	0	0	0
Beans (stock)	97	66	0	14	14	3	3	0	0	0	0	0
Mixed roots and green crops	57	9	6	17	20	21	11	5	8	3	2	0
Peas—vining	87	51	1	19	22	5	0	0	2	0	0	0
Peas—harvested dry	88	51	1	21	20	3	2	0	0	0	0	0
French beans	24	10	0	3	45	17	24	0	1	0	0	0
Brussels sprouts	27	0	0	0	28	3	4	2	25	24	4	11
Cabbage	68	4	9	2	14	11	14	4	22	13	4	1
Cauliflower	60	11	0	5	8	3	5	1	35	11	18	2
Carrots	34	26	0	5	15	21	5	17	7	5	0	0
Onions	39	0	0	5	3	6	9	13	15	4	42	4
Oilseed rape	108	42	0	10	34	7	2	5	0	0	0	0
Cereals for grazing	27	51	18	24	2	0	6	0	0	0	0	0
Forage maize	59	8	3	20	52	9	2	0	2	2	0	0
Arable silage	64	10	6	26	41	15	0	2	0	0	0	0
Lucerne	25	15	6	3	3	10	6	1	14	35	0	2
All tillage	9298	18	4	45	18	4	2	1	3	3	2	1
One year leys	74	38	3	23	18	12	5	0	2	0	0	0
Two to seven year leys	3557	34	11	31	12	6	3	2	1	0	0	0
Permanent grass	3502	53	14	26	6	1	1	0	0	0	0	0
All crops and grass	16431	31	8	36	13	4	2	1	2	1	1	1

ROTHAMSTED REPORT FOR 1977, PART 2

TABLE 7
Fertiliser use on grassland by years since seeded

Fields	Hectares ('000)	Overall* (kg ha ⁻¹)			% Area receiving			Actual* (kg ha ⁻¹)		
		N	P ₂ O ₅	K ₂ O	N	P	K	FYM	N	P ₂ O ₅
1 year leys	74	23	136	32	36	94	59	60	28	144
Other grassland	849	295	160	44	39	94	73	71	28	171
First year	770	305	180	33	35	96	68	68	30	188
Second year	1619	600	170	38	37	95	70	70	29	180
One or two years	617	318	162	27	31	94	62	64	47	172
Third year	437	232	169	27	29	95	65	64	45	178
Fourth year	351	218	171	34	33	96	70	69	39	177
Fifth year	1405	769	167	29	31	95	65	66	44	175
Three to five years	3103	1412	168	33	33	95	67	67	37	177
One to five years	405	236	143	29	23	93	65	57	51	153
Six to seven years	3942	2640	91	21	17	73	50	47	32	124
Over five years	3508	1648	164	32	32	95	67	66	39	173
One to seven years	3537	2404	85	20	16	71	48	46	30	120
Over seven years	7119	4075	118	25	22	81	56	54	33	145
All grassland										42

* The average application of any fertiliser component over all fields, including those receiving none, is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.