


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Rothamsted Experimental Station Report for 1984

[Full Table of Content](#)



Use of Fertilizers in England and Wales, 1984

B. M. Church

B. M. Church (1985) *Use of Fertilizers in England and Wales, 1984* ; Rothamsted Experimental Station Report For 1984, pp 277 - 284 - DOI: <https://doi.org/10.23637/ERADOC-1-25>

Use of fertilizers in England and Wales, 1984

B. M. CHURCH

The series of annual surveys done by staff of the ADAS Regional Soil Scientists and representatives of the Fertiliser Manufacturers' Association in collaboration with Rothamsted was continued in 1984, when a sample of 1352 farms was surveyed in England and Wales (Church & Lewis, 1977). It may perhaps be noted here that similar surveys of fertilizer practice were done on samples of 250 farms in Scotland in 1983 and in 1984 by representatives of the Fertiliser Manufacturers' Association, and that the 1983 results have been reported elsewhere (Church & Leech, 1984).

Compared with 1983, the survey data for England and Wales show an estimated increase of 6% in N use per hectare crops and grass, with corresponding increases of almost 8 and 14% in P and K per hectare. There have been average increases over the last four years of about 5, 3 and 6% per year in use of N, P and K per hectare crops and grass (Table 1). Straight N use per hectare has increased by a third since 1980 at an average of about 7.5% per year but use of N in compounds has changed little over this period.

TABLE 1
*Fertilizer use on tillage crops and grassland (kg ha⁻¹), 1980-84**

| | Tillage crops | | | | Grassland | | | | All crops and grass | | | |
|-------------------------------|---------------|------|------|------|-----------|------|------|------|---------------------|------|------|------|
| | 1980 | 1982 | 1983 | 1984 | 1980 | 1982 | 1983 | 1984 | 1980 | 1982 | 1983 | 1984 |
| N Straight | 77 | 99 | 116 | 128 | 69 | 71 | 69 | 71 | 73 | 85 | 91 | 99 |
| Compound | 44 | 42 | 37 | 34 | 50 | 52 | 57 | 61 | 47 | 47 | 48 | 48 |
| Total | 121 | 141 | 154 | 162 | 119 | 123 | 126 | 132 | 120 | 132 | 139 | 147 |
| P ₂ O ₅ | 49 | 55 | 54 | 61 | 27 | 24 | 26 | 25 | 37 | 39 | 39 | 42 |
| K ₂ O | 54 | 61 | 60 | 68 | 26 | 28 | 28 | 33 | 40 | 44 | 44 | 50 |

* Comparable data for 1981 are in the 1983 Rothamsted Report

Following the pattern of recent years, the increase in use of straight N between 1983 and 1984 was mainly on tillage crops, on which use of N in compounds continued to decline. These changes are partly explained by the continuing move from spring to winter sown cereals (about a quarter spring-sown in 1983, declining to about 15% in 1984). On grassland, use of straight N has hardly changed over the last four years but N use in compounds has increased by 10% a year since 1982. The increase in use of P between 1983 and 1984 was on tillage crops only, as has been the case since 1980, but K use continued to increase on both tillage crops and grassland.

On winter wheat, use of straight N per hectare was at least maintained, and use of both P and K is estimated to have increased sharply in 1984 (Table 2). On spring barley N use was at

TABLE 2
Fertilizer use on winter wheat and spring barley (kg ha⁻¹), 1980-84

| | Winter wheat | | | | Spring barley | | | |
|-------------------------------|--------------|------|------|------|---------------|------|------|------|
| | 1980 | 1982 | 1983 | 1984 | 1980 | 1982 | 1983 | 1984 |
| N Straight | 126 | 148 | 167 | 171 | 24 | 34 | 51 | 44 |
| Compound | 19 | 18 | 16 | 16 | 63 | 60 | 57 | 54 |
| Total | 145 | 162 | 182 | 187 | 87 | 94 | 107 | 98 |
| P ₂ O ₅ | 46 | 49 | 51 | 56 | 37 | 38 | 39 | 39 |
| K ₂ O | 39 | 42 | 46 | 53 | 40 | 41 | 44 | 44 |

ROTHAMSTED REPORT FOR 1984, PART 2

TABLE 3
Fertilizer use in England and Wales, 1984

| Fields | Hectares ('000) | Overall* (kg ha ⁻¹) | | | % Area receiving | | | Actual* (kg ha ⁻¹) | | | |
|----------------------------|--------------------|---------------------------------|-------------------------------|------------------|------------------|-----|-----|--------------------------------|-----|-------------------------------|------------------|
| | | N | P ₂ O ₅ | K ₂ O | N | P | K | FYM | N | P ₂ O ₅ | K ₂ O |
| Spring wheat | 35 | 21 | 132 | 26 | 28 | 98 | 56 | 56 | 135 | 47 | 51 |
| Winter wheat | 2867 | 1807 | 187 | 56 | 53 | 99 | 83 | 11 | 189 | 63 | 63 |
| Spring barley | 480 | 98 | 39 | 44 | 98 | 91 | 93 | 24 | 100 | 42 | 47 |
| Winter barley | 1127 | 921 | 150 | 57 | 59 | 100 | 95 | 13 | 150 | 61 | 64 |
| Spring oats | 66 | 14 | 72 | 33 | 35 | 91 | 84 | 25 | 79 | 39 | 42 |
| Winter oats | 149 | 57 | 114 | 56 | 59 | 98 | 97 | 11 | 116 | 58 | 60 |
| Rye | 40 | 10 | 123 | 49 | 49 | 98 | 94 | 3 | 126 | 42 | 52 |
| Maize | 25 | 6 | 89 | 35 | 49 | 81 | 72 | 80 | 53 | 110 | 49 |
| Early potatoes | 67 | 20 | 185 | 202 | 220 | 100 | 100 | 58 | 185 | 202 | 220 |
| Maincrop potatoes | 335 | 132 | 214 | 228 | 278 | 99 | 99 | 40 | 216 | 230 | 281 |
| Sugar beet | 414 | 206 | 148 | 74 | 160 | 98 | 94 | 28 | 150 | 79 | 164 |
| Oilseed rape | 422 | 259 | 279 | 63 | 60 | 99 | 97 | 5 | 281 | 65 | 68 |
| Swedes (stock) | 68 | 12 | 56 | 93 | 79 | 99 | 97 | 50 | 58 | 94 | 82 |
| Turnips (stock) | 54 | 12 | 85 | 62 | 56 | 97 | 74 | 74 | 88 | 83 | 76 |
| Kale and cow cabbage | 66 | 15 | 112 | 45 | 49 | 94 | 86 | 87 | 120 | 52 | 57 |
| Rape for stockfeed | 43 | 11 | 75 | 104 | 46 | 99 | 99 | 26 | 76 | 105 | 46 |
| Beans for stockfeed | 84 | 27 | 3 | 38 | 41 | 14 | 61 | 59 | 24 | 63 | 69 |
| Other stockfeed | 82 | 25 | 77 | 51 | 62 | 85 | 93 | 94 | 90 | 55 | 65 |
| Peas for human consumption | 156 | 79 | 3 | 26 | 32 | 15 | 49 | 51 | 7 | 21 | 53 |
| Broad beans | 25 | 5 | 2 | 75 | 109 | 2 | 71 | 69 | 4 | 84 | 106 |
| Runner and French beans | 34 | 10 | 107 | 56 | 94 | 81 | 84 | 4 | 133 | 73 | 113 |
| Brussels sprouts | 35 | 7 | 257 | 78 | 193 | 97 | 94 | 8 | 266 | 84 | 206 |
| Cabbages | 46 | 9 | 202 | 71 | 150 | 90 | 91 | 7 | 226 | 78 | 157 |
| Cauliflower | 33 | 8 | 222 | 82 | 190 | 98 | 98 | 12 | 226 | 83 | 194 |
| Onions | 33 | 6 | 147 | 105 | 148 | 99 | 94 | 98 | 149 | 112 | 151 |
| Small fruit | 54 | 9 | 93 | 25 | 49 | 84 | 69 | 77 | 12 | 111 | 36 |
| Top fruit | 110 | 41 | 55 | 19 | 39 | 74 | 46 | 45 | 9 | 74 | 40 |
| All tillage | 8385 | 4290 | 162 | 61 | 68 | 96 | 90 | 87 | 16 | 169 | 68 |
| 2-7 year leys | 2954 | 1510 | 190 | 33 | 51 | 96 | 77 | 77 | 48 | 198 | 43 |
| Permanent grass | 3666 | 2937 | 102 | 20 | 23 | 80 | 60 | 60 | 38 | 128 | 34 |
| All crops and grass | 15017 | 8739 | 147 | 42 | 50 | 91 | 77 | 76 | 29 | 162 | 55 |

*The average application of any fertilizer component over all fields including those receiving none is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.

USE OF FERTILIZERS IN ENGLAND AND WALES, 1984

TABLE 4
Percentages of crop area getting different amounts of N (kg ha⁻¹)

| | Fields | 0 | <25 | 25- | 50- | 75- | 100- | 125- | 150- | 200- | 250- | 300- | 400+ |
|----------------------------|--------|----|-----|-----|-----|-----|------|------|------|------|------|------|------|
| Spring wheat | 35 | 2 | 0 | 7 | 0 | 9 | 11 | 44 | 24 | 3 | 0 | 0 | 0 |
| Winter wheat | 2867 | 1 | 1 | 1 | 1 | 2 | 3 | 10 | 40 | 33 | 8 | 0 | 0 |
| Spring barley | 1127 | 2 | 1 | 7 | 23 | 18 | 26 | 16 | 6 | 1 | 0 | 0 | 0 |
| Winter barley | 1589 | 0 | 2 | 1 | 2 | 6 | 11 | 27 | 42 | 7 | 1 | 0 | 0 |
| Spring oats | 66 | 9 | 3 | 13 | 32 | 21 | 17 | 4 | 0 | 0 | 0 | 0 | 0 |
| Winter oats | 149 | 2 | 4 | 6 | 11 | 15 | 14 | 22 | 26 | 0 | 0 | 0 | 0 |
| Rye | 40 | 2 | 3 | 0 | 5 | 6 | 27 | 37 | 19 | 1 | 0 | 0 | 0 |
| Maize | 25 | 19 | 0 | 6 | 1 | 22 | 27 | 13 | 11 | 0 | 0 | 0 | 0 |
| Early potatoes | 67 | 0 | 0 | 4 | 0 | 6 | 13 | 6 | 35 | 22 | 7 | 4 | 3 |
| Maincrop potatoes | 335 | 1 | 0 | 1 | 1 | 3 | 3 | 6 | 24 | 25 | 5 | 0 | 0 |
| Sugar beet | 414 | 2 | 0 | 0 | 0 | 3 | 7 | 19 | 35 | 22 | 7 | 3 | 0 |
| Oilseed rape | 422 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 12 | 44 | 38 |
| Swedes (stock) | 68 | 3 | 5 | 43 | 33 | 13 | 2 | 1 | 0 | 1 | 0 | 0 | 0 |
| Turnips (stock) | 54 | 3 | 1 | 17 | 30 | 13 | 24 | 8 | 4 | 0 | 0 | 0 | 0 |
| Kale and cow cabbage | 66 | 6 | 1 | 4 | 13 | 17 | 21 | 13 | 19 | 5 | 0 | 0 | 0 |
| Rape for stockfeed | 43 | 1 | 1 | 25 | 38 | 21 | 5 | 4 | 0 | 0 | 0 | 0 | 0 |
| Beans for stockfeed | 84 | 86 | 11 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Other stockfeed | 82 | 15 | 4 | 10 | 24 | 12 | 11 | 20 | 5 | 0 | 0 | 0 | 0 |
| Peas for human consumption | 156 | 85 | 13 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Broad beans | 25 | 98 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Runner and French beans | 34 | 19 | 10 | 3 | 0 | 0 | 3 | 13 | 19 | 31 | 0 | 1 | 0 |
| Brussels sprouts | 35 | 3 | 0 | 2 | 0 | 0 | 0 | 3 | 0 | 26 | 16 | 7 | 33 |
| Cabbages | 46 | 10 | 0 | 0 | 0 | 0 | 2 | 4 | 3 | 29 | 23 | 11 | 18 |
| Cauliflower | 33 | 2 | 0 | 1 | 0 | 0 | 3 | 11 | 23 | 7 | 20 | 19 | 7 |
| Onions | 33 | 1 | 3 | 1 | 0 | 0 | 11 | 18 | 21 | 21 | 0 | 2 | 0 |
| Small fruit | 54 | 16 | 13 | 7 | 9 | 6 | 14 | 4 | 29 | 1 | 0 | 0 | 0 |
| Top fruit | 110 | 26 | 7 | 30 | 10 | 12 | 1 | 5 | 5 | 3 | 0 | 1 | 0 |
| All tillage | 8385 | 4 | 1 | 3 | 5 | 6 | 9 | 15 | 30 | 18 | 7 | 3 | 0 |
| 2-7 year leys | 2954 | 4 | 1 | 8 | 7 | 8 | 7 | 14 | 13 | 11 | 5 | 5 | 5 |
| Permanent grass | 3666 | 20 | 3 | 15 | 13 | 11 | 7 | 9 | 5 | 15 | 5 | 4 | 1 |
| All crops and grass | 15017 | 9 | 2 | 8 | 8 | 8 | 8 | 11 | 20 | 13 | 6 | 7 | 1 |

ROTHAMSTED REPORT FOR 1984, PART 2

TABLE 5

| | Percentages of crop area getting different amounts of P_2O_5 ($kg\ ha^{-1}$) | | | | | | | | | | | |
|----------------------------|--|-----|-----|-----|-----|------|------|------|------|------|------|------|
| Fields | 0 | <25 | 25- | 50- | 75- | 100- | 125- | 150- | 200- | 250- | 300- | 400+ |
| Spring wheat | 35 | 44 | 7 | 32 | 17 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| Winter wheat | 2867 | 12 | 2 | 18 | 53 | 13 | 2 | 0 | 0 | 0 | 0 | 0 |
| Spring barley | 1127 | 9 | 15 | 50 | 24 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Winter barley | 1589 | 5 | 2 | 22 | 57 | 10 | 2 | 0 | 0 | 0 | 0 | 0 |
| Spring oats | 66 | 16 | 16 | 52 | 15 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| Winter oats | 149 | 3 | 3 | 30 | 55 | 6 | 0 | 2 | 0 | 0 | 0 | 0 |
| Rye | 40 | 6 | 28 | 36 | 30 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Maize | 25 | 28 | 0 | 38 | 30 | 4 | 0 | 0 | 0 | 0 | 0 | 0 |
| Early potatoes | 67 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 11 | 39 | 28 | 5 |
| Maincrop potatoes | 335 | 1 | 0 | 1 | 1 | 3 | 3 | 7 | 10 | 24 | 20 | 2 |
| Sugar beet | 414 | 6 | 3 | 26 | 32 | 10 | 7 | 6 | 1 | 1 | 0 | 3 |
| Oilseed rape | 422 | 3 | 1 | 11 | 72 | 9 | 3 | 0 | 0 | 0 | 0 | 0 |
| Swedes (stock) | 68 | 1 | 8 | 12 | 16 | 25 | 6 | 6 | 0 | 1 | 1 | 0 |
| Turnips (stock) | 54 | 26 | 10 | 22 | 13 | 5 | 13 | 5 | 4 | 0 | 4 | 0 |
| Kale and cow cabbage | 66 | 14 | 8 | 46 | 28 | 2 | 0 | 0 | 0 | 2 | 0 | 0 |
| Rape for stockfeed | 43 | 1 | 16 | 20 | 5 | 8 | 13 | 6 | 23 | 6 | 0 | 0 |
| Beans for stockfeed | 84 | 39 | 5 | 16 | 31 | 3 | 1 | 2 | 0 | 1 | 0 | 0 |
| Other stockfeed | 82 | 7 | 20 | 31 | 32 | 1 | 2 | 5 | 2 | 1 | 0 | 0 |
| Peas for human consumption | 156 | 51 | 3 | 20 | 18 | 6 | 1 | 0 | 0 | 0 | 0 | 0 |
| Broad beans | 25 | 29 | 0 | 1 | 20 | 21 | 9 | 9 | 2 | 0 | 4 | 0 |
| Runner and French beans | 34 | 24 | 8 | 17 | 26 | 9 | 10 | 10 | 11 | 0 | 0 | 0 |
| Brussels sprouts | 35 | 6 | 2 | 23 | 29 | 9 | 12 | 14 | 12 | 2 | 0 | 0 |
| Cabbages | 46 | 9 | 8 | 23 | 21 | 40 | 23 | 14 | 7 | 2 | 3 | 0 |
| Cauliflower | 33 | 2 | 8 | 2 | 48 | 5 | 7 | 6 | 22 | 15 | 1 | 0 |
| Onions | 33 | 6 | 17 | 6 | 2 | 0 | 1 | 0 | 0 | 1 | 1 | 1 |
| Small fruit | 54 | 31 | 15 | 18 | 24 | 3 | 0 | 1 | 0 | 1 | 1 | 0 |
| Top fruit | 110 | 54 | 18 | 24 | 3 | 0 | 1 | 0 | 0 | 1 | 0 | 0 |
| All tillage | 8385 | 10 | 4 | 22 | 46 | 10 | 2 | 1 | 2 | 1 | 1 | 1 |
| 2-7 year leys | 2954 | 23 | 24 | 32 | 13 | 4 | 1 | 1 | 1 | 0 | 0 | 0 |
| Permanent grass | 3666 | 40 | 32 | 19 | 5 | 1 | 1 | 1 | 6 | 0 | 0 | 0 |
| All crops and grass | 15017 | 23 | 17 | 23 | 23 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

USE OF FERTILIZERS IN ENGLAND AND WALES, 1984

TABLE 6
Percentages of crop area getting different amounts of K₂O (kg ha⁻¹)

| | Fields | 0 | <25 | 25- | 50- | 75- | 100- | 125- | 150- | 200- | 250- | 300- | 400+ |
|----------------------------|--------|----|-----|-----|-----|-----|------|------|------|------|------|------|------|
| Spring wheat | 35 | 44 | 5 | 30 | 16 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Winter wheat | 2867 | 17 | 2 | 18 | 46 | 13 | 2 | 1 | 0 | 0 | 0 | 0 | 0 |
| Spring barley | 1127 | 7 | 11 | 43 | 33 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Winter barley | 1589 | 7 | 2 | 17 | 55 | 16 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| Spring oats | 66 | 16 | 16 | 44 | 20 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Winter oats | 149 | 3 | 3 | 32 | 48 | 9 | 1 | 4 | 0 | 0 | 0 | 0 | 0 |
| Rye | 40 | 7 | 1 | 41 | 48 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Maize | 25 | 20 | 0 | 22 | 53 | 2 | 0 | 4 | 0 | 0 | 0 | 0 | 0 |
| Early potatoes | 67 | 0 | 0 | 0 | 1 | 12 | 4 | 9 | 25 | 17 | 39 | 16 | 3 |
| Maincrop potatoes | 335 | 1 | 0 | 1 | 0 | 1 | 1 | 2 | 22 | 15 | 7 | 5 | 2 |
| Sugar beet | 414 | 2 | 1 | 0 | 9 | 14 | 14 | 12 | 12 | 15 | 0 | 0 | 0 |
| Oilseed rape | 422 | 11 | 1 | 4 | 68 | 9 | 3 | 2 | 1 | 0 | 0 | 0 | 0 |
| Swedes (stock) | 68 | 3 | 7 | 13 | 21 | 27 | 22 | 5 | 1 | 0 | 0 | 0 | 0 |
| Turnips (stock) | 54 | 26 | 6 | 17 | 17 | 15 | 15 | 1 | 4 | 0 | 0 | 0 | 0 |
| Kale and cow cabbage | 66 | 13 | 6 | 37 | 30 | 6 | 6 | 1 | 0 | 0 | 0 | 0 | 0 |
| Rape for stockfeed | 43 | 1 | 21 | 51 | 19 | 5 | 2 | 1 | 0 | 0 | 0 | 0 | 0 |
| Beans for stockfeed | 84 | 41 | 3 | 11 | 30 | 3 | 8 | 4 | 0 | 0 | 0 | 0 | 0 |
| Other stockfeed | 82 | 6 | 18 | 30 | 23 | 5 | 3 | 11 | 5 | 0 | 0 | 0 | 0 |
| Peas for human consumption | 156 | 49 | 2 | 7 | 32 | 7 | 1 | 1 | 0 | 0 | 0 | 0 | 0 |
| Broad beans | 25 | 31 | 0 | 2 | 10 | 0 | 0 | 0 | 20 | 32 | 5 | 0 | 0 |
| Runner and French beans | 34 | 16 | 3 | 3 | 35 | 17 | 2 | 1 | 7 | 1 | 5 | 6 | 4 |
| Brussels sprouts | 35 | 6 | 0 | 0 | 0 | 2 | 0 | 0 | 9 | 39 | 9 | 0 | 5 |
| Cabbages | 46 | 4 | 0 | 7 | 2 | 13 | 6 | 15 | 35 | 17 | 0 | 2 | 0 |
| Cauliflower | 33 | 2 | 0 | 0 | 0 | 3 | 7 | 6 | 52 | 28 | 3 | 0 | 0 |
| Onions | 33 | 2 | 0 | 7 | 13 | 2 | 6 | 22 | 29 | 13 | 5 | 0 | 0 |
| Small fruit | 54 | 23 | 11 | 33 | 19 | 3 | 1 | 0 | 16 | 1 | 0 | 0 | 1 |
| Top fruit | 110 | 55 | 5 | 19 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| All tillage | 8385 | 13 | 3 | 18 | 42 | 12 | 3 | 2 | 2 | 2 | 1 | 2 | 0 |
| 2-7 year leys | 2954 | 23 | 17 | 22 | 14 | 7 | 6 | 4 | 2 | 1 | 0 | 0 | 1 |
| Permanent grass | 3666 | 40 | 26 | 20 | 8 | 2 | 3 | 2 | 2 | 1 | 0 | 0 | 0 |
| All crops and grass | 15017 | 24 | 13 | 19 | 26 | 8 | 3 | 2 | 2 | 1 | 0 | 0 | 0 |

ROTHAMSTED REPORT FOR 1984, PART 2

TABLE 7
Fertilizer use on grassland classified by utilization[†]

| Fields | % Grassland area | Overall* (kg ha ⁻¹) | | | % Area receiving | | | Actual* (kg ha ⁻¹) | | | |
|---------------------|------------------|---------------------------------|-------------------------------|------------------|------------------|-----|----|--------------------------------|-----|-------------------------------|------------------|
| | | N | P ₂ O ₅ | K ₂ O | N | P | K | FYM | N | P ₂ O ₅ | K ₂ O |
| Paddock grazed | | | | | | | | | | | |
| Not mown | 153 | 2 | 213 | 23 | 96 | 73 | 74 | 32 | 221 | 32 | 47 |
| Mown | 84 | 1 | 180 | 26 | 98 | 76 | 83 | 54 | 184 | 35 | 59 |
| All paddock grazed | 237 | 3 | 202 | 24 | 97 | 74 | 77 | 39 | 208 | 33 | 52 |
| Strip grazed | | | | | | | | | | | |
| Not mown | 135 | 2 | 187 | 18 | 19 | 93 | 64 | 53 | 201 | 28 | 29 |
| Mown | 142 | 3 | 206 | 31 | 60 | 97 | 78 | 84 | 212 | 40 | 71 |
| All strip grazed | 277 | 4 | 198 | 25 | 42 | 95 | 71 | 75 | 207 | 36 | 56 |
| Set stocked | | | | | | | | | | | |
| Not mown | 876 | 15 | 139 | 19 | 19 | 86 | 55 | 54 | 162 | 34 | 35 |
| Mown | 730 | 9 | 180 | 31 | 52 | 96 | 75 | 77 | 187 | 41 | 67 |
| All set stocked | 1606 | 23 | 154 | 23 | 31 | 90 | 62 | 63 | 172 | 37 | 49 |
| Other grazings | | | | | | | | | | | |
| Not mown | 2289 | 39 | 85 | 21 | 19 | 75 | 61 | 59 | 28 | 113 | 35 |
| Mown | 1850 | 24 | 154 | 31 | 48 | 96 | 77 | 78 | 61 | 160 | 40 |
| All other grazings | 4139 | 63 | 111 | 25 | 30 | 83 | 67 | 66 | 41 | 134 | 37 |
| All grazings | 6259 | 94 | 129 | 24 | 31 | 86 | 66 | 66 | 41 | 150 | 37 |
| Cut for silage | | | | | | | | | | | |
| Not grazed | 155 | 2 | 302 | 45 | 105 | 100 | 83 | 85 | 48 | 303 | 54 |
| Grazed extensively | 943 | 11 | 202 | 37 | 67 | 100 | 85 | 88 | 68 | 203 | 43 |
| Grazed intensively | 602 | 8 | 235 | 34 | 69 | 99 | 79 | 84 | 65 | 239 | 44 |
| All cut for silage | 1700 | 22 | 223 | 37 | 71 | 99 | 83 | 86 | 65 | 224 | 44 |
| Cut for hay | | | | | | | | | | | |
| Not grazed | 85 | 1 | 85 | 21 | 23 | 90 | 60 | 58 | 41 | 94 | 35 |
| Grazed extensively | 851 | 11 | 97 | 24 | 26 | 92 | 67 | 67 | 52 | 105 | 40 |
| Grazed intensively | 402 | 5 | 113 | 25 | 29 | 94 | 72 | 73 | 51 | 119 | 34 |
| All cut for hay | 1338 | 17 | 101 | 24 | 27 | 93 | 69 | 68 | 51 | 109 | 40 |
| All mowings | 3069 | 40 | 171 | 31 | 52 | 96 | 77 | 78 | 59 | 177 | 41 |
| Not stated/not used | 160 | 3 | 67 | 17 | 16 | 53 | 35 | 35 | 22 | 128 | 44 |
| All grass | 6682 | 100 | 131 | 25 | 32 | 85 | 66 | 66 | 41 | 154 | 38 |

* The average application of any fertilizer component over all fields including those receiving none is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.

† Note that fields which are both grazed and mown appear in both the grazing and mowing sections of the table.

USE OF FERTILIZERS IN ENGLAND AND WALES, 1984

TABLE 8
Percentages of grassland area getting different amounts of N (kg ha⁻¹)

| | Fields | 0 | <25 | 25- | 50- | 75- | 100- | 125- | 150- | 200- | 250- | 300- | 400+ |
|---------------------|--------|----|-----|-----|-----|-----|------|------|------|------|------|------|------|
| Paddock grazed | | | | | | | | | | | | | |
| Not mown | 153 | 4 | 3 | 6 | 7 | 11 | 3 | 8 | 10 | 1 | 11 | 27 | 7 |
| Mown | 84 | 2 | 0 | 9 | 7 | 10 | 15 | 2 | 15 | 7 | 13 | 18 | 1 |
| All paddock grazed | 237 | 3 | 2 | 7 | 7 | 11 | 7 | 6 | 12 | 3 | 12 | 24 | 5 |
| Strip grazed | | | | | | | | | | | | | |
| Not mown | 135 | 7 | 0 | 12 | 6 | 1 | 14 | 10 | 9 | 6 | 12 | 19 | 5 |
| Mown | 142 | 3 | 0 | 4 | 3 | 4 | 4 | 8 | 29 | 14 | 12 | 18 | 2 |
| All strip grazed | 277 | 5 | 0 | 7 | 4 | 3 | 8 | 9 | 20 | 11 | 12 | 18 | 3 |
| Set stocked | | | | | | | | | | | | | |
| Not mown | 876 | 14 | 1 | 10 | 14 | 11 | 7 | 8 | 8 | 7 | 6 | 9 | 4 |
| Mown | 730 | 4 | 1 | 8 | 10 | 10 | 9 | 6 | 13 | 12 | 10 | 14 | 4 |
| All set stocked | 1606 | 10 | 1 | 9 | 13 | 10 | 8 | 7 | 10 | 9 | 8 | 11 | 4 |
| Other grazings | | | | | | | | | | | | | |
| Not mown | 2289 | 25 | 4 | 17 | 13 | 10 | 5 | 6 | 7 | 5 | 2 | 3 | 1 |
| Mown | 1850 | 4 | 1 | 11 | 11 | 12 | 8 | 9 | 16 | 11 | 8 | 9 | 1 |
| All other grazings | 4139 | 17 | 3 | 15 | 12 | 11 | 7 | 7 | 10 | 7 | 5 | 5 | 1 |
| All grazings | | | | | | | | | | | | | |
| Cut for silage | | | | | | | | | | | | | |
| Not grazed | 155 | 0 | 1 | 0 | 1 | 1 | 1 | 1 | 12 | 2 | 12 | 19 | 19 |
| Grazed extensively | 943 | 0 | 1 | 3 | 5 | 6 | 8 | 10 | 20 | 18 | 11 | 15 | 2 |
| Grazed intensively | 602 | 1 | 0 | 1 | 2 | 4 | 5 | 6 | 6 | 18 | 16 | 24 | 5 |
| All cut for silage | 1700 | 1 | 1 | 2 | 4 | 5 | 7 | 9 | 9 | 18 | 17 | 13 | 20 |
| Cut for hay | | | | | | | | | | | | | |
| Not grazed | 85 | 10 | 1 | 25 | 13 | 15 | 19 | 6 | 6 | 1 | 1 | 3 | 0 |
| Grazed extensively | 851 | 8 | 2 | 20 | 18 | 19 | 9 | 7 | 10 | 3 | 4 | 1 | 0 |
| Grazed intensively | 402 | 6 | 2 | 15 | 16 | 14 | 13 | 5 | 15 | 7 | 3 | 2 | 1 |
| All cut for hay | 1338 | 7 | 2 | 19 | 17 | 17 | 10 | 7 | 12 | 4 | 4 | 2 | 0 |
| All mowings | 3069 | 4 | 1 | 9 | 9 | 10 | 9 | 8 | 15 | 11 | 9 | 12 | 3 |
| Not stated/not used | | | | | | | | | | | | | |
| All grass | 160 | 47 | 3 | 6 | 7 | 14 | 3 | 5 | 4 | 7 | 1 | 3 | 1 |
| | 6682 | 15 | 2 | 12 | 12 | 10 | 7 | 7 | 10 | 8 | 6 | 8 | 2 |

ROTHAMSTED REPORT FOR 1984, PART 2

about the level of 1981, but more of this N was applied straight in 1984. The latest results confirm the conjecture that N use on spring barley was exceptional in 1983 because of the spring rainfall pattern.

The average amounts of fertilizer nutrients used per hectare in 1984 on individual tillage crops, and on grassland classified according to utilization, and the proportions of each crop which got different amounts of nutrient are summarized in Tables 3-8 in this paper.

REFERENCES

- CHURCH, B. M. & LEWIS, D. A. (1977) Fertiliser use on farm crops in England and Wales: Information from the Survey of Fertiliser Practice 1942-1976. *Outlook on Agriculture* **9**, 186-193.
CHURCH, B. M. & LEECH, P. K. (1984) Fertilizer use on farm crops in Scotland, 1983: *Rothamsted duplicated report*.