


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Rothamsted Experimental Station Report for 1985

[Full Table of Content](#)



## Use of Fertilizers in England and Wales, 1985

**J. I. Elsmere**

J. I. Elsmere (1986) *Use of Fertilizers in England and Wales, 1985* ; Rothamsted Experimental Station Report For 1985, pp 245 - 251 - DOI: <https://doi.org/10.23637/ERADOC-1-34076>

## Use of fertilizers in England and Wales, 1985

J. I. ELSMERE

This series of annual surveys continued in 1985 as a collaborative project between ADAS soil scientists, representatives of the Fertiliser Manufacturers' Association and Rothamsted. Due to reduced resources, ADAS withdrew from the fieldwork this year, but the size of the survey was almost maintained at previous levels. A sample of 1187 farms was surveyed by Farm Research Limited on behalf of the Fertiliser Manufacturers' Association during June and July. A similar survey was again done in Scotland on a sample of 264 farms, and the results of this will be reported elsewhere.

Compared with 1984, the survey estimates for England and Wales show little change in total use of N, P and K per hectare crops and grass (Table 1). However there continues to be some increase in N use per hectare on tillage crops. Following the trend of recent years, use of N in compound fertilizers decreased, while that of straight N increased. This is particularly noticeable on winter wheat (Table 2) where use of compound N decreased sharply from 16 to 11 kg ha<sup>-1</sup>; this was more than compensated for by an increase in use of straight N from 171 to 181 kg ha<sup>-1</sup>. The proportion of winter wheat which got any N in compounds decreased from 67% in 1984 to 57% in 1985. Apart from the trend towards straight N, this change resulted from use of relatively more PK and relatively less low-N compound fertilizers. Unlike recent years, the percentage of winter-sown cereals remained about the same as for 1984 at about 84%. After estimated increases in 1984, use of P and K per hectare on tillage crops was about the same as in 1983.

**TABLE 1**  
*Fertilizer use on tillage crops and grassland (kg ha<sup>-1</sup>), 1982–85*

	Tillage crops				Grassland				All crops and grass			
	1982	1983	1984	1985	1982	1983	1984	1985	1982	1983	1984	1985
N Straight	99	116	128	134	71	69	71	70	85	91	99	102
Compound	42	37	34	27	52	57	61	61	47	48	48	44
Total	141	154	162	161	123	126	132	131	132	139	147	146
P <sub>2</sub> O <sub>5</sub>	55	54	61	56	24	26	25	24	39	39	42	40
K <sub>2</sub> O	61	60	68	63	28	28	33	32	44	44	50	48

**TABLE 2**  
*Fertilizer use on winter wheat and spring barley (kg ha<sup>-1</sup>) 1982–85*

	Winter wheat				Spring barley			
	1982	1983	1984	1985	1982	1983	1984	1985
N Straight	148	167	171	181	34	51	44	53
Compound	18	16	16	11	60	57	54	49
Total	166	182	187	192	94	107	98	102
P <sub>2</sub> O <sub>5</sub>	51	51	56	54	38	39	39	38
K <sub>2</sub> O	45	46	53	52	41	44	44	44

Average N use on sugar beet, well above general recommendations for some time, has recently tended to fluctuate from year to year, but was well down at 126 kg ha<sup>-1</sup> in 1985 compared with 148 kg ha<sup>-1</sup> the previous year. Estimated use of P and K on sugar beet was also less in 1985.

Recent years have shown small increases in lime use, but in 1984/5, 5·5% of the crops and grass area was treated with lime compared with 7·5% in 1983/4. The smaller use this year was probably because of unfavourable weather conditions in autumn 1984.

The average amounts of fertilizer nutrients used per hectare in 1985 on individual tillage crops, and on grassland classified according to utilization, and the proportions of each crop which got different amounts of nutrients are summarized in Tables 3–8 at the end of this paper.

## ROTHAMSTED REPORT FOR 1985, PART 2

TABLE 3  
*Fertilizer use in England and Wales, 1985*

Fields	Hectares ('000)	Overall* (kg ha <sup>-1</sup> )			% Area receiving			Actual* (kg ha <sup>-1</sup> )
					N	P	K	
		N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O				
Spring wheat	43	13	145	45	52	99	95	26
Winter wheat	3651	1832	192	54	52	100	87	146
Spring barley	1187	490	102	38	44	99	91	48
Winter barley	1926	918	150	55	58	100	94	62
Spring oats	60	17	70	35	39	96	93	63
Winter oats	225	80	119	57	59	100	93	42
Rye	46	12	130	41	53	97	94	37
Maize	22	7	74	45	45	78	67	73
Early potatoes	57	18	194	186	243	100	100	45
Maincrop potatoes	331	94	198	209	278	99	99	194
Sugar beet	455	187	126	56	137	99	93	186
Oilseed rape	591	299	272	58	55	100	94	210
Swedes (stock)	63	14	56	93	73	89	93	278
Turnips (stock)	70	20	73	51	53	98	84	60
Kale and cow cabbage	48	10	106	45	47	95	79	64
Rape for stockfeed	28	8	94	50	43	100	65	53
Beans for stockfeed	121	47	47	2	35	29	13	57
Other stockfeed	83	27	73	52	61	71	87	102
Peas for human consumption	296	121	4	23	31	18	44	18
Runner and French beans	27	7	152	60	86	91	89	53
Brussels sprouts	36	8	209	109	204	100	100	67
Cabbages	58	12	175	71	152	98	86	205
Cauliflower	47	17	192	106	193	92	93	110
Onions	41	11	133	81	157	100	87	166
Small fruit	45	11	48	19	81	76	50	133
Top fruit	59	30	48	15	26	50	40	93
All tillage	9954	4397	161	56	63	95	88	162
1 year leys	66	24	180	34	46	99	73	38
2-7 year leys	2900	1510	185	33	50	96	77	100
Permanent grass	4131	2815	102	19	22	79	59	32
All crops and grass	17051	8746	146	40	48	90	77	38

\*The average application of any fertilizer component over all fields including those receiving none is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.

## USE OF FERTILIZERS IN ENGLAND AND WALES, 1985

**TABLE 4**  
*Percentages of crop area getting different amounts of N (kg ha<sup>-1</sup>)*

Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	43	1	0	3	2	8	13	22	47	4	0	0
Winter wheat	3651	0	0	0	0	2	4	9	37	40	6	1
Spring barley	1187	1	1	9	21	19	22	15	11	2	0	0
Winter barley	1926	0	0	1	2	6	13	28	41	8	1	0
Spring oats	60	4	3	24	34	28	3	2	0	0	0	0
Winter oats	225	0	0	2	7	14	36	20	18	1	0	0
Rye	46	3	4	1	5	8	26	23	15	10	4	0
Maize	22	22	8	12	4	21	15	11	8	0	0	0
Early potatoes	57	0	1	0	0	0	13	4	35	28	12	5
Maincrop potatoes	331	1	0	1	3	4	12	25	37	28	17	2
Sugar beet	455	1	1	1	3	4	12	25	32	20	2	0
Oilseed rape	591	0	0	0	0	1	0	1	2	22	44	29
Swedes (stock)	63	11	5	35	22	18	2	5	1	0	0	0
Turnips (stock)	70	2	7	34	13	21	9	9	5	0	0	0
Kale and cow cabbage	48	5	3	15	12	9	19	24	10	0	1	3
Rape for stockfeed	28	0	0	17	18	38	10	8	6	0	3	0
Beans for stockfeed	121	87	10	2	0	1	0	0	0	0	0	0
Other stockfeed	83	29	5	13	12	12	6	7	14	4	0	0
Peas for human consumption	296	82	15	1	2	0	0	0	0	0	0	0
Runner and French beans	27	9	0	0	0	0	18	11	58	2	3	0
Brussels sprouts	36	0	0	0	0	0	4	16	8	33	6	5
Cabbages	58	2	0	11	4	3	5	9	22	28	13	3
Cauliflower	47	8	0	3	7	0	22	3	9	7	26	15
Onions	41	0	0	8	9	27	19	0	13	9	15	0
Small fruit	45	24	24	10	6	4	29	0	2	0	0	0
Top fruit	59	50	1	19	4	4	0	8	13	0	0	0
All tillage	9954	5	1	2	4	6	9	14	28	21	6	2
1 year leys	66	1	0	6	7	6	13	8	25	13	8	4
2-7 year leys	2900	4	1	8	9	8	6	10	14	10	5	5
Permanent grass	4131	21	3	13	14	9	7	7	10	4	4	1
All crops and grass	17051	10	2	7	8	7	8	11	20	14	7	5

ROTHAMSTED REPORT FOR 1985, PART 2

	Percentages of crop area getting different amounts of $P_2O_5$ ( $kg\ ha^{-1}$ )											
Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	43	5	11	47	34	1	2	0	0	0	0	0
Winter wheat	3651	13	3	17	52	12	2	0	1	0	0	0
Spring barley	1187	9	21	44	24	2	0	0	0	0	0	0
Winter barley	1926	6	5	23	54	10	1	0	1	0	0	0
Spring oats	60	7	33	43	14	4	0	0	0	0	0	0
Winter oats	225	7	4	13	66	8	1	0	0	0	0	0
Rye	46	20	14	16	42	7	0	0	0	0	0	0
Maize	22	33	9	31	13	0	6	0	8	0	0	0
Early potatoes	57	0	1	1	7	0	1	17	30	37	1	6
Maincrop potatoes	331	0	0	1	2	3	3	4	35	24	9	9
Sugar beet	455	7	7	35	29	13	5	0	2	0	0	0
Oilseed rape	591	6	2	16	62	11	2	0	0	0	0	0
Swedes (stock)	63	7	6	8	22	23	17	7	4	1	0	4
Turnips (stock)	70	16	30	17	14	9	10	1	0	2	1	0
Kale and cow cabbage	48	21	18	23	22	7	4	0	3	0	1	0
Rape for stockfeed	28	35	7	30	4	7	8	0	7	2	0	0
Beans for stockfeed	121	43	2	14	31	7	1	0	2	0	0	0
Other stockfeed	83	13	8	17	47	8	4	0	2	0	0	0
Peas for human consumption	296	56	1	16	24	1	1	1	0	0	0	0
Runner and French beans	27	11	6	15	50	11	0	0	8	0	0	0
Brussels sprouts	36	0	0	13	24	20	8	6	25	5	0	0
Cabbages	58	14	1	33	23	5	8	1	14	7	0	0
Cauliflower	47	7	0	4	32	17	5	11	18	6	0	0
Onions	41	13	1	10	45	8	3	8	0	0	0	0
Small fruit	45	50	27	11	4	8	0	0	0	2	0	0
Top fruit	59	60	20	16	3	1	0	0	0	1	0	0
All tillage	9954	12	6	22	45	9	2	1	2	0	0	0
1 year leys	66	27	22	30	7	10	4	1	2	0	0	0
2-7 year leys	2900	23	28	30	11	5	1	1	1	0	0	0
Permanent grass	4131	41	33	17	22	26	6	1	0	0	0	0
All crops and grass	17051	23	19									

## USE OF FERTILIZERS IN ENGLAND AND WALES, 1985

TABLE 6  
Percentages of crop area getting different amounts of  $K_2O$  ( $kg\ ha^{-1}$ )

	Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Spring wheat	43	5	7	28	56	2	0	2	0	0	0	0	0
Winter wheat	3651	18	3	17	47	12	3	1	0	0	0	0	0
Spring barley	1187	10	13	36	36	3	1	0	0	0	0	0	0
Winter barley	1926	8	4	19	53	12	3	1	1	0	0	0	0
Spring oats	60	7	25	38	26	4	1	0	0	0	0	0	0
Winter oats	225	6	3	17	62	7	4	0	0	0	0	0	0
Rye	46	13	10	21	37	14	0	5	0	0	0	0	0
Maize	22	33	9	31	13	0	6	0	8	0	0	0	0
Early potatoes	57	0	1	0	0	0	2	6	16	46	3	25	0
Maincrop potatoes	331	0	1	0	1	1	1	3	7	22	29	34	2
Sugar beet	455	1	1	2	11	16	16	9	27	15	1	0	0
Oilseed rape	591	13	1	11	59	12	2	0	0	0	0	0	0
Swedes (stock)	63	19	6	3	20	33	5	10	2	0	1	0	0
Turnips (stock)	70	17	19	16	27	8	9	0	3	1	0	0	0
Kale and cow cabbage	48	21	18	10	34	6	10	0	0	0	0	1	0
Rape for stockfeed	28	19	7	37	25	3	5	0	0	0	0	0	0
Beans for stockfeed	121	49	1	16	28	4	1	0	0	0	0	0	0
Other stockfeed	83	17	3	16	34	15	4	5	2	2	0	0	0
Peas for human consumption	296	52	1	9	35	1	1	1	1	0	0	0	0
Runner and French beans	27	11	0	6	37	19	12	7	0	5	4	0	0
Brussels sprouts	36	0	0	0	0	2	11	8	17	55	7	0	0
Cabbages	58	8	0	0	0	2	7	0	3	16	14	0	0
Cauliflower	47	7	0	0	5	0	3	0	10	17	40	17	1
Onions	41	3	0	3	10	3	8	7	17	25	24	8	0
Small fruit	45	19	3	8	25	9	7	0	28	0	0	0	0
Top fruit	59	50	13	8	24	3	1	0	0	0	0	0	0
All tillage	9954	14	4	18	43	10	3	1	2	2	1	1	0
1 year leys	66	27	20	13	19	3	7	6	4	0	0	0	0
2-7 year leys	2900	20	20	21	16	7	7	6	3	0	0	0	0
Permanent grass	4131	42	27	19	7	2	2	1	2	1	0	0	0
All crops and grass	17051	24	14	19	27	3	2	2	1	0	0	0	0

ROTHAMSTED REPORT FOR 1985, PART 2

TABLE 7  
*Fertilizer use on grassland classified by utilization<sup>†</sup>*

Fields	% Grassland area	Overall* (kg ha <sup>-1</sup> )			% Area receiving			Actual* (kg ha <sup>-1</sup> )
		N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N	P	K	
Paddock grazed								
Not mown	121	2	181	15	22	83	46	30
Mown	95	1	209	26	44	95	68	74
All paddock grazed	216	4	192	20	31	88	55	42
Strip grazed								
Not mown	117	2	230	23	27	98	69	50
Mown	178	2	204	24	46	96	79	83
All strip grazed	295	4	216	23	37	97	74	77
Set stocked								
Not mown	905	13	161	24	23	89	67	66
Mown	851	11	188	33	57	95	79	82
All set stocked	1756	24	173	28	39	92	73	44
Other grazings								
Not mown	2512	39	80	19	17	74	57	55
Mown	1980	25	143	27	45	92	72	76
All other grazings	4492	64	105	22	28	81	63	38
All grazings	6759	95	130	24	31	85	66	41
Cut for seed								
Not grazed	22	0	149	59	69	94	87	76
All cut for seed	36	0	160	46	56	97	78	74
Cut for silage								
Not grazed	132	2	240	39	84	99	78	86
Grazed extensively	1090	13	192	33	62	98	79	84
Grazed intensively	757	10	230	36	66	100	85	89
All cut for silage	1979	25	211	34	65	99	82	86
Cut for hay								
Not grazed	119	1	112	12	18	95	33	45
Grazed extensively	834	11	84	21	25	84	65	66
Grazed intensively	409	5	103	18	25	85	60	63
All cut for hay	1362	17	91	19	25	85	62	64
All mowings								
Not stated/not used	108	2	68	23	19	62	48	47
All grass	7140	100	131	24	32	85	65	66

\*The average application of any fertilizer component over all fields including those receiving none is termed 'overall'. The average excluding fields with none of the component is termed 'actual'.

† Note that fields which are both grazed and mown will appear in both the grazing and mowing sections of the table.

USE OF FERTILIZERS IN ENGLAND AND WALES, 1985

TABLE 8  
*Percentages of grassland area getting different amounts of N (kg ha<sup>-1</sup>)*

	Fields	0	<25	25-	50-	75-	100-	125-	150-	200-	250-	300-	400+
Paddock grazed													
Not mown	121	17	1	10	2	4	9	6	5	9	16	17	5
Mown	95	5	2	2	4	4	5	12	25	11	5	14	10
All paddock grazed	216	12	2	7	3	4	8	9	13	10	11	16	7
Strip grazed													
Not mown	117	2	0	3	12	4	6	8	13	5	10	26	10
Mown	178	4	2	2	4	5	3	13	14	18	16	13	5
All strip grazed	295	3	1	3	8	5	4	10	14	12	13	19	7
Set stocked													
Not mown	905	11	2	6	13	7	8	6	15	9	9	9	5
Mown	851	5	1	5	8	10	6	7	17	13	11	15	4
All set stocked	1756	8	1	6	11	8	7	6	16	10	10	12	4
Other grazings													
Not mown	2512	26	3	18	15	10	6	8	6	3	4	2	1
Mown	1980	8	1	11	13	10	7	9	15	9	8	7	2
All other grazings	4492	19	2	15	14	10	6	8	10	5	5	4	1
All grazings	6759	15	2	12	13	9	6	8	11	7	7	7	3
Cut for seed													
Not grazed	22	6	0	2	0	3	15	14	47	12	0	0	0
All cut for seed	36	3	0	1	4	12	14	8	34	16	6	2	0
Cut for silage													
Not grazed	132	1	0	1	4	4	12	12	10	12	7	33	5
Grazed extensively	1090	2	0	5	7	9	7	10	18	12	14	12	5
Grazed intensively	757	0	0	1	4	7	4	6	20	17	15	19	6
All cut for silage	1979	1	0	3	5	8	6	9	18	14	14	16	5
Cut for hay													
Not grazed	119	5	0	9	24	19	13	7	10	8	6	1	0
Grazed extensively	834	16	2	19	22	12	7	8	9	4	2	1	0
Grazed intensively	409	15	3	12	14	11	8	12	14	6	1	3	0
All cut for hay	1362	15	2	17	20	12	7	9	11	5	2	1	0
All mowings	3377	7	1	8	11	9	7	9	15	10	9	10	3
Not stated/not used													
All grass	108	38	3	21	2	8	2	17	3	1	1	4	1
All grass	7140	15	2	12	9	6	8	11	7	7	7	7	3