

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Memoranda of the Plan and Results of the Rothamsted Field Experiments, June 1862



[Full Table of Content](#)

---

## Experiments on Oats; Hoos Field

### Rothamsted Research

Rothamsted Research (1862) *Experiments on Oats; Hoos Field* ; Memoranda Of The Plan And Results Of The Rothamsted Field Experiments, June 1862, pp 3 - 3 - DOI:

<https://doi.org/10.23637/ERADOC-1-230>

EXPERIMENTS ON THE GROWTH OF BARLEY YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE AND WITH DIFFERENT KINDS OF MANURE.  
HOOS FIELD.

Previous cropping—1847, Swedish turnips, with dung and superphosphate of lime, the roots carted off; 1848, Barley; 1849, Clover; 1850, Wheat; 1851, Barley manured with ammonia-salts.  
First experimental Barley-crop in 1852. Barley every year since; and, with one or two exceptions, the same manures on the same plots each year.  
(Area under experiment, about 4½ acres).

PLOTS.	Manures, per acre, for the growing (11th) Barley crop—1862. 1 Acre ... = (about) 0.40 Hectare. 1 Bushel... = (about) 0.36 Hectolitre. 1 lb. (pound, avoird.) = (about) 0.45 Kilogramme. 1 Bushel per acre = (nearly) 0.9 Hectolitre per Hectare. 1 lb. per acre = 1.12 Kilogramme per Hectare.	Average Produce per Acre, per Annum, during Ten Years, 1851-61.	
		Dressed Corn.	Total Corn.
1 O.	Unmanured, continuously	Bushels.	lbs.
2 O.	Superphosphate of Lime (1)	22½	1281
3 O.	Mixed Alkalies (2)	28	1562
4 O.	ditto	24½	1396
	; and "Superphosphate of Lime"	30½	1712
6 1	Unmanured, continuously	25	1414
7	Ashes (burnt soil, turf, and weeds)	24	1352
	Farm-yard dung (14 tons every year)	45	2541
1 A.	200 lbs. Ammonia-Salts (3)	33½	1908
2 A.	ditto	45½	2563
3 A.	; and "Superphosphate of Lime"	35	1989
4 A.	; "Superphosphate of Lime"; and "Mixed Alkalies"	46½	2593
1 A.A.	200 lbs. (4) ditto	39½	2244
2 A.A.	ditto	49	2744
3 A.A.	; "Superphosphate of Lime"	38½	2190
4 A.A.	; "Superphosphate of Lime"; and "Mixed Alkalies"	50	2772
1 C.	1000 lbs. (5) Rapecake	47	2619
2 C.	1000 lbs. (6) ditto	47½	2677
3 C.	1000 lbs. (6) ditto	44	2480
4 C.	1000 lbs. (6) ditto	47½	2652
1 N. (6)	275 lbs. Nitrate of Soda	37½ (6)	2125 (6)
2 N. (6)	ditto	42½ (6)	2376 (6)
5 O.	200 lbs. (9) Sulphate of Potass	24½ (11)	1373 (11)
5 A.	ditto	44½ (11)	2470 (11)
M.	100 lbs. each, Sulph. Soda and Sulph. Magnesia; and	22½ (12)	1262 (12)

(1) 200 lbs. bone-ash, 150 lbs. sulphuric acid (sp. gr. 1.7).  
 (2) 200 lbs. sulphate of potass, 100 lbs. sulphate of soda, and 100 lbs. sulphate of magnesia; for the first six years, 300 lbs., 200 lbs., and 100 lbs. respectively.  
 (3) Equal parts sulphate and muriate of commerce. (4) 400 lbs. per annum for the first six years, and 200 lbs. only each year since.  
 (5) 2000 lbs. per annum for the first six years, and 1000 lbs. only, each year since.  
 (6) 300 lbs. sulphate of potass, 200 lbs. bone-ash, and 150 lbs. sulphuric acid (sp. gr. 1.7), without nitrate of soda, the first year (1852); nitrate alone each year since.  
 (7) 350 lbs. nitrate of soda for 1853-4-5-6, and 7; and 275 lbs. only each year since.  
 (8) 300 lbs. per annum for the first six years, and 200 lbs. each year since.  
 (9) Average of 7 years only.  
 (10) Ammonia-salts also the first year, but not since.  
 (11) Average of 9 years only.  
 (12) Average of 9 years only.