

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Plan and Results of the Rothamsted Field Experiments, June 1862



[Full Table of Content](#)

Experiments on Rotation; Agdell Field

Rothamsted Research

Rothamsted Research (1862) *Experiments on Rotation; Agdell Field* ; Memoranda Of The Plan And Results Of The Rothamsted Field Experiments, June 1862, pp 6 - 6 - DOI:

<https://doi.org/10.23637/ERADOC-1-230>

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present crop is the 19th experimental one, or the 11th crop of the 1st Course. One-third of the land has been continuously unmanured; and one-third manured with superphosphate of lime alone once every four years, that is, for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described below. In the Second, Third, and Fourth Courses, instead of clover, half of each plot was sown with beans, and the other half left fallow. From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the unclean leaves were spread and ploughed in. In the case of all the other crops, the total produce was removed from the land. The abstract of results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the later Courses, beans (not fallow) replaced the clover.

(Area under experiment about 2½ acres.)

| Years. | Description of Crop. | PLOT I. Unmanured continuously. | | | PLOT 2. Superphosphate of Lime, (1) alone, for the Turnip crops only. | | | PLOT 3. Complex Manure, (2) for the Turnip crops only. | | |
|------------------------|---------------------------|------------------------------------|------------------|----------------|--------------------------------------------------------------------------|------------------|----------------|-----------------------------------------------------------|------------------|----------------|
| | | Corn (or Roots). | Straw (or Leaf). | Total Produce. | Corn (or Roots). | Straw (or Leaf). | Total Produce. | Corn (or Roots). | Straw (or Leaf). | Total Produce. |
| 1st Course, 1848-51. | | | | | | | | | | |
| 1848 | Swedish Turnips | 175½ cwts. | 19½ cwts. | 195 cwts. | 292 cwts. | 35 cwts. | 327 cwts. | 394½ cwts. | 46½ cwts. | 441 cwts. |
| 1849 | Barley | 1706 lbs. | 2088 lbs. | 3794 lbs. | 1705 lbs. | 1870 lbs. | 3575 lbs. | 2673 lbs. | 2988 lbs. | 5656 lbs. |
| 1850 | Clover (weighed green) .. | | | 194½ cwts. | | | 199½ cwts. | | | 219½ cwts. |
| 1851 | Wheat | 1958 lbs. | 3431 lbs. | 5389 lbs. | 1882 lbs. | 3371 lbs. | 5253 lbs. | 1948 lbs. | 3552 lbs. | 5500 lbs. |
| 2nd Course, 1852-55. | | | | | | | | | | |
| 1852 | Swedish Turnips | 26 cwts. | 4½ cwts. | 30½ cwts. | 223½ cwts. | 204 cwts. | 243½ cwts. | 396½ cwts. | 36½ cwts. | 483 cwts. |
| 1853 | Barley | 2035 lbs. | 2430 lbs. | 4465 lbs. | 1687 lbs. | 1873 lbs. | 3560 lbs. | 2269 lbs. | 2604 lbs. | 4873 lbs. |
| 1854 | Beans | 390 lbs. | 1055 lbs. | 1445 lbs. | 431 lbs. | 1103 lbs. | 1534 lbs. | 710 lbs. | 1355 lbs. | 2065 lbs. |
| 1855 | Wheat | 2240 lbs. | 3619 lbs. | 5859 lbs. | 2264 lbs. | 3525 lbs. | 5789 lbs. | 2429 lbs. | 3942 lbs. | 6371 lbs. |
| 3rd Course, 1856-59. | | | | | | | | | | |
| 1856 | Swedish Turnips | 32 cwts. | 2½ cwts. | 34½ cwts. | 136 cwts. | 7½ cwts. | 143½ cwts. | 333½ cwts. | 12½ cwts. | 346½ cwts. |
| 1857 | Barley | 2737 lbs. | 2600 lbs. | 5337 lbs. | 1601 lbs. | 1475 lbs. | 3076 lbs. | 2733 lbs. | 2435 lbs. | 5168 lbs. |
| 1858 | Beans | 415 lbs. | 1100 lbs. | 1515 lbs. | 450 lbs. | 1155 lbs. | 1605 lbs. | 837 lbs. | 1520 lbs. | 2357 lbs. |
| 1859 | Wheat | 2232 lbs. | 4030 lbs. | 6262 lbs. | 2190 lbs. | 3930 lbs. | 6120 lbs. | 2544 lbs. | 4610 lbs. | 7154 lbs. |
| 4th Course, 1860-(63). | | | | | | | | | | |
| 1860 | Swedish Turnips | 1 cwt. | 19½ lbs.) | 1 cwt. | 29½ cwts. | 1½ cwt. | 30½ cwts. | 87½ cwts. | 3½ cwts. | 90½ cwts. |
| 1861 | Barley | 2196 lbs. | 2522 lbs. | 4718 lbs. | 1775 lbs. | 2000 lbs. | 3775 lbs. | 3451 lbs. | 3940 lbs. | 7391 lbs. |
| 1862 | Beans | 1821 lbs. | 1840 lbs. | 3661 lbs. | 1890 lbs. | 2150 lbs. | 4040 lbs. | 2710 lbs. | 3280 lbs. | 5990 lbs. |
| 1863 | Wheat | 2883 lbs. | 3467 lbs. | 6350 lbs. | 2229 lbs. | 3390 lbs. | 5619 lbs. | 2929 lbs. | 4497 lbs. | 7426 lbs. |

(1) First Course—100 lbs. bone-ash, and 100 lbs. sulphuric acid (sp. gr. 1.7); Second Course—160 lbs. bone-ash, 120 lbs. sulphuric acid; Third and Fourth Courses—200 lbs. bone-ash, 150 lbs. sulphuric acid *per acre*.
 (2) First Course—100 lbs. pearlash, 100 lbs. sulphuric acid, 100 lbs. sulphate of ammonia, and 1000 lbs. rape-cake; Second Course—300 lbs. sulphate of potash, 100 lbs. sulphate of soda, 100 lbs. sulphate of ammonia, 100 lbs. sulphate of ammonia, 100 lbs. muriate of ammonia, and 2000 lbs. rape-cake; Third and Fourth Courses—300 lbs. sulphate of potash, 200 lbs. sulphate of soda, 100 lbs. sulphate of ammonia, 100 lbs. sulphuric acid, 100 lbs. sulphate of ammonia, and 2000 lbs. rape-cake *per acre*.