


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Plan and Results of the Rothamsted Field Experiments, May 1866



[Full Table of Content](#)

Experiments on Barley; Hoos Field

Rothamsted Research

Rothamsted Research (1867) *Experiments on Barley; Hoos Field* ; Memoranda Of The Plan And Results Of The Rothamsted Field Experiments, May 1866, pp 3 - 3 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-232>

EXPERIMENTS ON THE GROWTH OF **BARLEY** YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

HOOS FIELD.

Previous Cropping—1847, Swedish Turnips, with Dung and Superphosphate of Lime, the Roots carted off; 1848, Barley; 1849, Clover; 1850, Wheat; 1851, Barley manured with Ammonia-salts.

First Experimental Barley Crop in 1852. Barley every year since; and, with one or two exceptions, the same Manures on the same Plots each year.

(Area under experiment, about 4½ acres.)

PLOTS.	Manures, per acre; fifteen Year Season—1866.	PRODUCE PER ACRE.			
		Average per Annum, over 14 Years, 1852-1865.		Fourth Year Season, 1865.	
		Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.
		Quantity, Bushels.	Weight per Bushel.	Quantity, Bushels.	Weight per Bushel.
1 O.	Unmanured continuously	21½	52	18	54
2 O.	Superphosphate of Lime (1)	27½	53½	18	54
3 O.	Mixed Alkalies (2)	24½	52½	22	54½
4 O.	Do. do.	30½	53	24½	54
6(1)	Unmanured continuously	24½	52½	21	53½
7	Ashes (burnt soil, turf, and weeds)	23½	52½	19½	53½
	Farm-yard dung (14 tons every year)	48½	52½	52½	54½
1 A.	200 lbs. Ammonia-salts (3)	34½	51½	29½	53½
2 A.	do. do.	48½	52½	48½	52½
3 A.	do. do.	36½	52½	33½	54½
4 A.	do. do.	47½	52½	46½	53½
1 A.A.	200 lbs. (4) ditto	30½	51½	33½	53½
2 A.A.	do. do.	50½	52½	47½	52½
3 A.A.	do. do.	39½	51½	34½	52½
4 A.A.	do. do.	51	52½	49	53½
1 A.A.S.	200 lbs. (4) ditto	39½	55½	35	54½
2 A.A.S.	do. do.	51	54½	47½	52½
3 A.A.S.	do. do.	45½	56	41	54½
4 A.A.S.	do. do.	54½	55	50½	53
1 C.	1000 lbs. (6) Rape-cake	46½	52½	45	53½
2 C.	do. do.	48½	52½	46½	53½
3 C.	do. do.	44½	52½	48½	53½
4 C.	do. do.	48½	52½	48½	53½
1 N. (7)	275 lbs. Nitrate of Soda	38½ (11)	52½ (11)	37	54
2 N. (7)	do. do.	43 (11)	52 (11)	39½	53½
5 O.	200 lbs. (8) Sulphate of Potass	24½ (12)	52½ (12)	23	54½
5 A.	do. do.	46½ (12)	52½ (12)	48½	54½
M.	100 lbs. each, Sulph. Soda and Sulph. Magnesia; and	25½ (13)	52½ (13)	19½	54½

(1) 200 lbs. Bone-ash, 150 lbs. Sulphuric acid (sp. gr. 1.7). (2) 200 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, and 100 lbs. Magnesia (for the first six years, 300 lbs., 200 lbs., and 100 lbs., respectively).
 (3) Equal parts Sulphate and Muriate of Ammonia of Commerce. (4) 400 lbs. per annum for the first six years, and 200 lbs. only each year since.
 (5) The application of Silicates did not commence until 1864, so that the average produce given applies to two years only (1864 and 1865). These Silicated plots ("AAS") comprise, respectively, one half of the original "AA" plots, and, as will be seen, they continue to be, in other respects, manured in the same way as the remaining halves.
 (6) 2000 lbs. per annum for the first six years, and 1000 lbs. only, each year since. (7) 300 lbs. Sulphate of Potass, 200 lbs. Bone-ash, and 150 lbs. Sulphuric acid (sp. gr. 1.7), without Nitrate of Soda, the first year (1852); Nitrate alone each year since. (8) 550 lbs. Nitrate of Soda for 1853-4-5-6, and 7; and 275 lbs. only each year since.
 (9) Ammonia-salts also the first year, but not since. (10) Average of 13 years only. (11) Average of 11 years only. (12) Average of 13 years only. (13) Average of 11 years only.