


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Plan and Results of the Field Experiments, May 1869



[Full Table of Content](#)

Experiments on the Growth of Root-crops

Rothamsted Research

Rothamsted Research (1870) *Experiments on the Growth of Root-crops* ; Memoranda Of The Plan And Results Of The Field Experiments, May 1869, pp 6 - 6 - DOI:

<https://doi.org/10.23637/ERADOC-1-233>

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS.

I.—BEANS, PEAS, AND TARES.

EXPERIMENTS on the growth of Leguminous corn-crops, with different descriptions of manure, were commenced in 1847, about nine acres being devoted to the purpose.

Experiments with BEANS were continued for thirteen consecutive seasons, to 1859 inclusive; but, during the later years, the crop fell off very much, and the land became very foul.

In 1860 the land was fallowed.

In 1861 a crop of wheat, without manure, was taken.

In 1862 beans were again sown, but with some variation in the manuring.

In 1863 the land was fallowed.

In 1864, and since, beans have been grown, with much the same manures on the same plots, each year, as in 1862.

The general result of the experiments with BEANS has been, that mineral constituents added as manure (more particularly potash, and, to some extent, phosphoric acid also), increased the crop very much during the early years; and, to a certain extent, afterwards, whenever the season was favourable for the crop. Ammonia-salts, on the other hand, produced very little effect; notwithstanding that a Leguminous crop contains two, three, or more times as much nitrogen as a Gramineous one grown under parallel circumstances. Nitrate of soda, however, has produced very striking effects. But Leguminous crops grown too frequently on the same land seem to be peculiarly subject to disease, which no combination of manuring that we have hitherto tried seems to obviate.

Experiments with PEAS were soon abandoned, owing to the difficulty of keeping the land free from weeds, and an alternation of BEANS and WHEAT was substituted; the beans being manured much as in the experiments with the same crop above described.

In alternating WHEAT with BEANS, the remarkable result has been obtained, that nearly as much wheat, and nearly as much nitrogen, were yielded in eight crops of wheat in alternation with the highly nitrogenous beans, as in sixteen crops of wheat grown consecutively without manure in another field, and also nearly as much as were obtained in a third field in eight crops alternated with bare fallow.

Experiments with TARES were also soon abandoned, for the same reason; beans being at first substituted, with some variation in the description of the manures employed; but of late this experiment has likewise been abandoned.

II.—RED CLOVER (*Trifolium pratense*).

Experiments on the growth of Clover, with different descriptions of manure, were commenced in 1849, and, with the occasional interposition of a corn-crop, or fallow, have been continued up to the present time. As with beans, the result was, that mineral constituents applied as manures (particularly potash, and, more or less, phosphoric acid also), considerably increased the early crops; whereas ammonia-salts had little or no effect. But since the first few years all attempts to grow Clover year after year on this land have failed to give anything like a fair crop, or a plant that would stand the usual time on the ground, notwithstanding that fresh seed has been sown again and again.

In one year a portion of the land was trenched 2 feet deep; one-third of the manure being applied at a depth of 16 inches, one-third at a depth of 8 inches, and the remainder on the surface.

The general result of the experiments is, that neither ammonia-salts, nor nitrate of soda, nor organic matter rich in carbon as well as other constituents, nor mineral manures, nor a complex mixture, has availed to restore the clover-yielding capabilities of the land.

It is, however, worthy of remark that, in 1854, Red Clover was sown in a kitchen-garden only a few hundred yards distant from the experimental field, on soil which has been under ordinary garden cultivation for, probably, two or three centuries, and it has every year since shown very luxuriant growth; and, after re-sowing three times during the period (in 1860, 1865, and 1868), there is, at the present time, little or no indication of failure.

Lastly, in the winter of 1867-8, small portions of the experimental land were dug, some to the depth of 9 inches, some to the depth of 18, some to the depth of 27, and some to the depth of 36 inches, and sown to the respective depths with different manurial mixtures. From other similarly sized plots the soil was removed to the depths of 9, 18, and 27 inches respectively, and replaced by soil from the same kitchen-garden border, on a portion of which Clover has been successfully grown since 1854, as above referred to. Clover was sown in April, 1868, over the whole of these, and some other portions not so treated; but the plant has, for the most part, died off during the winter, and Clover has been again sown (April, 1869).

EXPERIMENTS ON THE GROWTH OF ROOT-CROPS.

EXPERIMENTS with TURNIPS were commenced in 1843. Eight acres, divided into numerous plots, were set apart for the purpose; and the crop was grown for ten consecutive years on the same land ("Norfolk Whites" 1843-1848, and "Swedes" 1849-1852); on some plots without manure, and on others with different descriptions of manure. Barley was then grown for three consecutive seasons (1853-1855) without manure, in order to test the comparative corn-growing condition of the different plots, and also to equalize their condition, as far as possible, by the exhaustion of some of the most active and immediately available constituents supplied by the previous manuring. A new series of experiments with Swedes was then arranged, having regard to the character of the manures previously applied on the different plots, and to the results previously obtained. This second series was commenced in 1856, and is still in progress.

It is impossible adequately to state the bearing of the results in a few words, but the following are some of the most characteristic indications:—

1. Without manure of any kind, the produce of roots was reduced in a few years to a few cwts. per acre; but the diminutive plants (both root and leaf) contained a very unusually high percentage of nitrogen.
2. Of "mineral" constituents, phosphoric acid (in the form of superphosphate of lime) was by far the most effective manure; but, when this manure is used alone, the immediately available nitrogen of the soil is rapidly exhausted.
3. Really large crops of turnips can only be obtained when the soil supplies a liberal amount of both carbonaceous and nitrogenous matter (as well as mineral constituents); and when they are already available within the soil, or are supplied in the form of farmyard manure, rape-cake, Peruvian guano, ammonia-salts, &c., the rapidity of growth and the amount of the crop are greatly increased by the use of superphosphate of lime applied near to the seed.