Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



## Memoranda of the Plan and Results of the Field Experiments, May 1869



Full Table of Content

## **Experiments on Rotation; Agdell Field**

## **Rothamsted Research**

Rothamsted Research (1870) *Experiments on Rotation; Agdell Field;* Memoranda Of The Plan And Results Of The Field Experiments, May 1869, pp 7 - 7 **- DOI:** 

https://doi.org/10.23637/ERADOC-1-233

(7)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION-TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

## AGDELL FIELD.

These Experiments were commenced in 1848; so that the present crop (1869) is the 22nd experimental one, or the second crop of the Sixth Course One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in

the foot-note, No. 2.

In the Second, Third, Fourth, and Fifth Courses, instead of clover, half of each plot was sown with beans, and the other half left fallow.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uncaten leaves were spread and ploughed in. In the case of all the other crops, the total produce was removed from the land.

The abstract of results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the later courses, beans (not fallow) replaced the clover.

(Area under experiment, about 21 acres.)

1 lb. (pound avoir.) per acre = (about) 1·12 Kilogramme per Hectare, or 0·57 Zollverein Pfund. per Prussian Morgen. 1 cwt. (hundredweight) per acre = (about) 125·5 Kilogrammes per Hectare, or 0·64 Centner per Pr. Morgen.										
	Phoduce per Acre.									
Years.	P.Lor 1, Description of Crop. Unmanured continuously.			usly.	PLOT 2. Superphosphate of Lime (1), alone, for the Turnip Crops only.			PLOT 3.  Complex Manure (2), for the Turnip Crops only.		
		Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4).	Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4),	Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4).
1st Course, 1848-51.										
1848 1849 1850 1851	Norfolk White Turnips Barley Clover (calcd, as hay) Wheat.	65½ cwts. 44½ bush. 28½ bush.	45% cwts. 2983 lbs. 3431 lbs.	111½ cwts. 5656 lbs. 54 cwts. 5389 lbs.	2254 cwts. 29% bush. 28 bush.	1064 cwts. 2111 lbs. 3371 lbs.	332 cwts. 3841 lbs. 574 cwts. 5253 lbs.	218 cwts, 28 <sup>7</sup> / <sub>8</sub> bush. 28 <sup>7</sup> / <sub>8</sub> bush.	1513 cwts. 2088 1bs. 3552 1bs.	369% cwts. 3794 lbs. 63 cwts. 5500 lbs.
2nd Course, 1852-55.										
1852 1853 1854 1855	Swedish Turnips. Barley Beans. Wheat	26 cwts. 34% bush, 5% bush. 35% bush.	4½ cwts. 2430 lbs. 1055 lbs. 3619 lbs.	304 cwts, 4465 lbs, 1445 lbs, 5859 lbs.	2234 cwts. 28§ bush. 57 bush. 354 bush.	20½ cwts, 1873 lbs, 1103 lbs, 3525 lbs.	243½ cwts. 3560 lbs. 1534 lbs. 5789 lbs.	390½ cwts. 39½ bush. 9½ bush. 37% bush.	36½ cwts, 2604 lbs. 1355 lbs, 3942 lbs.	433 cwts, 4873 lbs. 2065 lbs. 6371 lbs.
3rd Course, 1856-59.										
1856 1857 1858 1859	Swedish Turnips Barley Beans Wheat	32 cwts. 48½ bush. 6½ bush. 35½ bush.	2½ cwts. 2600 lbs. 1100 lbs. 4030 lbs.	34½ cwts 5337 lbs. 1515 lbs. 6262 lbs.	136 cwts. 28½ bush. 6½ bush. 34½ bush.	7½ cwts, 1475 lbs, 1155 lbs, 3930 lbs.	142½ cwts. 3076 lbs. 160 <b>5</b> lbs. 6120 lbs,	333% ewts. 48 bush. 12% bush. 39% bush.	12½ cwts, 2435 lbs. 1520 lbs. 4610 lbs.	3464 cwts. 5168 lbs. 2357 lbs. 7154 lbs.
4ти Course, 1860-63.										
1860 1861 1862 1863	Swedish Turnips. Barloy. Beans. Wheat	1 cwt. 39§ bush. 29 bush. 44¼ bush.	(64 lbs.) 2522 lbs. 1840 lbs. 3467 lbs.	1 cwt. 4718 lbs. 3661 lbs. 6350 lbs.	294 cwts. 308 bush. 293 bush. 347 bush.	1½ cwts. 2000 lbs. 2150 lbs. 3390 lbs.	30% cwis. 3775 lbs. 4040 lbs. 5619 lbs.	87½ cwts. 60% bush, 43% bush, 46% bush.	3½ cwts, 3940 lbs. 3280 lbs. 4697 lbs.	904 cwts. 7391 lbs. 5990 lbs. 7626 lbs.
5тн Course, 1864-67.										
1864 1865 1806 1867	Swedish Turnips Barley Beans Wheat	8½ cwts. 39 bush. 10½ bush. 21 bush.	04 cwts, 2154 lbs, 1013 lbs, 2143 lbs.	0½ cwts. 4182 lbs. 1629 lbs. 3473 lbs.	68 cwts. 334 bush. 78 bush. 194 bush.	44 cwts, 1615 lbs. 978 lbs. 1966 lbs.	724 cwts, 3394 lbs. 1463 lbs. 3222 lbs.	1764 cwts. 47½ bush. 20½ bush. 23½ bush.	84 cwts, 2595 lbs, 1990 lbs, 3003 lbs,	185 cwts. 5148 lbs. 3343 lbs. 4567 lbs.
Summary—Average of the 5 Courses, 1848-1867.										
1848, '59, '56, '64 } 1849, '56, '66, '64 } '57, '61, '65 } 1850, '54 } '57, '62, '66 } 1851, '55, '59, '63, '67 }	Swedish Turnips  Barley (Clover,1850 (calcd.as hay) (Beans Wheat	26% cwts. 41% bush. 12% bush. 33 bush.	10½ cwts. 2538 lbs. 1252 lbs. 3338 lbs.	374 cwts. 4872 lbs, 54 cwts. 2078 lbs, 5467 lbs,	1364 cwts. 304 bush. 128 bush. 304 bush.	28 cwts. 1815 Ibs. 1347 lbs. 3236 lbs.	164½ cwts. 3529 lbs. 57¼ cwts. 2161 lbs, 5200 lbs.	242½ cwts.  44½ bush.  21½ bush.  35½ bush.	42½ cwts. 2732 lbs. 2036 lbs. 3961 lbs.	285 cwts. 5275 lbs. 63 cwts. 3439 lbs. 6244 lbs.

<sup>(</sup>t) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1-7); Second Course—150 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, and Sixth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-Cake; Second Course—300 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 180 lbs. Sulphate of December 120 lbs. Sulphate of Magnesia, 180 lbs. Sulphate of December 120 lbs. Sulphate of Magnesia 180 lbs. Sulphate of Magnesia 180 lbs. Sulphate of Ammonia 180 lbs. Muriate Acid. 180 lbs. Sulphate of Ammonia 180 lbs. Muriate. 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate

of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, and Sixth Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Maguesia, 200 lbs. Boneash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

<sup>(3)</sup> The quantitles given in Bushels represent the Dressed Corn only.

(4) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, and Total