
Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# **Experiments on the Economy of Nitrogenous Manures**

## **Rothamsted Research**

Rothamsted Research (1873) *Experiments on the Economy of Nitrogenous Manures ;* Memoranda Of The Field Experiments At Rothamsted, May 1872, pp 11 - 11 - **DOI:** https://doi.org/10.23637/ERADOC-1-236

### (11)

#### EXPERIMENTS WITH A VIEW TO ECONOMY IN THE USE OF EXPENSIVE NITROGENOUS MANURES.

IT is found that generally less than half the nitro- commence a series of experiments to determine whether salts, or nitrate of soda, is recovered in the increase of the crop for which they are used; that a considerable quantity may remain in the soil in a comparatively harrowing-in. inactive state, yielding increase very slowly; and that a considerable quantity may be carried away by drainage, and lost. It seemed desirable, therefore, to

gen supplied in such manures as guano, ammonia- any saving can be effected by applying comparatively

Plot No.			PRODUCE PER ACRE.		
	Manures per Acre, &c.	Dressed Corn.			
		Quantity.	Weight per Bushel.	Total Straw,	
1	Unmanured. Seed 1 bushel, dibbled 6 inches apart in the rows	Bushels, 23 <sup>3</sup> / <sub>4</sub>	1bs. 59 · 3	cwts. 24 <del>1</del>	
2	146 lbs, (1) Sulphate Ammonia. Seed 1 bushel;	31 <u>1</u>	59.1	36 <u>1</u>	
3	(292 lbs. Sulphate Ammonia. Seed I bushel;	$28\frac{3}{4}$	58.3	35§	

FIRST SEASON, 1871.

Experiments upon Wheat. Little Hoos Field. Plots 1 acre each.

(1) Containing Nitrogen equal to that in 15 bushels of grain, with its average proportion of Straw.

Experiments upon	Barley.	Thirty-acres Field.	Plots ½ acre each.
------------------	---------	---------------------	--------------------

Plot, No,		PRODUCE PER ACRE.		
			Dressed Corn.	
	MANURES PER ACRE, &c.	Quantity.	Weight per Bushel.	Total Straw.
1	Unmanured. Seed 3 bushels; drilled	Bushels.	1bs. 53 · 9	cwts. 245
2	{1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; } Manures mixed with Ashes and sown broadcast; seed drilled }	49 <del>7</del>	53.3	30 <del>1</del>
3	(1 ewt. Superphosphate, 1 ewt. Nitrate Soda. Seed 3 bushels;	49 <u>1</u>	53 <sup>.</sup> 4	28 <u>1</u>
4	{1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; [Manures, Ashes, and Seed mixed, and drilled together	51	53·0	303
5	{1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 14 bushel; Holes dibbled, 6 inches apart in the rows; Manures (mixed with Ashes) put in, and Seed above }	51 <u>‡</u>	53.3	28 <del>1</del>
	2 cwts. Superphosphate, 2 cwts. Nitrate Soda. Seed 3 bushels; Manures mixed with Ashes and sown broadcast; seed drilled	56‡	51.6	322

#### SECOND SEASON, 1872.

Experiments upon Barley. Thirty-acres Field. Plots 1/2 acre each.

PLOT.	MANURES PER AGRE, &c.	Dressed Corn.		
No.		Quantity.	Weight per Bushel,	Total Straw.
1	Unmanured. Seed 2 <sup>1</sup> / <sub>2</sub> bushels, drilled	Bushels,	lbs.	cwts.
2	(3 cwts. Superphosphate, 2 cwts. Nitrate Soda. Seed 21 bushels;	-		
3	3 ewis. Superphosphate, 2 ewis. Nitrate Soda. Seed 2½ bushels; The Superphosphate mixed with 40 lbs. slaked Lime to neutralize the acid, the Nitrate added, and the whole made up to 15 bushels per acre with Ashes, and sown broadcast; Seed drilled			
4	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 2 <sup>1</sup> / <sub>2</sub> bushels; Manures and Seed made up to 15 bushels per acre with Ashes, and the whole (Manure, Seed, and Ashes) drilled together			
5	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 24 bushels; Manures and Seed made up to 15 bushels per acre with a mixture of half Lime and half Ashes, and the whole (Manure, Seed, Lime, and Ashes) drilled together		-	