Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Memoranda of the Field Experiments at Rothamsted, May 1872



Full Table of Content

Experiments on the Economy of Nitrogenous Manures

Rothamsted Research

Rothamsted Research (1873) *Experiments on the Economy of Nitrogenous Manures*; Memoranda Of The Field Experiments At Rothamsted, May 1872, pp 11 - 11 - **DOI:**

https://doi.org/10.23637/ERADOC-1-236

(11)

EXPERIMENTS WITH A VIEW TO ECONOMY IN THE USE OF EXPENSIVE NITROGENOUS MANURES.

It is found that generally less than half the nitro- | commence a series of experiments to determine whether gen supplied in such manures as guano, ammonia- any saving can be effected by applying comparatively salts, or nitrate of soda, is recovered in the increase small quantities near to the seed, instead of larger of the crop for which they are used; that a considerable quantity may remain in the soil in a comparatively harrowing-in. inactive state, yielding increase very slowly; and that a considerable quantity may be carried away by drainage, and lost. It seemed desirable, therefore, to

amounts in the usual mode of broadcast sowing and

FIRST SEASON, 1871. Experiments upon Wheat. Little Hoos Field. Plots 1 acre each.

PLOT No.	Manures per Acre, &c.	PRODUCE PER ACRE.		
		Dressed Corn.		
		Quantity.	Weight per Bushel.	Total Straw.
1	Unmanured. Seed I bushel, dibbled 6 inches apart in the rows	Bushels,	lbs. 59 · 3	cwts. 24½
2	146 lbs. (1) Sulphate Ammonia. Seed 1 bushel;	31½	59.1	361
3	(292 lbs. Sulphate Ammonia, Seed I bushel;	$28\frac{3}{4}$	58.3	$35\frac{5}{8}$

⁽¹⁾ Containing Nitrogen equal to that in 15 bushels of grain, with its average proportion of Straw.

Experiments upon Barley. Thirty-acres Field. Plots 1/2 acre each.

			Produce per Acre.		
PLOT.	Manures per Acre, &c.	Dressed Corn.			
		Quantity.	Weight per Bushel.	Total Straw.	
1	Unmanured. Seed 3 bushels; drilled	Bushels.	1bs, 53·9	cwts. 245	
2	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; Manures mixed with Ashes and sown broadcast; seed drilled	497	53.3	30½	
3	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels;	49½	53.4	281	
4	{1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 3 bushels; Manures, Ashes, and Seed mixed, and drilled together	51	53.0	30g	
5	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 1½ bushel;	51½	53.3	281	
6	(2 cwts. Superphosphate, 2 cwts. Nitrate Soda. Seed 3 bushels; Manures mixed with Ashes and sown broadcast; seed drilled	56‡	51.6	327	

SECOND SEASON, 1872.

Experiments upon Barley. Thirty-acres Field. Plots ½ acre each.

PLOT. No.	Manures per Acre, &c.	PRODUCE PER ACRE.		
		Dressed Corn.		
		Quantity.	Weight per Bushel,	Total Straw.
1	Unmanured. Seed 2½ bushels, drilled	Bushels,	lbs.	cwts.
2	(3 cwts. Superphosphate, 2 cwts. Nitrate Soda. Seed 2½ bushels;			100
3	3 cwts. Superphosphate, 2 cwts. Nitrate Soda. Seed 2½ bushels; The Superphosphate mixed with 40 lbs. slaked Lime to neutralize the acid, the Nitrate added, and the whole made up to 15 bushels per acre with Ashes, and sown broadcast; Seed drilled			
4	(1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed $2\frac{1}{2}$ bushels; Manures and Seed made up to 15 bushels per acre with Ashes, and the whole (Manure, Seed, and Ashes) drilled together			
5	1 cwt. Superphosphate, 1 cwt. Nitrate Soda. Seed 24 bushels; Manures and Seed made up to 15 bushels per acre with a mixture of half Lime and half Ashes, and the whole (Manure, Seed, Lime, and Ashes) drilled together			