


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Field Experiments at Rothamsted, May 1872



[Full Table of Content](#)

Experiments on Permanent Meadow Land; the Park

Rothamsted Research

Rothamsted Research (1873) *Experiments on Permanent Meadow Land; the Park* ; Memoranda Of The Field Experiments At Rothamsted, May 1872, pp 2 - 2 - DOI:
<https://doi.org/10.23637/ERADOC-1-236>

THE PARK.
EXPERIMENTS WITH DIFFERENT MANURES ON PERMANENT MEADOW LAND.

The Land has probably been laid down with Grass for some centuries. No fresh seed has been artificially sown within the last 40 years certainly; nor is there record of any having been sown since the Grass was first laid down. The experiments commenced in 1856, at which time the character of the herbage appeared uniform over all the Plots. Excepting as explained in the Table, and in the foot-notes, the same description of Manure has been applied year after year to the same Plot.

PLOTS.	Manures, per acre, per Annum.				Produce per Acre, weighed as Hay.				Average per Annum; 16 Years 1856-1871.	PLOTS.
	13th Season; 1868.	14th Season; 1869.	15th Season; 1870.	16th Season; 1871.	Cwts.	Cwts.	Cwts.	Cwts.		
1	(1856-63, 8 years, 14 tons Farnyard Manure, and 200 lbs. Ammonia-salts ⁽¹⁾ ; average produce 49½ cwts. } (1864 and since, 200 lbs. Ammonia-salts alone; average produce (8 years, 1864-71) 43½ cwts. }	41½	61	16½	43½	46½	1			
2	(1856-63, 8 years, 14 tons Farnyard Manure; average produce 42½ cwts. (1864 and since, unmanured; average produce (8 years, 1864-71) 38½ cwts. }	36½	55½	13½	33½	40½	2			
3	Unmanured, continuously	17½	38	5½	25½	22½	3			
4	3½ cwts. Superphosphate of Lime ⁽²⁾	19½	40½	7½	24½	24½	1 } 4			
4	3½ cwts. Superphosphate of Lime, and 400 lbs. Ammonia-salts	29½	45½	8½	38½	36½				
5	400 lbs. Ammonia-salts	24	35½	5½	29½	28½	5			
6	(1856-63, 13 years, 400 lbs. Ammonia-salts; average produce 30½ cwts. (1869 and since, 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph. ; av. prod. (3 yrs., 1869-71) 36½ cwts. }	27½	56½	16½	37½	31½	6			
7	300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	38	54½	17½	39½	35½	7			
8	(1856-61, 6 years, 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate; average produce 36 cwts. (1862 and since, 250 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate; average produce (10 years, 1862-71) 39 cwts. }	27½	46½	12½	30	32½	8			
9	300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, 3½ cwts. Superphosphate, and 400 lbs. Ammonia-salts	59½	68½	29½	58½	52½	9			
10	(1856-61, 6 yrs, 300 lbs. Sulph. Potass, 200 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph., 400 lbs. Amm.-salts; av. prod. 55½ cwts. (1862 and since, 250 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph., 400 lbs. Amm.-salts; av. prod. (10 yrs., 1862-71) 45½ cwts. }	44½	57½	21½	46½	49½	10			
11	(300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph., 800 lbs. Ammonia-salts (300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph., 800 lbs. Ammonia-salts, and 400 lbs. Silicate Soda ⁽³⁾	63½	75½	42½	56½	60½	1 } 11			
11	Unmanured continuously	72½	78½	49	65½	64½				
12	Unmanured continuously	23½	38½	11½	26½	25½	12			
13	300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosph., 400 lbs. Ammonia salts, 2000 lbs. Cut Wheat-straw	61	77½	48	63	56½	13			
14	550 lbs. Nitrate of Soda ⁽⁴⁾ , 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	69	76½	56½	61½	57½	14			
15	550 lbs. Nitrate of Soda	31½	53½	15½	38½	36½	15			
16	275 lbs. Nitrate of Soda, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	51½	74½	33½	57	48½	16			
17	275 lbs. Nitrate of Soda	28½	54½	19½	38½	36½	17			
18	Mixture supplying the quantity of Potass, Soda, Lime, Magnesia, Phosphoric acid, Silica, and Nitrogen, contained in 1 ton of Hay (commencing 1865)	27½	53½	14½	37½	33½	18			
19	275 lbs. Nitrate of Soda, 290 lbs. Sulphate of Potass, and 3½ cwts. Superphosphate (commencing 1872)	19			
20	327 lbs. Nitrate of Potass, and 3½ cwts. Superphosphate (commencing 1872)	20			

(1) "Ammonia-salts"—in all cases equal parts Sulphate and Muriate of Ammonin of Commerce.
 (2) The "Superphosphate of Lime," is, in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid Sp. gr. 1.7 (and water).
 (3) Plots 6, 8, and 10, had, besides the Manures specified, 2000 lbs. Sawdust per acre per annum for the first 7 years, 1856-1862, but without effect.
 (4) 200 lbs. 1856-63 inclusive.
 (5) 500 lbs. in 1862 and 1863.
 (6) Only 400 lbs. in 1859-60-61.
 (7) The application of Silicates did not commence until 1862.
 (8) 550 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 400 lbs. of "Ammonia-salts,"
 (9) Average of 13 years only, as the manures specified were first applied in 1859 (previously, 1856-7 and 8, Sawdust only).
 (10) Average of 14 years only, as these experiments did not commence until 1858.
 (11) Average of 7 years only, as the experiment only commenced in 1865.