Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



## Memoranda of the Field Experiments at Rothamsted, May 1872



Full Table of Content

## **Experiments on Rotation; Agdell Field**

## **Rothamsted Research**

Rothamsted Research (1873) *Experiments on Rotation; Agdell Field;* Memoranda Of The Field Experiments At Rothamsted, May 1872, pp 9 - 9 - **DOI:** https://doi.org/10.23637/ERADOC-1-236

(9)

## AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION-TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present crop (1872) is the 25th experimental one, or the first crop of the Seventh Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in

In the Second, Third, Fourth, Fifth, and Sixth Courses, instead of clover, half of each plot was sown with beans, and the other half left fallow.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land. The abstract of results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the later courses, beans (not fallow) replaced the clover.

(Area under experiment, about 2½ acres.)

	1 lb. (pound avoir, 1 cwt. (hundredwe		= (about) = (about)	1·12 Kilogi 125·5 Kilogi	amme per Hec ammes per He	tare, or 0.57 ctare, or 0.64	Zollverein Pfu Centner per P	nd. per Prussia r. Morgen.	m Morgen.		
	Produce per Acre.										
Years.	Description of Crop.	Prov 1. Unmanured continuously.			PLOT 2. Superphosphate of Lime (1), alone, for the Turnip Crops only.			PLOT 3.  Complex Manure (®), for the Turnip Crops only.			
		Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4).	Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4),	Corn (3) (or Roots).	Straw (or Leaf).	Total Produce (4).	
1st Course, 1848-51.											
1848 1849 1850 1851	Norfolk White Turnips Barley Clover (calcd. as hay) Wheat.	65½ cwts. 44% bush. 28½ bush.	45% cwts. 2983 lbs. 3431 lbs.	1114 cwts. 5656 lbs. 54 cwts. 5389 lbs.	2254 cwts, 297 bush, 28 bush.	106½ cwts. 2111 lbs. 3371 lbs.	332 cwts. 3841 lbs. 574 cwts. 5253 lbs.	218 cwts, 287 bush. 287 bush.	151% cwts. 2088 lbs. 3552 lbs.	369\(\frac{1}{2}\) cwts, 379\(\frac{1}{2}\) lbs, 63\(\cdot \cdot \	
2nd Course, 1852-55.											
1952 1853 1854 1855	Swedish Turnips, Barley Beans, Wheat	26 cwts, 34% bush, 5% bush, 35% bush,	4½ cwts. 2430 lbs. 1055 lbs. 3619 lbs.	304 cwts, 4465 lbs, 1445 lbs, 5859 lbs,	223½ cwts. 28½ bush. 5½ bush. 35½ bush.	204 cwts. 1873 lbs. 1103 lbs. 3525 lbs.	243½ cwts. 3560 lbs. 1524 lbs. 5789 lbs.	396½ cwta. 38½ bush. 9½ bush. 37% bush.	36½ cwts, 2604 lbs, 1355 lbs, 3942 lbs.	433 cwts, 4873 lbs. 2065 lbs. 6371 lbs.	
3rd Course, 1856-59.											
1856 1857 1858 1859	Swedish Turnips Barley Beans Wheat	32 cwts. 48½ bush. 6½ bush. 35½ bush.	2½ cwts, 2600 lbs. 1100 lbs. 4030 lbs.	34½ cwts. 5337 lbs. 1515 lbs. 6262 lbs.	136 cwts, 28½ bush, 6½ bush, 34½ bush.	7½ cwts, 1475 lbs, 1155 lbs. 3930 lbs.	143½ cwts. 3076 lbs. 1605 lbs. 6120 lbs,	3334 cwts. 48 bush. 128 bush. 395 bush.	12½ cwts, 2435 lbs, 1520 lbs, 4610 lbs,	3464 cwts, 5163 lbs. 2357 lbs, 7154 lbs.	
4TH COURSE, 1860-63.											
1860 1861 1862 1863	Swedish Turnips, Barley, Beans, Wheat	1 cwt. 38% bush. 29 bush. 44% bush.	(64 lbs.) 2522 lbs. 1840 lbs. 3467 lbs.	1 cwt. 4718 lbs. 3661 lbs. 6350 lbs.	29½ cwts. 30½ bush. 29¾ bush. 34% bush.	1½ cwts. 2000 lbs, 2150 lbs. 3590 lbs.	30% cwts, 3775 lbs, 4040 lbs, 5619 lbs.	87½ cwts. 60% bush. 43% bush. 46% bush.	34 cwts, 3940 lbs. 3270 lbs. 4697 lbs.	904 cwts. 7394 lbs. 5990 lbs. 7626 lbs.	
5тн Course, 1864-67.											
1864 1865 1866 1867	Swedish Turnips Barley Beans Wheat	8½ cwts. 39 bush. 10¼ bush. 21 bush.	0% cwts. 2154 lbs. 1013 lbs. 2143 lbs.	9½ cwts. 4182 lbs. 1689 lbs. 3473 lbs.	68 cwts. 334 bush. 78 bush. 194 bush.	4½ cwts. 1615 lbs. 978 lbs. 1966 lbs.	723 cwts. 3394 lbs. 1463 lbs. 3222 lbs.	1764 cwts. 47½ bush. 20¾ bush. 23¾ bush.	84 cwts, 2595 lbs, 1990 lbs, 3003 lbs,	185 cwts. 5148 lbs. 3343 lbs. 4567 lbs.	
6TH COURSE, 1863-71.											
1868 1869 1870 1871	Swedish Turnips Barley Beans Wheat	Faile 24& bush, 13& bush, 20& bush.	ed, and ploughed 1948 lbs. 738 lbs. 2799 lbs.	up.   3358 lbs.   1591 lbs.   4092 lbs.	Falle 284 bush. 158 bush. 232 bush.	d, and ploughed 20:5 lbs. 768 lbs. 3048 lbs.	up.  3686 lbs.  1778 lbs.  4521 lbs.	Fail 427 bush. 248 bush. 23 bush.	ed, and ploughed 3309 lbs. 1056 lbs. 3440 lbs.	1 up.   5800 lbs.   2664 lbs.   4883 lbs.	
Summary—Average of the 6 Courses, 1848-1871.											
1848, '52, '56, \\ '60, '64 \\ 1849, '53, '57, \\ 1850, '54 '58, \\ '62, '66, '70 \\ 1851, '55, '59, \\ '63, '67, '71 \\	Swedish Turnips Barley (Clover, 1850 (calcd as hay) Beans Wheat	26% cwts. 38% bush. 12% bush. 30% bush.	10½ cwts. 2440 lbs. 1149 lbs. 3248 lbs.	37‡ cwts, 4619 lbs. 54 cwts, 1980 lbs. 5238 lbs.	136½ cwts. 30 bush. 13 bush. 29¾ bush.	28 cwts. 1850 lbs. 1231 lbs. 3205 lbs.	164½ cwts. 3555 lbs. 57% cwts. 2084 lbs. 5087 lbs.	242½ cwts. 44½ bush. 22½ bush. 33½ bush.	42½ cwts. 2829 lbs. 1840 lbs. 3874 lbs.	285 cwts, 5362 lbs, 63 cwts, 3284 lbs, 6017 lbs,	

<sup>(1)</sup> First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1-7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, and Seventh Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphuric A 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate

of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, and Seventh Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

(3) The quantities given in Buskels represent the Dressed Corn only.

<sup>(4)</sup> The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, and Total