Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Memoranda of the Field Experiments at Rothamsted May 1874



Full Table of Content

Experiments on Oats; Geescroft Field

Rothamsted Research

Rothamsted Research (1875) Experiments on Oats; Geescroft Field; Memoranda Of The Field Experiments At Rothamsted May 1874, pp 5 - 5 - DOI: https://doi.org/10.23637/ERADOC-1-238

GEESCROFT FIELD.

Previous Cropping—1847 and 1848, Clover, Experimental Manures; 1849—1859, Beans, Experimental Manures; 1860, Fallow; 1861 and 1862, Wheat, Unmanured; 1866, Fallow; 1864, Beans, Dunged; 1865, Wheat, Unmanured; 1866, Beans, Unmanured; 1866, Mananured; 1866, Beans, Dunged; 1865, Wheat, Unmanured; 1866, Beans, Unmanured; 1867, Wheat, Unmanured; 1868, Wheat, Unmanured; 1868, Wheat, Unmanured; 1868, Wheat, Unmanured; 1869, Wh EXPERIMENTS ON THE GROWTH OF OATS YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

First Experimental Oat Crop in 1869.

(Area under Experiment, 4 acre.)

(5,)									
	NNUM 1873.	Total Straw.		cwts. 103	138	284	411	273	35
Produce per Acre.	AVERAGE PER ANNUM 5 YEARS, 1869-1873.	Dressed Corn.	Weight per Bushel.	1bs.	35	352	87	853	352
			Quantity.	Bushele.	243	47	59	471	573
	5TH SEASON, 1873.	Total Straw.		cwts.	80 800	163	278	163	24
		Dressed Corn.	Weight per Bushel.	1bs. 271	288888888888888888888888888888888888888	325	343	301	33%
	5тв	Dressec	Quantity.	Bushels.	17	363	484	393	638
	4TH SEASON, 1872.	Total Straw.		cwts.	103	308	458	208	24
		Corn.	Weight per Bushel.	1bs. 36½	373	373	394	362	374
	4тн S	Dressed Corn.	Quantity.	Bushels.	193	555	623	421	448
ODUCE P	3RD SEASON, 1871.	Total Straw.		cwts. 114	133	408	20	343	80 80 80
PB		Dressed Corn.	Weight per Bushel.	1bs. 383	857	363	3558	368	53 84
			Quantity.	Bushels.	22	571	58	55	\$09
	2nd Season, 1870.	Total Straw.		cwta.		174	22 88 88	23	23 88 88
		Corn.	Weight per Bushel.	1bs.	351	343	36	35‡	55 84
	2nd S	Dressed Corn.	Quantity.	Busbels.	191	30	508	361	20
	1sr Season, 1869.	Total Straw.		cwts. 194	243	367	54	423	497
		Corn.	Weight per Bushel,	1bs. 363	383	873	\$6g	383	381
	18T S	Dressed Corn.	Quantity.	Bushels. 363	45	563	75}	624	69
MANURES, PER ACRE, PER ANNUM.				Unmanured	(200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts.) Superphosphate of Lime (1)	400 lbs. Ammonia-salts (2)	400 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	550 lbs. Nitrate of Soda (3)	(550 lbs. Nitrate of Soda, 200 lbs. Sulphate Pokass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate
	Prore	-		, =	64	ec:	4	2	9

6TH SEASON, 1874.

same as in previous years, Only 200 lbs. instead of 400 lbs. Ammonia-salts to Plots 3 and 4, and only 275 lbs. instead of 550 lbs., Nitrate of Soda to Plots 5 and 6; in all other respects the manures are the and as stated above.

(¹) "Superphosphate of Lime"—in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water), (²) "Ammonia-salls"—in each case, equal parts Sulphate and Muriate of Ammonia of Commerce.
(³) 550 lbs. Nitrate of Soda is rediconed to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts,"