


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Field Experiments at Rothamsted May 1874



[Full Table of Content](#)

Experiments on Rotation; Agdell Field

Rothamsted Research

Rothamsted Research (1875) *Experiments on Rotation; Agdell Field* ; Memoranda Of The Field Experiments At Rothamsted May 1874, pp 9 - 9 - DOI: <https://doi.org/10.23637/ERADOC-1-238>

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present crop (1874) is the 27th experimental one, or the third crop of the Seventh Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in the foot-note, No. 2.

In the Second, Third, Fourth, Fifth, and Sixth Courses, instead of clover, half of each plot was sown with beans, and the other half left fallow; for the third crop of the Seventh Course clover was again sown (spring 1873), on half of each plot, the other half being left fallow.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land.

The abstract of the results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the second, third, fourth, fifth, and sixth courses, beans (not fallow) replaced the clover.

(Area under experiment, about 2½ acres.)

1 lb. (pound avoird.) per acre .. = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.								
		Plot 1. Unmanured continuously.			Plot 2. Superphosphate of Lime, ¹ alone, for the Turnip Crops only.			Plot 3. Complex Manure, ² for the Turnip Crops only.		
		Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴
1ST COURSE, 1848-51.										
1848	Norfolk White Turnips	65½ cwt.	45½ cwt.	111½ cwt.	225½ cwt.	106½ cwt.	332 cwt.	218 cwt.	151½ cwt.	369½ cwt.
1849	Barley	44½ bush.	2983 lbs.	5656 lbs.	29½ bush.	2111 lbs.	3841 lbs.	26½ bush.	2088 lbs.	3794 lbs.
1850	Clover (calc. as hay)	54 cwt.	5½ cwt.	63 cwt.
1851	Wheat	28½ bush.	3431 lbs.	5389 lbs.	28 bush.	3371 lbs.	5253 lbs.	28½ bush.	3553 lbs.	5500 lbs.
2ND COURSE, 1852-55.										
1852	Swedish Turnips	26 cwt.	44 cwt.	304 cwt.	223 cwt.	204 cwt.	243½ cwt.	397½ cwt.	361 cwt.	433 cwt.
1853	Barley	343 bush.	2430 lbs.	4465 lbs.	284 bush.	1873 lbs.	3560 lbs.	38½ bush.	2804 lbs.	4573 lbs.
1854	Beans	5½ bush.	1055 lbs.	1445 lbs.	5½ bush.	1103 lbs.	1534 lbs.	3½ bush.	1355 lbs.	2065 lbs.
1855	Wheat	35½ bush.	3619 lbs.	5839 lbs.	35½ bush.	3595 lbs.	5789 lbs.	37½ bush.	3942 lbs.	6371 lbs.
3RD COURSE, 1856-59.										
1856	Swedish Turnips	32 cwt.	2½ cwt.	34½ cwt.	136 cwt.	7½ cwt.	143½ cwt.	333½ cwt.	12½ cwt.	346½ cwt.
1857	Barley	48½ bush.	2600 lbs.	5337 lbs.	284 bush.	1475 lbs.	3076 lbs.	48 bush.	2435 lbs.	5163 lbs.
1858	Beans	6½ bush.	1100 lbs.	1515 lbs.	6½ bush.	1185 lbs.	1605 lbs.	12½ bush.	1320 lbs.	2357 lbs.
1859	Wheat	35½ bush.	4030 lbs.	6262 lbs.	34½ bush.	3890 lbs.	6120 lbs.	39½ bush.	4610 lbs.	7154 lbs.
4TH COURSE, 1860-63.										
1860	Swedish Turnips	1 cwt.	(6½ lbs.)	1 cwt.	294 cwt.	1½ cwt.	304 cwt.	87½ cwt.	3½ cwt.	904 cwt.
1861	Barley	32½ bush.	2322 lbs.	4718 lbs.	30½ bush.	2000 lbs.	3775 lbs.	60½ bush.	3940 lbs.	7391 lbs.
1862	Beans	29 bush.	1840 lbs.	3661 lbs.	29½ bush.	2150 lbs.	4040 lbs.	43½ bush.	3280 lbs.	5990 lbs.
1863	Wheat	44½ bush.	3437 lbs.	6330 lbs.	34½ bush.	3890 lbs.	5119 lbs.	46½ bush.	4597 lbs.	7626 lbs.
5TH COURSE, 1864-67.										
1864	Swedish Turnips	8½ cwt.	6½ cwt.	94 cwt.	68 cwt.	44 cwt.	72½ cwt.	176½ cwt.	8½ cwt.	185 cwt.
1865	Barley	39 bush.	2174 lbs.	4182 lbs.	334 bush.	1615 lbs.	3394 lbs.	47½ bush.	2535 lbs.	5148 lbs.
1866	Beans	10½ bush.	1013 lbs.	1629 lbs.	7½ bush.	978 lbs.	1463 lbs.	20½ bush.	1990 lbs.	3343 lbs.
1867	Wheat	21 bush.	2143 lbs.	3473 lbs.	19½ bush.	1966 lbs.	3222 lbs.	22½ bush.	3003 lbs.	4567 lbs.
6TH COURSE, 1868-71.										
1868	Swedish Turnips	Failed, and ploughed up.			Failed, and ploughed up.			Failed, and ploughed up.		
1869	Barley	24½ bush.	1948 lbs.	3358 lbs.	28½ bush.	2025 lbs.	3686 lbs.	42½ bush.	3309 lbs.	5800 lbs.
1870	Beans	13½ bush.	738 lbs.	1591 lbs.	15½ bush.	768 lbs.	1778 lbs.	24½ bush.	1056 lbs.	2664 lbs.
1871	Wheat	20½ bush.	2793 lbs.	4092 lbs.	23½ bush.	3048 lbs.	4521 lbs.	23 bush.	3440 lbs.	4883 lbs.
7TH COURSE, 1872-75.										
1872	Swedish Turnips	34½ cwt.	8½ cwt.	42½ cwt.	170½ cwt.	17½ cwt.	188 cwt.	339½ cwt.	85½ cwt.	375½ cwt.
1873	Barley	28½ bush.	1343 lbs.	2717 lbs.	20½ bush.	1565 lbs.	2875 lbs.	31½ bush.	1723 lbs.	3573 lbs.
1874	Clover
1875	Wheat

SUMMARY—AVERAGE OF THE FIRST 6 COURSES, 1848-1871.

1848, '52, '56, '60, '64	Swedish Turnips	26½ cwt.	10½ cwt.	37½ cwt.	136½ cwt.	28 cwt.	164½ cwt.	242½ cwt.	4½ cwt.	255 cwt.
1849, '53, '57, '61, '65, '69	Barley	38½ bush.	2440 lbs.	4619 lbs.	20 bush.	1850 lbs.	3553 lbs.	44½ bush.	2829 lbs.	5362 lbs.
1850, '54, '58, '62, '66, '70	Clover, 1850 (calc. as hay)	54 cwt.	57½ cwt.	63 cwt.
1851, '55, '59, '63, '67, '71	Beans	12½ bush.	1149 lbs.	1930 lbs.	13 bush.	1231 lbs.	2084 lbs.	22½ bush.	1340 lbs.	3244 lbs.
	Wheat	30½ bush.	3248 lbs.	6238 lbs.	29½ bush.	3205 lbs.	5087 lbs.	33½ bush.	3874 lbs.	6017 lbs.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, and Seventh Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-Cake; Second Course—300 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, and Seventh Courses—320 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

(3) The quantities given in Bushels represent the Dressed Corn only.

(4) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, and Total Straw.