


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1875

[Full Table of Content](#)



Experiments on Rotation; Agdell Field

Rothamsted Research

Rothamsted Research (1876) *Experiments on Rotation; Agdell Field* ; Yields Of The Field Experiments 1875, pp 9 - 9 - DOI: <https://doi.org/10.23637/ERADOC-1-239>

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present crop (1875) is the 28th experimental one, or the fourth crop of the Seventh Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in the foot-note, No. 2.

In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead, on half of each plot, and the other half left fallow; for the third crop of the Seventh Course clover was again sown (spring 1873), on half of each plot, the other half being left fallow.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land.

The abstract of the results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the second, third, fourth, fifth, and sixth courses, beans (not fallow) replaced the clover.

(Area under experiment, about 2½ acres.)

1 lb. (pound avoird.) per acre .. = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.								
		PLOT 1. Unmanured continuously.			PLOT 2. Superphosphate of Lime, alone, for the Turnip Crops only.			PLOT 3. Complex Manure, ² for the Turnip Crops only.		
		Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴
1ST COURSE, 1848-51.										
1848	Norfolk White Turnips	65½ cwt.	45½ cwt.	111½ cwt.	22½ cwt.	10½ cwt.	33½ cwt.	218 cwt.	15¼ cwt.	369½ cwt.
1849	Barley	44½ bush.	2983 lbs.	5656 lbs.	29½ bush.	2111 lbs.	3841 lbs.	28½ bush.	2083 lbs.	3791 lbs.
1850	Clover (calc ^d . as hay)	54 cwt.	5½ cwt.	63 cwt.
1851	Wheat	28½ bush.	3481 lbs.	5389 lbs.	28 bush.	3371 lbs.	5233 lbs.	28½ bush.	3552 lbs.	5500 lbs.
2ND COURSE, 1852-55.										
1852	Swedish Turnips	26 cwt.	4 cwt.	30½ cwt.	22½ cwt.	20½ cwt.	24½ cwt.	39½ cwt.	36½ cwt.	433 cwt.
1853	Barley	34½ bush.	2430 lbs.	4463 lbs.	28½ bush.	1873 lbs.	3560 lbs.	38½ bush.	2604 lbs.	4873 lbs.
1854	Beans	5½ bush.	1055 lbs.	1443 lbs.	5½ bush.	1103 lbs.	1534 lbs.	9½ bush.	1355 lbs.	2065 lbs.
1855	Wheat	35½ bush.	3619 lbs.	5859 lbs.	35½ bush.	3523 lbs.	5789 lbs.	37½ bush.	3942 lbs.	6371 lbs.
3RD COURSE, 1856-59.										
1856	Swedish Turnips	32 cwt.	2½ cwt.	34½ cwt.	136 cwt.	7½ cwt.	14½ cwt.	33½ cwt.	12½ cwt.	346½ cwt.
1857	Barley	48½ bush.	2600 lbs.	5337 lbs.	28½ bush.	1475 lbs.	3076 lbs.	44 bush.	2455 lbs.	5163 lbs.
1858	Beans	64 bush.	1100 lbs.	1513 lbs.	6½ bush.	1153 lbs.	1605 lbs.	12½ bush.	1320 lbs.	2357 lbs.
1859	Wheat	35½ bush.	4030 lbs.	6282 lbs.	34½ bush.	3030 lbs.	6120 lbs.	29½ bush.	4610 lbs.	7154 lbs.
4TH COURSE, 1860-63.										
1860	Swedish Turnips	1 cwt.	6½ lb.	1 cwt.	29½ cwt.	14 cwt.	30½ cwt.	87½ cwt.	34 cwt.	90½ cwt.
1861	Barley	38½ bush.	2522 lbs.	4712 lbs.	30½ bush.	2000 lbs.	3775 lbs.	60½ bush.	3930 lbs.	7391 lbs.
1862	Beans	29 bush.	1840 lbs.	3661 lbs.	29½ bush.	2150 lbs.	4040 lbs.	4½ bush.	3280 lbs.	5990 lbs.
1863	Wheat	44½ bush.	3467 lbs.	6350 lbs.	34½ bush.	3390 lbs.	5619 lbs.	46½ bush.	4697 lbs.	7636 lbs.
5TH COURSE, 1864-67.										
1864	Swedish Turnips	8½ cwt.	0½ cwt.	9 cwt.	68 cwt.	4½ cwt.	72½ cwt.	176½ cwt.	84 cwt.	185 cwt.
1865	Barley	39 bush.	2154 lbs.	4182 lbs.	33½ bush.	1615 lbs.	3394 lbs.	47½ bush.	2595 lbs.	5148 lbs.
1866	Beans	10½ bush.	1013 lbs.	1689 lbs.	7½ bush.	978 lbs.	1463 lbs.	20½ bush.	1990 lbs.	3343 lbs.
1867	Wheat	21 bush.	2143 lbs.	3473 lbs.	19½ bush.	1866 lbs.	3222 lbs.	23½ bush.	3003 lbs.	4567 lbs.
6TH COURSE, 1868-71.										
1868	Swedish Turnips	Failed, and ploughed up.			Failed, and ploughed up.			Failed, and ploughed up.		
1869	Barley	24½ bush.	1949 lbs.	3358 lbs.	28½ bush.	2023 lbs.	3686 lbs.	42½ bush.	3309 lbs.	5900 lbs.
1870	Beans	13½ bush.	738 lbs.	1591 lbs.	15½ bush.	768 lbs.	1778 lbs.	24½ bush.	1036 lbs.	2664 lbs.
1871	Wheat	20½ bush.	2799 lbs.	4092 lbs.	20½ bush.	3048 lbs.	4521 lbs.	23 bush.	3440 lbs.	4853 lbs.
7TH COURSE, 1872-75.										
1872	Swedish Turnips	34½ cwt.	8½ cwt.	42½ cwt.	170½ cwt.	17½ cwt.	183 cwt.	339½ cwt.	35½ cwt.	375½ cwt.
1873	Barley	22½ bush.	1343 lbs.	2717 lbs.	20½ bush.	1563 lbs.	2873 lbs.	31½ bush.	1723 lbs.	3573 lbs.
1874	Clover	31½ cwt.	52½ cwt.	84½ cwt.
1875	Wheat

SUMMARY—AVERAGE OF THE FIRST 6 COURSES, 1848-1871.

1848, '52, '56, '60, '64	Swedish Turnips	26½ cwt.	10½ cwt.	37½ cwt.	136½ cwt.	28 cwt.	164½ cwt.	242½ cwt.	42½ cwt.	285 cwt.
1849, '53, '57, '61, '65, '69	Barley	38½ bush.	2440 lbs.	4619 lbs.	30 bush.	1850 lbs.	3555 lbs.	44½ bush.	2329 lbs.	5362 lbs.
1850, '54, '58, '62, '66, '70	Clover, 1850 (calc ^d . as hay)	54 cwt.	57½ cwt.	63 cwt.
1851, '55, '59, '63, '67, '71	Beans	12½ bush.	1149 lbs.	1990 lbs.	13 bush.	1231 lbs.	2084 lbs.	22½ bush.	1840 lbs.	3284 lbs.
	Wheat	30½ bush.	3248 lbs.	5233 lbs.	29½ bush.	3205 lbs.	5087 lbs.	33½ bush.	3874 lbs.	6017 lbs.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, and Seventh Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-Cake; Second Course—300 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate

of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, and Seventh Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

(3) The quantities given in *Bushels* represent the *Dressed* Corn only.

(4) The "Total Produce" of the Corn-crops includes *Dressed* Corn, *Offal* Corn, and *Total* Straw.