

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1876

[Full Table of Content](#)



## Experiments on Rotation; Agdell Field

### Rothamsted Research

Rothamsted Research (1877) *Experiments on Rotation; Agdell Field* ; Yields Of The Field Experiments 1876, pp 10 - 10 - DOI: <https://doi.org/10.23637/ERADOC-1-240>

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present crop (1876) is the 29th experimental one, or the first crop of the Eighth Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in the foot-note, No. 2.

In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead, on half of each plot, and the other half left fallow; for the third crop of the Seventh Course clover was again sown (spring 1873), on half of each plot, the other half being left fallow.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land.

The abstract of the results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the second, third, fourth, fifth, and sixth courses, beans (not fallow) replaced the clover.

(Area under experiment, about 2½ acres.)

1 lb. (pound avoird.) per acre .. = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.  
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.								
		Plot 1. Unmanured continuously.			Plot 2. Superphosphate of Lime, alone, for the Turnip Crops only.			Plot 3. Complex Manure <sup>2</sup> for the Turnip Crops only.		
		Corn <sup>3</sup> (or Roots).	Straw (or Leaf).	Total Produce. <sup>4</sup>	Corn <sup>3</sup> (or Roots).	Straw (or Leaf).	Total Produce. <sup>4</sup>	Corn <sup>3</sup> (or Roots).	Straw (or Leaf).	Total Produce. <sup>4</sup>
1ST COURSE, 1848-51.										
1848	Norfolk White Turnips	6½ cwt.	45½ cwt.	111½ cwt.	22½ cwt.	106½ cwt.	332 cwt.	218 cwt.	151½ cwt.	369½ cwt.
1849	Barley	4½ bush.	293½ lbs.	563½ lbs.	2½ bush.	211½ lbs.	384½ lbs.	28½ bush.	2088½ lbs.	3794½ lbs.
1850	Clover (calc. as hay)	..	..	54 cwt.	..	..	5½ cwt.	..	..	63 cwt.
1851	Wheat	28½ bush.	343½ lbs.	588½ lbs.	29 bush.	337½ lbs.	523½ lbs.	28½ bush.	3552½ lbs.	5500½ lbs.
2ND COURSE, 1852-55.										
1852	Swedish Turnips	26 cwt.	44 cwt.	304 cwt.	22½ cwt.	204 cwt.	243½ cwt.	398½ cwt.	364 cwt.	433 cwt.
1853	Barley	34½ bush.	2430 lbs.	4465 lbs.	28½ bush.	1873½ lbs.	5350½ lbs.	38½ bush.	2604½ lbs.	4573½ lbs.
1854	Beans	5½ bush.	1065 lbs.	1445 lbs.	5½ bush.	1103½ lbs.	1334½ lbs.	9½ bush.	1355½ lbs.	2065½ lbs.
1855	Wheat	35½ bush.	3619 lbs.	5959 lbs.	33½ bush.	3325½ lbs.	5789½ lbs.	37½ bush.	3942½ lbs.	6371½ lbs.
3RD COURSE, 1856-59.										
1856	Swedish Turnips	32 cwt.	24 cwt.	344 cwt.	136 cwt.	74 cwt.	1424 cwt.	233½ cwt.	124 cwt.	346½ cwt.
1857	Barley	48½ bush.	2600 lbs.	5337 lbs.	28½ bush.	1475½ lbs.	3076½ lbs.	48 bush.	2435½ lbs.	5168½ lbs.
1858	Beans	6½ bush.	1100 lbs.	1515 lbs.	6½ bush.	1155½ lbs.	1605½ lbs.	12½ bush.	1320½ lbs.	2357½ lbs.
1859	Wheat	35½ bush.	4030 lbs.	6262 lbs.	34½ bush.	3930½ lbs.	6120½ lbs.	39½ bush.	4610½ lbs.	7154½ lbs.
4TH COURSE, 1860-63.										
1860	Swedish Turnips	1 cwt.	(64 lbs.)	1 cwt.	20½ cwt.	14 cwt.	30½ cwt.	87½ cwt.	34 cwt.	90½ cwt.
1861	Barley	38½ bush.	2522 lbs.	4719 lbs.	30½ bush.	2000 lbs.	3775 lbs.	60½ bush.	3940 lbs.	7391 lbs.
1862	Beans	29 bush.	1840 lbs.	3661 lbs.	29½ bush.	2159 lbs.	4040 lbs.	43½ bush.	3280 lbs.	5990 lbs.
1863	Wheat	44½ bush.	3467 lbs.	6350 lbs.	34½ bush.	3390 lbs.	5619 lbs.	46½ bush.	4687 lbs.	7626 lbs.
5TH COURSE, 1864-67.										
1864	Swedish Turnips	8½ cwt.	6½ cwt.	94 cwt.	68 cwt.	43 cwt.	72½ cwt.	1764 cwt.	8½ cwt.	185 cwt.
1865	Barley	39 bush.	2154 lbs.	4182 lbs.	33½ bush.	1615½ lbs.	3394½ lbs.	47½ bush.	2565½ lbs.	5148½ lbs.
1866	Beans	10½ bush.	1013 lbs.	1689 lbs.	7½ bush.	973½ lbs.	1463½ lbs.	20½ bush.	1900½ lbs.	3343½ lbs.
1867	Wheat	21 bush.	2143 lbs.	3473 lbs.	19½ bush.	1966½ lbs.	3222½ lbs.	23½ bush.	3003½ lbs.	4567½ lbs.
6TH COURSE, 1868-71.										
1868	Swedish Turnips	Failed, and ploughed up.			Failed, and ploughed up.			Failed, and ploughed up.		
1869	Barley	24½ bush.	1948 lbs.	3338 lbs.	28½ bush.	2045½ lbs.	3586½ lbs.	42½ bush.	3509½ lbs.	5800½ lbs.
1870	Beans	13½ bush.	738 lbs.	1591 lbs.	13½ bush.	768½ lbs.	1778½ lbs.	24½ bush.	1036½ lbs.	2664½ lbs.
1871	Wheat	20½ bush.	2739 lbs.	4092 lbs.	23½ bush.	3043½ lbs.	4521½ lbs.	23 bush.	3440½ lbs.	4883½ lbs.
7TH COURSE, 1872-75.										
1872	Swedish Turnips	34½ cwt.	8½ cwt.	42½ cwt.	170½ cwt.	112 cwt.	188 cwt.	359½ cwt.	35½ cwt.	375½ cwt.
1873	Barley	23½ bush.	1343 lbs.	2717 lbs.	20½ bush.	1563½ lbs.	2875½ lbs.	31½ bush.	1723½ lbs.	3573½ lbs.
1874	Clover	..	..	31½ cwt.	..	..	52½ cwt.	..	..	84½ cwt.
1875	Wheat	21½ bush.	2430 lbs.	3784 lbs.	28½ bush.	3536½ lbs.	5328½ lbs.	31½ bush.	4685½ lbs.	6699½ lbs.

SUMMARY—AVERAGE OF THE FIRST 7 COURSES, 1848-1875.

1848, '52, '56, '60, '64, '72	} Swedish Turnips	27½ cwt.	10½ cwt.	384 cwt.	142½ cwt.	26½ cwt.	168½ cwt.	258½ cwt.	41½ cwt.	300 cwt.
1849, '53, '57, '61, '65, '69, '73		} Barley	36½ bush.	2283 lbs.	4349 lbs.	28½ bush.	1809 lbs.	3458 lbs.	42½ bush.	2671 lbs.
1850, '54, '58, '62, '66, '70, '74	} (Clover, 1850 and '74) { Clover (calc. as hay)		..	..	42½ cwt.	..	..	55 cwt.	..	..
1851, '55, '59, '63, '67, '71, '75		} Beans	12½ bush.	1149 lbs.	1990 lbs.	13 bush.	1231 lbs.	2084 lbs.	22½ bush.	1340 lbs.
	} Wheat		30 bush.	3131 lbs.	5030 lbs.	29½ bush.	3252 lbs.	5122 lbs.	33 bush.	3990 lbs.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, and Seventh Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-Cake; Second Course—300 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lb. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate

of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, and Seventh Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

(3) The quantities given in Bushels represent the Dressed Corn only.

(4) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, and Total Straw.