


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Field Experiments at Rothamsted: May 1877



[Full Table of Content](#)

Experiments on Permanent Meadow Land; the Park

Rothamsted Research

Rothamsted Research (1878) *Experiments on Permanent Meadow Land; the Park* ; Memoranda Of The Field Experiments At Rothamsted: May 1877, pp 2 - 2 - DOI:
<https://doi.org/10.23637/ERADOC-1-241>

THE PARK.

EXPERIMENTS WITH DIFFERENT MANURES ON PERMANENT MEADOW LAND.

The Land has probably been laid down with Grass for some centuries. No fresh seed has been artificially sown within the last 40 years certainly; nor is there record of any having been sown since the Grass was first laid down. The experiments commenced in 1856, at which time the character of the herbage appeared uniform over all the Plots. Excepting as explained in the Table and in the foot-notes, the same description of Manure has been applied year after year to the same Plot.

(Area under experiment, about 7 acres.)

PLOTS.	Manures, per acre, per Annum.	PRODUCE PER ACRE, WEIGHED AS HAY.						PLOTS.
		Average per Annum.			Twenty-fourth Season, 1875 (18).			
		10 Years, 1866-66, (C)	20 Years, 1866-76, (C)	Cwts.	First Crop.	Second Crop.	Total.	
1	{1856-63, 8 years, 14 tons Farmyard Manure, and 200 lbs. Ammonia-salts; average produce 49½ cwts. } {1864 and since, 200 lbs. Ammonia-salts alone; average produce (12 years, 1864-75) 38½ cwts. }	48½	37½	33½	33½	17½	51½	1
2	{1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. } {1864 and since, unmanured; average produce (12 years, 1864-75) 32½ cwts. }	41½	32	26½	26½	11½	38½	2
3	Unmanured, continuously	22½	20	21½	21½	20	19½	3
4	3½ cwts. Superphosphate of Lime (1)	23½	21½	22½	22½	21	15½	16½
5	3½ cwts. Superphosphate of Lime, and 400 lbs. Ammonia-salts	33½	30½	32½	32½	18	51	16½
6	400 lbs. Ammonia-salts	30½	22	26½	26½	24½	18	17½
7	{1856-63, 13 years, 400 lbs. Ammonia-salts; average produce 30½ cwts. } {1869 and since, 300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	31½	30½	30½	30½	35½	15	50½
8	{1856-61, 6 years, 300 lbs. Sulph. Potass, 200 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	33½	36½	35½	36½	40½	24	64½
9	{1862 and since, 250 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	38½	26½	30½	26½	16	44½	24½
10	300 lbs. Sulphate Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 400 lbs. Ammonia-salts	53½	48½	51	48½	52	24½	76½
11	{1856-61, 6 yrs. 300 lbs. Sulph. Potass, 200 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate; average produce 36 cwts. } {1862 and since, 250 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate; average produce (14 years, 1862-75) 27½ cwts. }	52½	39½	46	39½	43	24½	67½
12	300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	61½	58	46	57½	46	50	97½
13	300 lbs. Sulph. Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	63½	61½	62½	62½	60	41	101
14	Unmanured continuously	25	22½	24	24	25	14½	37½
15	300 lbs. Sulphate Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	55½	59	57½	59	65	30	95½
16	550 lbs. Nitrate of Soda (2), 800 lbs. Sulphate Potass, 100 lbs. Sulph. Soda, 100 lbs. Sulph. Magnesia, and 3½ cwts. Superphosphate	53½	60½	57	60½	62½	17	86½
17	275 lbs. Nitrate of Soda, 300 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	36	35	38	35	29½	13	42½
18	Mixture supplying the quantity of Potass, Soda, Lime, Magnesia, Phosphoric acid, Silica, and Nitrogen, contained in 1 ton of Hay (commencing 1865)	45½	47	46	47	45	16	63½
19	275 lbs. Nitrate of Soda, 290 lbs. Sulphate of Potass, and 3½ cwts. Superphosphate (commencing 1872)	34½	33	33	33	30	13	43
20	327 lbs. Nitrate of Potass, and 3½ cwts. Superphosphate (commencing 1872)	21	33	32	33	34	15	50
		38½	38½	41	20	61
		36½	36½	42	21	63

(1) "Ammonia-salts" - in all cases equal parts Sulphate and Nitrate of Ammonia of Commerce.
 (2) The "Superphosphate of Lime" is, in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid Sp. Gr. 1.7 (and water).
 (3) Plots 6, 8, and 10, laid, besides the Manures specified, 2000 lbs. Sawdust per acre per annum for the first 7 years, 1856-1862, but without effect.
 (4) 200 lbs. 1856-63 inclusive.
 (5) 500 lbs. in 1862 and 1863.
 (6) Only 400 lbs. in 1859-60-61.
 (7) The application of Silicates did not commence until 1862.
 (8) 550 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 400 lbs. of "Ammonia-salts."
 (9) The manures specified were first applied in 1859 (previously, 1856-7 and 8, Sawdust only).
 (10) Averages of 8 years, 10 years, and 18 years, as these experiments did not commence until 1858.
 (11) Averages of 4 years only, 1872-75.
 (12) In previous years the second crop has either been fed off by sheep, without other food, or mown and left on the ground; but in the twentieth season, 1875, it was so unusually heavy, that it was cut, weighed as hay, and removed.
 (13) The second crop of the twentieth season (1875) is not included in these averages, as in all other years the first crop only was weighed and removed.