


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1878

[Full Table of Content](#)



Experiments on Oats; Geescroft Field

Rothamsted Research

Rothamsted Research (1879) *Experiments on Oats; Geescroft Field* ; Yields Of The Field Experiments 1878, pp 11 - 11 - DOI: <https://doi.org/10.23637/ERADOC-1-242>

GEESCROFT FIELD.

EXPERIMENTS ON THE GROWTH OF OATS YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1847 and 1848, Clover, Experimental Manures; 1849—1869, Beans, Experimental Manures; 1860, Fallow; 1861 and 1862, Wheat, Unmanured; 1863, Fallow; 1864, Beans, Dunged; 1865, Wheat, Unmanured; 1866, Beans, Unmanured; 1867 and 1868, Wheat, Unmanured.
First Experimental Oat Crop in 1869.

(Area under Experiment, $\frac{1}{2}$ acre.)

PLOTS.	MANURES, PER ACRE, PER ANNUM.	PRODUCE PER ACRE.											
		1ST SEASON, 1869.		2ND SEASON, 1870.		3RD SEASON, 1871.		4TH SEASON, 1872.		5TH SEASON, 1873.		AVERAGE PER ANNUM 5 YEARS, 1869-1873.	
		Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.
1	Unmanured	36 $\frac{1}{2}$	36 $\frac{1}{2}$	16 $\frac{1}{2}$	35	20 $\frac{1}{2}$	33 $\frac{1}{2}$	15	36 $\frac{1}{2}$	10 $\frac{1}{2}$	27 $\frac{1}{2}$	19 $\frac{1}{2}$	33 $\frac{1}{2}$
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate of Lime (1)	45	38 $\frac{1}{2}$	19 $\frac{1}{2}$	35 $\frac{1}{2}$	22	35 $\frac{1}{2}$	19 $\frac{1}{2}$	37 $\frac{1}{2}$	10 $\frac{1}{2}$	28 $\frac{1}{2}$	24 $\frac{1}{2}$	35
3	400 lbs. Ammonia-salts (2)	56 $\frac{1}{2}$	37 $\frac{1}{2}$	30	34 $\frac{1}{2}$	57 $\frac{1}{2}$	36 $\frac{1}{2}$	55 $\frac{1}{2}$	37 $\frac{1}{2}$	30 $\frac{1}{2}$	39 $\frac{1}{2}$	47	35 $\frac{1}{2}$
4	{ 400 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate	75 $\frac{1}{2}$	39 $\frac{1}{2}$	50 $\frac{1}{2}$	36	58 $\frac{1}{2}$	35 $\frac{1}{2}$	62 $\frac{1}{2}$	39 $\frac{1}{2}$	45 $\frac{1}{2}$	34 $\frac{1}{2}$	59	37
5	550 lbs. Nitrate of Soda (3)	62 $\frac{1}{2}$	38 $\frac{1}{2}$	38 $\frac{1}{2}$	35 $\frac{1}{2}$	55	36 $\frac{1}{2}$	42 $\frac{1}{2}$	36 $\frac{1}{2}$	20 $\frac{1}{2}$	30 $\frac{1}{2}$	47 $\frac{1}{2}$	35 $\frac{1}{2}$
6	{ 550 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate	69 $\frac{1}{2}$	38 $\frac{1}{2}$	50	35 $\frac{1}{2}$	60 $\frac{1}{2}$	33 $\frac{1}{2}$	44 $\frac{1}{2}$	37 $\frac{1}{2}$	24	38 $\frac{1}{2}$	57 $\frac{1}{2}$	35 $\frac{1}{2}$
SECOND 5 YEARS; MINERAL MANURES AS BEFORE, AMMONIA-SALTS AND NITRATE OF SODA ONLY HALF AS MUCH AS PREVIOUSLY.													
		6TH SEASON, 1874.		7TH SEASON, 1875.		8TH SEASON, 1876 (4).		9TH SEASON, 1877 (5).		10TH SEASON, 1878.		AVERAGE PER ANNUM 5 YEARS, 1874-1878.	
		Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.	Quantity, Bushels.	Weight per Bushel, lbs.
1	Unmanured	12	31 $\frac{1}{2}$	12 $\frac{1}{2}$	29 $\frac{1}{2}$	8 $\frac{1}{2}$	32
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate of Lime (1)	13 $\frac{1}{2}$	31 $\frac{1}{2}$	13 $\frac{1}{2}$	29 $\frac{1}{2}$	7 $\frac{1}{2}$	30
3	200 lbs. Ammonia-salts (2)	37 $\frac{1}{2}$	33 $\frac{1}{2}$	30 $\frac{1}{2}$	32 $\frac{1}{2}$	17 $\frac{1}{2}$	34 $\frac{1}{2}$
4	{ 200 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate	46 $\frac{1}{2}$	34 $\frac{1}{2}$	30 $\frac{1}{2}$	34 $\frac{1}{2}$	29 $\frac{1}{2}$	35 $\frac{1}{2}$
5	275 lbs. Nitrate of Soda (3)	35 $\frac{1}{2}$	30 (4)	23 $\frac{1}{2}$ (4)	31 $\frac{1}{2}$ (4)	12 $\frac{1}{2}$	30 $\frac{1}{2}$
6	{ 275 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwt. Superphosphate	28 $\frac{1}{2}$ (4)	33 $\frac{1}{2}$ (4)	28 $\frac{1}{2}$ (4)	33 $\frac{1}{2}$ (4)	19 $\frac{1}{2}$	33 $\frac{1}{2}$

(1) "Superphosphate of Lime"—in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water).

(2) "Ammonia-salts"—in each case, equal parts Sulphate and Muriate of Ammonia of Commerce.

(3) 550 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."

(4) On these plots, where large quantities of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got into favourable condition for sowing, and the plant was very irregular. Owing to the extremely wet condition of the land, especially on the Nitrate plots, it was not sown until April 6, and then with a very unfavourable seed bed; and there being a heavy fall of snow a week later, the plant came up very irregularly, and much of it perished from standing surface-water.

(5) Owing to the very wet winter, 1876-7, the land could not be worked in time for sowing, and was therefore left fallow in 1877.