Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Memoranda of the Field Experiments at Rothamsted: May 1879



Full Table of Content

Experiments on Mangold-wurzel; Barn-field

Rothamsted Research

Rothamsted Research (1880) *Experiments on Mangold-wurzel; Barn-field;* Memoranda Of The Field Experiments At Rothamsted: May 1879, pp 18 - 19 - **DOI:**

https://doi.org/10.23637/ERADOC-1-243

(18)

EXPERIMENTS ON MANGOLD WURZEL.—BARN FIELD (after Sugar-Beet); commencing 1876.

The arrangement of the Plots is precisely the same as previously for Sugar-beet, excepting that Plot 9, which was unmanured for Sugar-beet, and also previously for Swedes, is now added as a manured Plot. With this exception, the manures are also substantially the same as previously for Sugar-beet; in fact, precisely the same as for the Sugar-beet in 1872 and 1873. Seed, Yellow Globe; dibbled on ridges, rows 26 inches apart; plants 11 inches apart in the rows (*). Area under experiment about 8 acres. Roots all carted off; Leaves weighed, spread on the respective Plots, and ploughed in.

1											
		MANURE	S PER ACR	E PER AN	NUM.						
PLOTS.	Series 1.	As S	IES 2. eries 1, dressed with litrate Soda.	As Se and Cross- 400 lbs.	ries 3, eries 1, dressed with Ammonia- ts."	As Se and Cross- 2000 lbs and 400	eries 1, dressed with Rape-cake lbs. "Am- a-salts."	SERIES 5. As Series 1, and Cross-dressed with 2000 lbs, Rape-cake.			
	First Season, 1876.	Seed dibbl	ed, May 25	2–26. Cro	op taken up	, Nov. 3-1	.7,	4 2	6.2	- 1	
7						PRODUCE	PER ACRE.	7 1		-	
		Roots.	Leaves.	Roots.	Leaves.	Roots.	Leaves.	Roots.	Leaves.	Roots.	Leaves.
1 2 3 4 5 6 7 8 9	Farmyard Manure (14 tons). Farmyard Manure (14 tons), and 3½ cwts. Superphosphate (¹) Without Manure (1846, and since) (3½ cwts. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride) Sodium (common salt), 200 lbs. Sulphate Magnesia 3½ cwts. Superphosphate 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphos, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate (²)	Tons. cwts. 19 12 19 13 6 10 8 8 7 10 6 16 8 13 5 9	Tons. cwts. 4 9 4 6 1 14 1 15 1 14 1 12 2 3 1 10	Tons ewts. 25 2 27 13 20 13 25 1 21 0 21 2 22 11 15 16	Tons. cwts. 7 5 7 3 5 12 6 0 5 14 5 8 5 14 5 3	Tons. cwts 29 19 29 8 14 3 19 19 13 10 17 15 19 2 11 17 25 14	Tons. cwts. 7 12 7 10 4 10 4 9 5 1 4 13 5 11 4 16 7 6	Tons. cwts. 31 9 30 18 19 19 30 8 17 2 26 8 27 2 18 2	Tons, cwts, 10 5 9 16 7 7 8 13 7 14 9 0 9 9 7 11	Tons. cwts. 24 9 29 19 17 4 25 8 17 17 20 10 20 12 15 12	Tons. cwts. 5 19 6 12 4 15 5 10 5 17 5 4 5 15 4 18
	Second Season, 1877. Seed dibbled	June 4-6	(Plots 8 a	nd 9, Jun	e 11th). (Crop taken	up, Nov. 1	4-23.	T 14 9	L B	11.00
1 2 3 4 5 6 7 8 9	Farmyard Manure (14 tons) Farmyard Manure (14 tons) and 3½ cwts. Superphosphate (¹) Without Manure (1846, and since) (3½ cwts. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride) Sodium (common salt), 200 lbs. Sulphate Magnesia 3½ cwts. Superphosphate 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphos, 500 lbs. Sulphate Potass, 36½ lbs. Amsalts (²) Urmanured, 1853, and since; previously part Umman, part Superphos. Farmyard Manure (14 tons), 3½ cwts. Superphosphate (³)	Tons. cwts. 15 7 16 14 5 9 6 16 6 1 5 8 7 0 3 19	Tons. cwts. 2 1 1 19 1 0 1 3 0 19 0 18 1 3 1 3	Tons. cwts. 24 13 26 8 16 17 21 10 20 5 20 19 22 2 9 17	Tons, cwts. 3 14 3 12 3 14 3 10 3 1 2 18 3 16 5 4	Tons. cwts. 27 1 26 18 8 16 16 10 12 2 15 6 16 13 7 4 13 17	Tons, cwts. 4 4 4 6 3 0 2 2 2 10 1 16 2 7 3 10 4 0	Tons, ewis. 30 5 28 15 13 9 27 9 15 3 24 18 25 15 11 9	Tons. cwts. 5 5 5 9 3 19 3 8 3 8 3 16 5 0 4 11	Tons. cwts. 25 18 24 12 13 17 21 14 15 3 19 3 20 13 10 3	Tons. cwts. 3 4 2 19 2 10 1 17 2 2 1 12 2 8 3 3
	THIRD SEASON, 1878. Seed dibb	led, June 8	3-9 (Plot 9	, June 11	th). Crop	taken up,	Nov. 7–20				
5 6 7 8	Farmyard Manure (14 tons) Farmyard Manure (14 tons), and 3½ cwts. Superphosphate (¹) Without Manure (1846, and since) (3½ cwts. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride) Sodium (common salt), 200 lbs. Sulphate Magnesia 3½ cwts. Superphosphate 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphos, 500 lbs. Sulphate Potass 3½ cwts. Superphos, 500 lbs. Sulphate Potass 3½ cwts. Superphos, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate (²) Unmanured, 1853, and since; previously part Unman, part Superphos. Farmyard Manure (14 tons), 3½ cwts. Superphosphate (²)	Tons. cwts. 13 5 14 16 3 10 5 9 4 14 3 18 5 8 2 13	Tons. cwts. 2 16 2 19 1 4 1 7 1 8 1 3 1 9 1 4	Tons. cwts. 18 15 21 4 10 2 18 10 14 11 15 1 13 18 11 19	Tons. ewts. 4 4 4 15 2 16 4 6 3 18 3 7 3 1 4 7	Tons. cwts. 20 11 19 15 4 7 14 3 8 2 12 0 11 18 6 13 15 17	Tons. ewts. 5 6 5 3 2 11 2 12 3 6 2 14 2 18 3 5 5 9	Tons, cwts. 22 4 20 18 6 11 21 2 8 4 15 3 14 0 6 12	Tons. cwts. 6 3 5 17 3 7 4 14 3 3 4 11 4 5 4 10	Tons cwts 17 1 18 17 6 3 15 19 8 1 12 5 11 19 6 4	Tons. cwts. 3 13 3 15 2 17 3 2 3 6 3 3 3 8 3 5
	Fourth S.	EASON, 187	9. Seed	dibbled, M	ay 13-15.						
5 6 7 8	Farmyard Manure (14 tons), and 3½ cwts. Superphosphate (¹) Without Manure (1846, and since) (3) cwts. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride) Sodium (common salt), 200 lbs. Sulphate Magnesia 3½ cwts. Superphosphate. 3½ cwts. Superphosphate. 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass, 36½ lbs. Am-salts (²) Unmanured, 1853, and since; previously part Unman., part Superphos. Farmyard Manure (14 tons), 3½ cwts. Superphosphate (³)	Tons, cwts.	Tons. cwts.	Tons, cwts.	Tons, cwts.	Tons, ewts.	Tons. cwts.	Tons, cwts.	Tons. cwts.	Tons, cwts.	Tons. cwts.
		Fifth	SEASON,	1880.							
1 2 3 4 5 6 7 8 9	Farmyard Manure (14 tons) Farmyard Manure (14 tons), and 3½ cwts. Superphosphate (¹) Without Manure (1846, and since) (3½ cwts. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride) Sodium (common salt), 200 lbs. Sulphate Magnesia 3½ cwts. Superphosphate 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate, 500 lbs. Sulphate Potass 3½ cwts. Superphosphate Unmanured, 1853, and since; previously part Unman., part Superphos. Farmyard Manure (14 tons), 3½ cwts. Superphosphate (³)	Tons, cwts.	Tons, cwts.	Tons. cwts.	Tons. cwts.	Tons. cwts.	Tons, cwts.	Tons. cwts.	Tons, cwts.	Tons. cwts.	Tons, cwts.
64	 (¹) "Superphosphate of Lime"—in all cases made fror (²) "Ammonia-salts"—in each case equal parts Sulpho (²) Plot 9 sown on the flat instead of on ridges; plants 	ate and Muriat	e of Ammonia	a of Commerc	e.						

(19)

EXPERIMENTS ON MANGOLD WURZEL.—BARN FIELD—continued.

SUMMARY OF THE COMPOSITION OF THE MANGEL ROOTS.

- As it will be some time before we shall be able to report fully the results obtained, or to be yet obtained, illustrating the influence of different manures, and of different seasons, on the composition of Mangels, an abstract of some of the analytical results, at present at command, is given below. The dry matter, ash, and nitrogen, are of course determined in the roots themselves. The sugar is determined in the expressed juice; and calculated into its percentage in the roots, on the assumption that they contain uniformly 96 per cent. of juice. But, with roots varying so much in character of growth, size, and ripeness, this will not be the case. Nevertheless, the results so calculated, approximately, and usefully, represent both the actual and relative amounts of sugar in the various roots. The amounts of dry matter, ash, and nitrogen, have also, in many cases, been determined in the expressed juice. In many cases also, the amount of the nitrogen existing as albuminoids has been determined. It may be observed that by far the larger proportion of both the mineral matter and the nitrogen of the roots is found in the juice; and of the nitrogen in the juice a variable proportion, ranging from less than one-fifth to not more than one-third of the total, is found to exist as albuminoids.

 In interpreting the figures, it must be borne in mind, that, with forty different experiments each year, and, in each year four, or five, or more, times, as much produce on some plots as on others, it would be impossible to sample each at its best, and all in the same condition of ripeness. Each year the seed was sown on all the plots at the same time. The sample analysed was in each case a mixture of vertical sections of ten or fifteen roots, and all the samples were as a rule taken within a period of from one to two weeks; as far as practicable beginning with the ripest. It is obvious, however, that the smaller crops would be much riper than the larger ones.

For								CROSS-	DRESSED M	ANURES,	PER ACE	RE, PER	ANNUM.							
Manures and Produce, se facing page.	SERIES 1. No cross-dressing.				SERIES 2. As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.				SERIES 3. As Series 1, and Cross-dressed with 400 lbs. Ammonia-salts.				SERIES 4. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake and 400 lbs. Ammonia-salts,				SERIES 5. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake,			
									First S	EASON,	1876.				1					
			1	1	7 1		Cent. To				_	De.N.), and Nitro	_	_	1		-		
PLOTS.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter, Per cent.	Sugar. Per cent.	Ash.	Nitrogen.	Dry Matter. Per cent.	Sugar. Per cent.		Nitrogen. Per cent.	Dry Matter. Per cent.	Sugar. Per cent.	Ash.	Nitrogen. Per cent.	Dry Matter.	Sugar. Per cent.	Ash, Per cent,	Nitrog Per ce
1 2 3	Per cent. 12:14 12:41 15:14	Per cent. 7:14 7:19	0.969 0.943 0.828	rer cent.	10·54 9·35 11·94	4.85	1.031 1.020 0.903	rei cent.	10.65 9.64 12.16	5:72	1.080 1.018 0.90±	Ter cont.	8·98 8·92 11·60		1.065 1.034 0.811	I GI GGM	11·30 10·51 12·42	::	0.989 1.005 0.751	
5 6 7 8 9	13·99 13·51 13·67 13·63 13·06	8·98 9·48 8·74	0.905 0.818 0.928 0.882 0.900		11·36 10·99 11·23 11·61 11·23	6·32 6·36 7·67	1·013 0·917 0·929 0·922 0·945		12·23 11·73 11·02 10·62 11·43 11·59	7·03 7·93 7·41 7·80	0 989 0.735 0.993 0.969 0.905 0.876		9·91 10·93 10·5; 10·66 10·20	5·62 6·05 5·40	1.067 0.816 1.036 1.015 0.856		11·28 10·65 11·55 11·58 11·61	6·94 6·84 7·30	1·003 0·744 0·911 0·936 0·757	
					1 1				SECOND	1				79.5						
1 2 3	Per cent. 14·48 13·85 16·58	Per cent. 9:01 10:02 11:19	Per cent. 0.988 0.961 0.827	Per cent.	Per cent. 12·01 12·91 14·06	Per cent. 8:21 8:22 8:76	Per cent. 1:122 1:107 1:072	Per cent.	Per cent. 12.95 13.24 17.11	Per cent. 8:95 7:84 10:16	1.097 1.089 0.888	Per cent.	Per cent. 12:44 11:78 14:44	7·97 7·68 9·80	1·114 1·126 0·834	Per cent.	Per cent. 13·34 14·08 16·41	Per cent. 7:79 8:51 10:21	Per cent. 1:010 1:000 0:819	Per ce
4 5 6 7 8 9	15·42 15·84 16·15 15·88 16·23	10·92 11·62 11·31	0.948 0.797 0.891 0.943 0.933		12·25 12·90 12·53 12·74 14·01	7·26 8·54 9·10	1·121 0·889 1·135 1·034 1·023	.4-1	13·11 15·63 15·05 13·96 14·95 14·84	9·35 10·00 9·45 	1.085 0.838 1.095 1.098 0.932 1.011		12·69 14·36 14·27 12·58 14·51	7·51 8·24 8·90	1·221 0·786 1·061 1·136 0·811		13·45 15·35 14·10 13·83 14·87	9·81 10·66 9·94	1.046 0.784 0.978 1.036 0.807	
	1 3								THIRD	Season,	1878.									
1 2 3	Per cent, 12·26 11·51 15·25	Per cent. 7-32 6-97 10-20	0.995 0.981 0.824	Per cent. 0:170 0:182 0:186	Per cent. 11:47 19:05 12:02	6.36 5.21 7.08	Per cent. 1:036 1:072 0:908	0·218 0·216 0 211	Per cent. 11:17 11:00 13:47	6·27 6·08 8·09	1.013 1.034 0.811	Per cent. 0·206 0·206 0·261	Per cent. 10.83 10.50 12.86	5·65 5·9± 7·61	1·046 0·987 0·802	Per cent. 0·2±1 0·217 0·247	Per cent. 11:98 10:66 14:10	6·90 6·14 8·82	Per cent. 0.985 0.948 0.846	0·18 0·17 0·29
4 5 6 7 8 9	13·56 13·91 14·23 13·42 14·50	9·01 9·17 9·12	0.928 0.810 0.989 0.976 0.903	0·129 0·144 0·173	11·03 11·61 11·04 11·26 11·10	6·24 6·90 6·23	1.084 0.873 0.986 0.982 0.937	0·188 0·188 0·193	11.90 13.00 13.55 11.92 12.81 10.77	7·27 8·14 8·67 6·21	0·975 0·845 0·988 0·932 0·869 -0·939	0·144 0·187 0·184	10·33 12·69 12·09 12·03 11·93	5·88 7·68 6·96	1·027 0·739 1·016 0·986 0·879	0·181 0·244 0·235	11·22 13·87 12·18 12·05 12·52	6·53 8·66 7·36	0*786 0*940 0*977 0*863	0·17 0·21 0·19
							V		Fourth	Season,	1879.									
1 2 3 4 5 6 7 8 9	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per ceut.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per_cent.	Per cent.	Per cent.	Per ce
3					<u> </u>				Fifth S	SEASON,	1880.									
1	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.			Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per o
1 2 3 4 5 6 7	, 6				•				Ser.	+										
8 9					,									-	-			*		60